

# 9.2.6

Puni Aditya - EE25BTECH11046

## Question:

Area of the region in the first quadrant enclosed by the x-axis, the line  $y = x$  and the circle  $x^2 + y^2 = 32$  is \_\_\_\_\_.

## Solution:

Let the conic section be  $g(\mathbf{x}) = \mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0$ . Let the line be  $\mathbf{x} = \mathbf{h} + \kappa \mathbf{m}$ . To find the points of intersection, we substitute the line equation into the conic equation.

$$g(\mathbf{h} + \kappa \mathbf{m}) = (\mathbf{h} + \kappa \mathbf{m})^T \mathbf{V} (\mathbf{h} + \kappa \mathbf{m}) + 2\mathbf{u}^T (\mathbf{h} + \kappa \mathbf{m}) + f = 0 \quad (1)$$

$$= (\mathbf{h}^T + \kappa \mathbf{m}^T) \mathbf{V} (\mathbf{h} + \kappa \mathbf{m}) + 2\mathbf{u}^T \mathbf{h} + 2\kappa \mathbf{u}^T \mathbf{m} + f = 0 \quad (2)$$

$$= \mathbf{h}^T \mathbf{V} \mathbf{h} + 2\kappa \mathbf{m}^T \mathbf{V} \mathbf{h} + \kappa^2 \mathbf{m}^T \mathbf{V} \mathbf{m} + 2\mathbf{u}^T \mathbf{h} + 2\kappa \mathbf{u}^T \mathbf{m} + f = 0 \quad (3)$$

$$= (\mathbf{m}^T \mathbf{V} \mathbf{m}) \kappa^2 + 2(\mathbf{m}^T \mathbf{V} \mathbf{h} + \mathbf{m}^T \mathbf{u}) \kappa + (\mathbf{h}^T \mathbf{V} \mathbf{h} + 2\mathbf{u}^T \mathbf{h} + f) = 0 \quad (4)$$

$$= (\mathbf{m}^T \mathbf{V} \mathbf{m}) \kappa^2 + 2\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}) \kappa + g(\mathbf{h}) = 0 \quad (5)$$

This is a quadratic equation in  $\kappa$ .

$$\kappa_{1,2} = \frac{-2\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}) \pm \sqrt{4(\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}))^2 - 4(\mathbf{m}^T \mathbf{V} \mathbf{m}) g(\mathbf{h})}}{2\mathbf{m}^T \mathbf{V} \mathbf{m}} \quad (6)$$

$$\kappa_{1,2} = \frac{-\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}) \pm \sqrt{(\mathbf{m}^T (\mathbf{V} \mathbf{h} + \mathbf{u}))^2 - (\mathbf{m}^T \mathbf{V} \mathbf{m}) g(\mathbf{h})}}{\mathbf{m}^T \mathbf{V} \mathbf{m}} \quad (7)$$

Using (7) to find the intersection points that define the boundaries of the area,

$$\text{Circle: } x^2 + y^2 - 32 = 0 \implies \mathbf{V} = \mathbf{I}, \mathbf{u} = \mathbf{0}, f = -32$$

$$\text{Lines: } \mathbf{x} = \kappa \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \text{ so } \mathbf{h}_1 = \mathbf{0}, \mathbf{m}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ and } \mathbf{x} = \kappa \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \text{ so } \mathbf{h}_2 = \mathbf{0}, \mathbf{m}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$g(\mathbf{h}_1) = g(\mathbf{0}) = -32 \quad (8)$$

$$\mathbf{m}_1^T \mathbf{V} \mathbf{m}_1 = \begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{I} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = 1 \quad (9)$$

$$\mathbf{m}_1^T (\mathbf{V} \mathbf{h}_1 + \mathbf{u}) = \begin{pmatrix} 1 & 0 \end{pmatrix} (\mathbf{I} \mathbf{0} + \mathbf{0}) = 0 \quad (10)$$

$$\kappa = \frac{0 \pm \sqrt{0^2 - (1)(-32)}}{1} = \pm \frac{\sqrt{32}}{1} = \pm 4\sqrt{2} \quad (11)$$

$$g(\mathbf{h}_2) = g(\mathbf{0}) = -32 \quad (12)$$

$$\mathbf{m}_2^\top \mathbf{V} \mathbf{m}_2 = \begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{I} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 2 \quad (13)$$

$$\mathbf{m}_2^\top (\mathbf{V} \mathbf{h}_2 + \mathbf{u}) = \begin{pmatrix} 1 & 1 \end{pmatrix} (\mathbf{I} \mathbf{0} + \mathbf{0}) = 0 \quad (14)$$

$$\kappa = \frac{0 \pm \sqrt{0^2 - (2)(-32)}}{2} = \pm \frac{\sqrt{64}}{2} = \pm 4 \quad (15)$$

The intersection points are

$$\mathbf{x}_i = \kappa \mathbf{m}_i \quad (16)$$

In the first quadrant, the intersection points defining the region are:

$$\mathbf{x}_1 = \begin{pmatrix} 4\sqrt{2} \\ 0 \end{pmatrix}, \quad \mathbf{x}_2 = \begin{pmatrix} 4 \\ 4 \end{pmatrix} \quad (17)$$

The area is the sum of two integrals, split at the x-coordinate of  $\mathbf{x}_2$ .


$$A = \int_0^4 x \, dx + \int_4^{4\sqrt{2}} \sqrt{32 - x^2} \, dx \quad (18)$$

$$= \left[ \frac{x^2}{2} \right]_0^4 + \left[ \frac{x}{2} \sqrt{32 - x^2} + 16 \sin^{-1} \left( \frac{x}{4\sqrt{2}} \right) \right]_4^{4\sqrt{2}} \quad (19)$$

$$= \frac{16}{2} + \left[ 0 + 16 \sin^{-1}(1) \right] - \left[ \frac{4}{2} \sqrt{16} + 16 \sin^{-1} \left( \frac{1}{\sqrt{2}} \right) \right] \quad (20)$$

$$= 8 + 16 \left( \frac{\pi}{2} \right) - \left[ 8 + 16 \left( \frac{\pi}{4} \right) \right] \quad (21)$$

$$= 8 + 8\pi - 8 - 4\pi = 4\pi \quad (22)$$



Plot