

Matgeo Presentation - Problem 5.8.19

ee25btech11021 - Dhanush sagar

October 2, 2025

Problem Statement

Let a, b, c be real numbers. Consider the following system of equations in x, y, z :

$$\begin{aligned}\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} &= 1, \\ \frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} &= 1, \\ -\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} &= 1.\end{aligned}$$

The system has:

- 1) no solution
- 2) unique solution
- 3) infinitely many solutions
- 4) finitely many solutions

solution

Let

$$A = \frac{x^2}{a^2}, \quad (0.1)$$

$$B = \frac{y^2}{b^2}, \quad (0.2)$$

$$C = \frac{z^2}{c^2}. \quad (0.3)$$

Then the system becomes

$$A + B - C = 1, \quad (0.4)$$

$$A - B + C = 1, \quad (0.5)$$

$$-A + B + C = 1. \quad (0.6)$$

solution

The augmented matrix is

$$\begin{aligned} &\begin{pmatrix} 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ -1 & 1 & 1 & 1 \end{pmatrix} \xrightarrow{R_2 \rightarrow R_2 - R_1} \begin{pmatrix} 1 & 1 & -1 & 1 \\ 0 & -2 & 2 & 0 \\ -1 & 1 & 1 & 1 \end{pmatrix} \xrightarrow{R_3 \rightarrow R_3 + R_1} \begin{pmatrix} 1 & 1 & -1 & 1 \\ 0 & -2 & 2 & 0 \\ 0 & 2 & 2 & 2 \end{pmatrix} \\ &\xrightarrow{R_3 \rightarrow R_3 + R_2} \begin{pmatrix} 1 & 1 & -1 & 1 \\ 0 & -2 & 2 & 0 \\ 0 & 0 & 2 & 2 \end{pmatrix} \xrightarrow{R_2 \rightarrow -\frac{1}{2}R_2} \begin{pmatrix} 1 & 1 & -1 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 2 & 2 \end{pmatrix} \\ &\xrightarrow{R_3 \rightarrow \frac{1}{2}R_3} \begin{pmatrix} 1 & 1 & -1 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \xrightarrow{R_2 \rightarrow R_2 + R_3} \begin{pmatrix} 1 & 1 & -1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \\ &\xrightarrow{R_1 \rightarrow R_1 + R_3} \begin{pmatrix} 1 & 1 & 0 & 2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \xrightarrow{R_1 \rightarrow R_1 - R_2} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}. \end{aligned}$$

(0.7)

solution

From the final matrix we read

$$A = 1, \quad B = 1, \quad C = 1. \quad (0.8)$$

Therefore,

$$\frac{x^2}{a^2} = 1, \quad \frac{y^2}{b^2} = 1, \quad \frac{z^2}{c^2} = 1, \quad (0.9)$$

which gives

$$x = \pm a, \quad y = \pm b, \quad z = \pm c. \quad (0.10)$$

Hence there are $2^3 = 8$ distinct solutions for (x, y, z) , so the correct choice is

(d) finitely many solutions

(0.11)

C Source Code: fraction matrix.c

```
#include <stdio.h>

void gen_system_points(double *A, double *B) {
    // Coefficient matrix (3x3)
    double tempA[9] = {
        1,  1, -1,    // Eqn 1
        1, -1,  1,    // Eqn 2
        -1, 1,  1     // Eqn 3
    };
    // RHS vector
    double tempB[3] = {1, 1, 1};

    for (int i = 0; i < 9; i++) A[i] = tempA[i];
    for (int i = 0; i < 3; i++) B[i] = tempB[i];
}
```

Python Script: fraction matrix.py

```
import ctypes
import numpy as np
import itertools
# --- Load C shared library ---
lib = ctypes.CDLL("./gen_system_points.so")
lib.gen_system_points.argtypes = [ctypes.POINTER(ctypes.c_double) * 9,
                                   ctypes.POINTER(ctypes.c_double) * 3]
# Storage
A_storage = (ctypes.c_double * 9)()
B_storage = (ctypes.c_double * 3)()
lib.gen_system_points(A_storage, B_storage)
# Convert to numpy
A = np.array(A_storage).reshape(3, 3)
B = np.array(B_storage)
print("Coefficient matrix A:\n", A)
print("RHS vector B:\n", B)
# Solve AX = B
X, Y, Z = np.linalg.solve(A, B)
```

Python Script: fraction matrix.py

```
print("\nSolution (X,Y,Z) = ({}, {}, {})".format(X, Y, Z))
# Symbolic answer
symbolic_points = []
for sx, sy, sz in itertools.product([1, -1], repeat=3):
    symbolic_points.append((f"{'+' if sx>0 else '-'}a",
                           f"{'+' if sy>0 else '-'}b",
                           f"{'+' if sz>0 else '-'}c"))
print("\nSymbolic solutions (in terms of a,b,c):")
for p in symbolic_points:
    print(p)
a, b, c = 2.0, 3.0, 1.0    # change these values
numeric_points = [(sx*a, sy*b, sz*c) for sx, sy, sz in itertools.product([1, -1], repeat=3)]
print("\nNumeric solutions (for a={}, b={}, c={}):".format(a, b, c))
for p in numeric_points:
    print(p)
np.savetxt("points_abc.txt", numeric_points)
print("\nSaved numeric points to points_abc.txt")
```