

Matgeo Presentation - Problem 5.8.19

ee25btech11021 - Dhanush sagar

October 3, 2025

Problem Statement

If we add 1 to the numerator and subtract 1 from the denominator, a fraction reduces to 1. It becomes $\frac{1}{2}$ if we only add 1 to the denominator. What is the fraction?

solution

Let the unknown fraction be represented as

$$\mathbf{u} = \begin{pmatrix} x \\ y \end{pmatrix}, \quad \frac{x}{y}. \quad (0.1)$$

Affine transformations are written as a vector plus a translation.

case 1: add 1 to numerator, subtract 1 from denominator

$$T_1(\mathbf{u}) = \mathbf{u} + \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad (0.2)$$

case 2: add 1 to denominator

$$T_2(\mathbf{u}) = \mathbf{u} + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (0.3)$$

Condition for a fraction: For any vector $\begin{pmatrix} a \\ b \end{pmatrix}$, the requirement $\frac{a}{b} = k$ is equivalent to the linear equation

$$\mathbf{r}_k \begin{pmatrix} a \\ b \end{pmatrix} = 0, \quad \mathbf{r}_k = (1 \quad -k), \quad (0.4)$$

solution

since $\mathbf{r}_k \begin{pmatrix} a \\ b \end{pmatrix} = a - kb = 0 \iff \frac{a}{b} = k$.

Case 1: $T_1(\mathbf{u})$ must yield fraction 1. use $\mathbf{r}_1 = (1 \quad -1)$

$$\mathbf{r}_1(T_1(\mathbf{u})) = 0 \quad (0.5)$$

$$\implies \mathbf{r}_1 \mathbf{u} + \mathbf{r}_1 \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 0 \quad (0.6)$$

$$(1 \quad -1) \mathbf{u} + 2 = 0 \quad (0.7)$$

This gives the first equation.

Case 2: $T_2(\mathbf{u})$ must yield fraction $\frac{1}{2}$. To avoid fractions, multiply the functional by 2, i.e., use $\mathbf{r}_2 = (2 \quad -1)$.

$$\mathbf{r}_2(T_2(\mathbf{u})) = 0 \quad (0.8)$$

$$\implies \mathbf{r}_2 \mathbf{u} + \mathbf{r}_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} = 0 \quad (0.9)$$

$$(2 \quad -1) \mathbf{u} - 1 = 0 \quad (0.10)$$

This gives the second equation.

solution

System of equations: Both conditions together form the system

$$\begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix} \mathbf{u} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}. \quad (0.11)$$

Gaussian elimination: Form the augmented matrix

$$\left[\begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix} \mid \begin{pmatrix} -2 \\ 1 \end{pmatrix} \right]. \quad (0.12)$$

Eliminate the entry below the pivot:

$$R_2 \leftarrow R_2 - 2R_1 \Rightarrow \left[\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \mid \begin{pmatrix} -2 \\ 5 \end{pmatrix} \right]. \quad (0.13)$$

Now eliminate above the pivot:

$$R_1 \leftarrow R_1 + R_2 \Rightarrow \left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \mid \begin{pmatrix} 3 \\ 5 \end{pmatrix} \right]. \quad (0.14)$$

Final solution:

$$\mathbf{u} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}, \frac{x}{y} = \frac{3}{5}. \quad (0.15)$$

C Source Code: fraction matrix.c

```
#include <stdio.h>

// Generate augmented matrix for fraction problem
void generate_matrix(double matrix[2][3]) {
    // Equation 1:  $x - y = -2$ 
    matrix[0][0] = 1;
    matrix[0][1] = -1;
    matrix[0][2] = -2;

    // Equation 2:  $2x - y = 1$ 
    matrix[1][0] = 2;
    matrix[1][1] = -1;
    matrix[1][2] = 1;
}
```

Python Script: fraction_matrix.py

```
import ctypes
import numpy as np

# Load the shared library
lib = ctypes.CDLL("./libfraction_matrix.so")

# Prepare 2x3 matrix
MatrixType = ctypes.c_double * 3 * 2
matrix = MatrixType()

# Call C function to generate matrix
lib.generate_matrix(matrix)
```

Python Script: fraction matrix.py

```
# Convert to numpy array
aug_matrix = np.array([[matrix[i][j] for j in range(3)] for i

# Solve system
A = aug_matrix[:, :2]
B = aug_matrix[:, 2]
solution = np.linalg.solve(A, B)
x, y = solution

print(f"The fraction is {x}/{y}")

# Optionally save fraction to file for plotting
with open("fraction.txt", "w") as f:
    f.write(f"{x} {y}")
```


Python Script: plot matrix.py

```
import numpy as np
import matplotlib.pyplot as plt

# Load fraction solution
with open("fraction.txt", "r") as f:
    x_sol, y_sol = map(float, f.read().split())

# Define the two lines from the equations:
# Equation 1:  $x - y = -2 \rightarrow y = x + 2$ 
# Equation 2:  $2x - y = 1 \rightarrow y = 2x - 1$ 

x_vals = np.linspace(0, 5, 100)
y1 = x_vals + 2
y2 = 2*x_vals - 1
```

Python Script: plot matrix.py

```
# Plot the lines
```

```
plt.plot(x_vals, y1, label="x - y = -2")
```

```
plt.plot(x_vals, y2, label="2x - y = 1")
```

```
# Plot the intersection point
```

```
plt.scatter(x_sol, y_sol, color='red', s=100, label=f'Intersection')
```

```
plt.xlabel("x (Numerator)")
```

```
plt.ylabel("y (Denominator)")
```

```
plt.title("Intersection of Two Lines (Fraction Solution)")
```

```
plt.grid(True)
```

```
plt.legend()
```

```
plt.show()
```

Result Plot

