5.13.15

SAMYAK GONDANE - AI25BTECH11029

Question

Find the equation of the circle passing through (0,0) and making intercepts a and b on the coordinate axes.

Solution

We use the general matrix form of a conic:

$$\mathbf{x}^{\mathsf{T}}\mathbf{V}\mathbf{x} + 2\mathbf{u}^{\mathsf{T}}\mathbf{x} + f = 0 \tag{1}$$

where:

$$\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}, \quad \mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}, \quad f \in \mathbb{R}$$
 (2)

Solution

Substitute the three points:

- At (0,0): f=0
- At (a,0): $a^2 + 2u_1a = 0 \Rightarrow u_1 = -\frac{a}{2}$
- At (0, b): $b^2 + 2u_2b = 0 \Rightarrow u_2 = -\frac{b}{2}$

So,

$$\mathbf{u} = \begin{pmatrix} -\frac{a}{2} \\ -\frac{b}{2} \end{pmatrix}, \quad f = 0 \tag{3}$$

Solution

Substitute back into the general form:

$$\mathbf{x}^{T} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} -\frac{a}{2} & -\frac{b}{2} \end{pmatrix} \mathbf{x} = 0$$
 (4)

Simplify:

$$x^2 + y^2 - ax - by = 0 (5)$$




Figure: