

Question 4.13.5

AI25BTECH11040 - Vivaan Parashar

October 1, 2025

1 Question:

The set of lines $ax + by + c = 0$, where $3a + 2b + 4c = 0$ are concurrent at the point _____.

2 Solution:

We are given the fact that $3a + 2b + 4c = 0$. This can be written as:

$$\implies \begin{pmatrix} 3 & 2 & 4 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = 0 \quad (1)$$

Let the point of concurrency be \mathbf{P} , at coordinates $\begin{pmatrix} p_x \\ p_y \end{pmatrix}$. Because \mathbf{P} lies on all lines $ax + by + c = 0$, we can write the following system of equations:

$$\begin{pmatrix} 3 & 2 & 4 \\ p_x & p_y & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (2)$$

For this system to have a non-trivial solution in a , b and c , and for the two original equations to be linearly dependent (since equation 2 should be true whenever 1 is true), the rank of the coefficient matrix must be 1. Applying row reduction:

$$\begin{pmatrix} 3 & 2 & 4 \\ p_x & p_y & 1 \end{pmatrix} \xrightarrow{R_3 \rightarrow R_3 - \frac{R_1}{4}} \begin{pmatrix} 3 & 2 & 4 \\ p_x - \frac{3}{4} & p_y - \frac{1}{2} & 0 \end{pmatrix} \quad (3)$$

Clearly, for the rank to be 1, the last row must be all zeros. Therefore the point of concurrency \mathbf{P} is $\begin{pmatrix} \frac{3}{4} \\ \frac{1}{2} \end{pmatrix}$.

3 Plot:




Figure 1: Graph of lines with randomly generated values of a and b satisfying $3a + 2b + 4c = 0$. All lines are concurrent at the point $\begin{pmatrix} \frac{3}{4} \\ \frac{1}{2} \end{pmatrix}$ (marked in red).