

10.3.21

EE25BTECH11020 - Darsh Pankaj Gajare

Question:

Find the point at which the line $y = x + 1$ is a tangent to the curve $y^2 = 4x$.

Solution: The given conic can be expressed as

$$\mathbf{x}^T V \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (1)$$

where

$$V = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} -2 \\ 0 \end{pmatrix}, \quad f = 0 \quad (2)$$

and $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$.

The given line is

$$y = x + 1 \quad (3)$$

which can be parameterized as

$$\mathbf{x} = \mathbf{h} + t\mathbf{m} \quad (4)$$

where

$$\mathbf{h} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \mathbf{m} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}. \quad (5)$$

Substituting $\mathbf{x} = \mathbf{h} + t\mathbf{m}$ in the conic equation,

$$(\mathbf{h} + t\mathbf{m})^T V (\mathbf{h} + t\mathbf{m}) + 2\mathbf{u}^T (\mathbf{h} + t\mathbf{m}) + f = 0 \quad (6)$$

Expanding,

$$t^2 (\mathbf{m}^T V \mathbf{m}) + 2t (\mathbf{m}^T V \mathbf{h} + \mathbf{u}^T \mathbf{m}) + (\mathbf{h}^T V \mathbf{h} + 2\mathbf{u}^T \mathbf{h} + f) = 0 \quad (7)$$

Compute each term:

$$\mathbf{m}^T V \mathbf{m} = 1, \quad \mathbf{m}^T V \mathbf{h} = 1, \quad \mathbf{u}^T \mathbf{m} = -2, \quad (8)$$

$$\mathbf{h}^T V \mathbf{h} = 1, \quad \mathbf{u}^T \mathbf{h} = 0 \quad (9)$$

Substituting,

$$t^2 - 2t + 1 = 0 \quad (10)$$


$$\Rightarrow (t - 1)^2 = 0 \Rightarrow t = 1 \quad (11)$$

Hence, the point of contact is

$$\mathbf{q} = \mathbf{h} + t\mathbf{m} \quad (12)$$

$$\mathbf{q} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}. \quad (13)$$

Plot using C libraries:



Plot using Python:

