

## 1.8.22: Equidistant Points Problem

ee25btech11005 - Aditya Mishra

September 9, 2025

### Question

Find all points that are equidistant from the points

$$\mathbf{A} = \begin{pmatrix} -5 \\ 4 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} -1 \\ 6 \end{pmatrix}.$$

How many such points exist?

### Solution

Let the desired point be

$$\mathbf{O} = \begin{pmatrix} x \\ y \end{pmatrix}.$$

The equidistance condition is:

$$\|\mathbf{O} - \mathbf{A}\| = \|\mathbf{O} - \mathbf{B}\|.$$

Squaring both sides:

$$\|\mathbf{O} - \mathbf{A}\|^2 = \|\mathbf{O} - \mathbf{B}\|^2.$$

Using vector dot product,

$$(\mathbf{O} - \mathbf{A})^\top (\mathbf{O} - \mathbf{A}) = (\mathbf{O} - \mathbf{B})^\top (\mathbf{O} - \mathbf{B}).$$

Expanding,

$$\mathbf{O}^\top \mathbf{O} - 2\mathbf{A}^\top \mathbf{O} + \mathbf{A}^\top \mathbf{A} = \mathbf{O}^\top \mathbf{O} - 2\mathbf{B}^\top \mathbf{O} + \mathbf{B}^\top \mathbf{B}.$$

Canceling terms,

$$-2\mathbf{A}^\top \mathbf{O} + \mathbf{A}^\top \mathbf{A} = -2\mathbf{B}^\top \mathbf{O} + \mathbf{B}^\top \mathbf{B}.$$

Rearranged,

$$2(\mathbf{B} - \mathbf{A})^\top \mathbf{O} = \mathbf{B}^\top \mathbf{B} - \mathbf{A}^\top \mathbf{A}.$$

Substituting values,

$$\begin{pmatrix} 4 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \frac{37 - 41}{2} = -2.$$

Simplified line equation:

$$2x + y = -1.$$

**Number of such points:** Infinitely many points lying on the above line.




Figure 1: Points  $A, B$  and equidistant line  $2x + y = -1$ .