

2.10.23

AI25BTECH11034 - Sujal Chauhan

September 24, 2025

# Question

The vector(s) which is/are coplanar with the vectors  $\hat{i} + \hat{j} + 2\hat{k}$  and  $\hat{i} + 2\hat{j} + \hat{k}$ , and perpendicular to vector  $\hat{i} + \hat{j} + \hat{k}$  is/are.

①  $\hat{j} - \hat{k}$

②  $\hat{i} + \hat{j}$

③  $\hat{i} - \hat{j}$

④  $\hat{j} + \hat{k}$

# Solution

Variable	Vector
<b>A</b>	$\begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$
<b>B</b>	$\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$
<b>C</b>	$\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$

# Solution

Listing options as vectors  $\mathbf{D}_i$ :

Input	Vector
$\mathbf{D}_1$	$\begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$
$\mathbf{D}_2$	$\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$
$\mathbf{D}_3$	$\begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$
$\mathbf{D}_4$	$\begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$

# Checking conditions

Let equation of plane be given by:

$$\mathbf{n}^\top \mathbf{X} = 1 \quad (1)$$

Let's find general solution  $\mathbf{n}$  which is perpendicular to the plane

$$\begin{pmatrix} A & B \end{pmatrix}^\top \mathbf{n} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (2)$$

$$\text{Given: } \begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & 1 \end{pmatrix} \mathbf{n} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (3)$$

(4)

We want to solve:  $\begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & 1 \end{pmatrix} \mathbf{n} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad \mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix} \quad (5)$

Form the augmented matrix: (6)

$$\left( \begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 1 & 2 & 1 & 1 \end{array} \right) \quad (7)$$

$$\xrightarrow{R_2 \rightarrow R_2 - R_1} \left( \begin{array}{ccc|c} 1 & 1 & 2 & 1 \\ 0 & 1 & -1 & 0 \end{array} \right) \quad (8)$$

$$\xrightarrow{R_1 \rightarrow R_1 - R_2} \left( \begin{array}{ccc|c} 1 & 0 & 3 & 1 \\ 0 & 1 & -1 & 0 \end{array} \right) \quad (9)$$

Thus the equations are: (10)

$$n_1 + 3n_3 = 1, \quad n_2 - n_3 = 0 \quad (11)$$

$$\text{Let } n_3 = a \in \mathbb{R}, \quad n_2 = a, \quad n_1 = 1 - 3a \quad (12)$$

So the general solution is: (13)

$$\mathbf{n} = \begin{pmatrix} 1 - 3a \\ a \\ a \end{pmatrix}, \quad a \in \mathbb{R} \quad (14)$$

Now any vector following both condition will be solution of the equation:

$$\begin{pmatrix} n & C \end{pmatrix}^T \mathbf{D}_i = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (15)$$

Checking Values for all options:

$$\begin{pmatrix} 1-3a & a & a \\ 1 & 1 & 1 \end{pmatrix} \mathbf{D}_i = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (16)$$



Vector	$\begin{pmatrix} 1-3a & a & a \\ 1 & 1 & 1 \end{pmatrix} \mathbf{D}_i$	Satisfies
$\mathbf{D}_1$	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	Yes
$\mathbf{D}_2$	$\begin{pmatrix} 1-2a \\ 2 \end{pmatrix}$	No
$\mathbf{D}_3$	$\begin{pmatrix} 1-4a \\ 0 \end{pmatrix}$	No
$\mathbf{D}_4$	$\begin{pmatrix} 2a \\ 2 \end{pmatrix}$	No

So only  $\mathbf{D}_1$  satisfies both conditions.




Figure: Vector  $\mathbf{D}_1$  in plane




Figure: Vector  $\mathbf{D}_2$  not coplanar




Figure: Vector  $\mathbf{D}_3$  not coplanar




Figure: Vector  $\mathbf{D}_4$  not coplanar

**Only 1 satisfies both conditions**