

# Matgeo Presentation - Problem 4.7.45

Revanth Siva Kumar.D – EE25BTECH11048

October 1, 2025

# QUESTION

For any three vectors  $\mathbf{a}$ ,  $\mathbf{b}$ ,  $\mathbf{c}$ , prove or disprove

$$(\mathbf{a} - \mathbf{b}) \cdot ((\mathbf{b} - \mathbf{c}) \times (\mathbf{c} - \mathbf{a})) = 2\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}).$$

## Solution:

We write the scalar triple product in determinant form:

$$(\mathbf{a} - \mathbf{b}) \cdot ((\mathbf{b} - \mathbf{c}) \times (\mathbf{c} - \mathbf{a})) = \det \begin{pmatrix} (\mathbf{a} - \mathbf{b})^T \\ (\mathbf{b} - \mathbf{c})^T \\ (\mathbf{c} - \mathbf{a})^T \end{pmatrix}. \quad (0.1)$$

Now observe that

$$(\mathbf{a} - \mathbf{b}) + (\mathbf{b} - \mathbf{c}) + (\mathbf{c} - \mathbf{a}) = \mathbf{0}. \quad (0.2)$$

Thus the three rows of the determinant are linearly dependent. From matrix theory, the determinant of a matrix with linearly dependent rows is zero. Hence

$$(\mathbf{a} - \mathbf{b}) \cdot ((\mathbf{b} - \mathbf{c}) \times (\mathbf{c} - \mathbf{a})) = 0. \quad (0.3)$$

## Solution:

On the other hand, the right-hand side is

$$2\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = 2 \det \begin{pmatrix} \mathbf{a}^T \\ \mathbf{b}^T \\ \mathbf{c}^T \end{pmatrix}, \quad (0.4)$$

which is not identically zero for arbitrary  $\mathbf{a}, \mathbf{b}, \mathbf{c}$ .

## Conclusion

The given statement is **false**.

The left-hand side  $(\mathbf{a} - \mathbf{b}) \cdot ((\mathbf{b} - \mathbf{c}) \times (\mathbf{c} - \mathbf{a}))$  is always zero, while the right-hand side  $2\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$  can be nonzero.