

2.10.33

Aditya Appana - EE25BTECH11004

September 8,2025

Question

Let α, β, γ be distinct real numbers. The points with position vectors $\alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$, $\beta\hat{i} + \gamma\hat{j} + \alpha\hat{k}$, $\gamma\hat{i} + \alpha\hat{j} + \beta\hat{k}$:

- ① are collinear
- ② form an equilateral triangle
- ③ form a scalene triangle
- ④ form a right angled triangle

Solution

To answer this question, we need to find the distance between each of these points.

Let \mathbf{A} be $\begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix}$, \mathbf{B} be $\begin{pmatrix} \beta \\ \gamma \\ \alpha \end{pmatrix}$, and \mathbf{C} be $\begin{pmatrix} \gamma \\ \alpha \\ \beta \end{pmatrix}$.

First, we need to check when the three points are collinear. We can do this using the collinearity matrix:

$$(\mathbf{C} - \mathbf{A} \quad \mathbf{B} - \mathbf{A})^T \quad (1)$$

If the rank of the matrix is 1, then the points are collinear.

$$\begin{pmatrix} \gamma - \alpha & \alpha - \beta & \beta - \gamma \\ \beta - \alpha & \gamma - \beta & \alpha - \gamma \end{pmatrix} \quad (2)$$

Solution

The rank of this matrix will be 1 only when all the elements in the bottom row of the matrix are equal to 0. This occurs only when $\alpha = \beta = \gamma$, which contradicts the fact that α, β, γ are distinct.

Therefore the points must be non-collinear and form a triangle.

The sides of the triangle are **$\mathbf{A} - \mathbf{B}$** , **$\mathbf{B} - \mathbf{C}$** , **$\mathbf{C} - \mathbf{A}$** .

Solution

$$\mathbf{A} - \mathbf{B} \text{ is } \begin{pmatrix} \alpha - \beta \\ \beta - \gamma \\ \gamma - \alpha \end{pmatrix} \quad (3)$$

$$\mathbf{B} - \mathbf{C} \text{ is } \begin{pmatrix} \beta - \gamma \\ \gamma - \alpha \\ \alpha - \beta \end{pmatrix} \quad (4)$$

$$\mathbf{C} - \mathbf{A} \text{ is } \begin{pmatrix} \gamma - \alpha \\ \alpha - \beta \\ \beta - \gamma \end{pmatrix} \quad (5)$$

$\|\mathbf{A} - \mathbf{B}\|$, $\|\mathbf{B} - \mathbf{C}\|$, $\|\mathbf{C} - \mathbf{A}\|$ are all equal, and equal to

$$\sqrt{(\alpha - \beta)^2 + (\beta - \gamma)^2 + (\gamma - \alpha)^2}$$

The three points therefore form an **equilateral triangle**, so option (2) is correct.

Solution

Then $\|\mathbf{A} - \mathbf{B}\|$, $\|\mathbf{B} - \mathbf{C}\|$, $\|\mathbf{C} - \mathbf{A}\|$ are all equal, and equal to

$$\sqrt{(\alpha - \beta)^2 + (\beta - \gamma)^2 + (\gamma - \alpha)^2}$$

The three points therefore form an equilateral triangle, so option (2) is correct.

Python Code

```
import numpy as np

vector = np.zeros(3)
vector[0] = input()
vector[1] = input()
vector[2] = input()

print(np.linalg.norm(vector))
```

```
#include<stdio.h>
#include<math.h>

float norm(float a, float b, float c){

float answer;
answer = pow(a,2) + pow(b,2) + pow(c,2);
answer = sqrt(answer);

return answer;

}
```


Python and C Code

```
import numpy as np
import ctypes
c_lib=ctypes.CDLL('./5c.so')

c_lib.norm.argtypes = [ctypes.c_float, ctypes.c_float, ctypes.c_float]
c_lib.norm.restype = ctypes.c_float

vector = np.zeros(3)
vector[0] = input()
vector[1] = input()
vector[2] = input()

answer = c_lib.norm(
    ctypes.c_float(vector[0]),
    ctypes.c_float(vector[1]),
    ctypes.c_float(vector[2]))
print(answer)
```