

# 4.7.43

Puni Aditya - EE25BTECH11046

## Question:

Show that the points  $(\hat{i} - \hat{j} + 3\hat{k})$  and  $3(\hat{i} + \hat{j} + \hat{k})$  are equidistant from the plane  $\mathbf{r} \cdot (5\hat{i} + 2\hat{j} - 7\hat{k}) + 9 = 0$  and lie on opposite sides of it.

## Solution:

Let the given points be  $\mathbf{P}_1 = \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$  and  $\mathbf{P}_2 = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix}$ . The equation of the given plane is

$$(5 \ 2 \ -7)\mathbf{x} + 9 = 0 \quad (1)$$

This can be written in the standard form  $\mathbf{n}^\top \mathbf{x} = k$ . Here,  $\mathbf{n} = \begin{pmatrix} 5 \\ 2 \\ -7 \end{pmatrix}$  and  $k = -9$ .

$$(5 \ 2 \ -7)\mathbf{x} = -9 \quad (2)$$

The reflection of point  $\mathbf{Q}$  with respect to the plane  $\mathbf{n}^\top \mathbf{x} = k$  is given by

$$\mathbf{R} = \mathbf{Q} - \frac{2(\mathbf{n}^\top \mathbf{Q} - k)}{\|\mathbf{n}\|^2} \mathbf{n} \quad (3)$$

Let the reflection of point  $\mathbf{P}_1$  with respect to the plane be  $\mathbf{Q}$ .

$$\mathbf{Q} = \mathbf{P}_1 - \frac{2(\mathbf{n}^\top \mathbf{P}_1 - k)}{\|\mathbf{n}\|^2} \mathbf{n} \quad (4)$$

$$= \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} - \frac{2 \left( (5 \ 2 \ -7) \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} - (-9) \right)}{5^2 + 2^2 + (-7)^2} \begin{pmatrix} 5 \\ 2 \\ -7 \end{pmatrix} \quad (5)$$

$$= \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} - \frac{-18}{78} \begin{pmatrix} 5 \\ 2 \\ -7 \end{pmatrix} \quad (6)$$

$$= \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix} + \frac{3}{13} \begin{pmatrix} 5 \\ 2 \\ -7 \end{pmatrix} \quad (7)$$

$$= \begin{pmatrix} \frac{28}{13} \\ -\frac{7}{13} \\ \frac{18}{13} \end{pmatrix} \quad (8)$$

Let a plane parallel to given plane pass through  $\mathbf{P}_1$ . Let this be  $\mathbf{n}^\top \mathbf{x} = c$

$$\mathbf{n}^\top \mathbf{Q} = c \quad (9)$$

$$(5 \quad 2 \quad -7) \begin{pmatrix} \frac{28}{13} \\ -\frac{7}{13} \\ \frac{18}{13} \end{pmatrix} = c \quad (10)$$

$$c = \frac{140}{13} - \frac{14}{13} - \frac{126}{13} \quad (11)$$

$$c = 0 \quad (12)$$


$$\mathbf{n}^\top \mathbf{P}_2 = (5 \quad 2 \quad -7) \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} \quad (13)$$

$$= 15 + 6 - 21 \quad (14)$$

$$= 0 = c \quad (15)$$

$\therefore \mathbf{P}_2$  lies in the plane  $\mathbf{n}^\top \mathbf{x} = c$ , the point  $\mathbf{P}_2$  and  $\mathbf{P}_1$  are equidistant from the plane  $\mathbf{n}^\top \mathbf{x} = k$  and lie on the opposite sides of the plane.

Points  $P_1$ ,  $P_2$  and Plane P



Plot