

Matrices in Geometry - 1.9.26

EE25BTECH11037 Divyansh

Aug, 2025

1 Problem

2 Solution

3 Final Answer

Problem Statement

Find the value of k , if the point $\mathbf{P}(2, 4)$ is equidistant from point $\mathbf{A}(5, k)$ and $\mathbf{B}(k, 7)$

Solution

Given: $\mathbf{P} \begin{pmatrix} 2 \\ 4 \end{pmatrix}$, $\mathbf{A} \begin{pmatrix} 5 \\ k \end{pmatrix}$ and a point $\mathbf{B} \begin{pmatrix} k \\ 7 \end{pmatrix}$ such that \mathbf{P} is equidistant from \mathbf{A} and \mathbf{B} .

$$\therefore \|\mathbf{A} - \mathbf{P}\| = \|\mathbf{B} - \mathbf{P}\|$$

On squaring both the sides, we get

$$\|\mathbf{A} - \mathbf{P}\|^2 = \|\mathbf{B} - \mathbf{P}\|^2$$

$$(\mathbf{A} - \mathbf{P})^\top (\mathbf{A} - \mathbf{P}) = (\mathbf{B} - \mathbf{P})^\top (\mathbf{B} - \mathbf{P})$$

Solution

$$(\mathbf{A} - \mathbf{P}) = \begin{pmatrix} 3 \\ k - 4 \end{pmatrix}$$

$$(\mathbf{B} - \mathbf{P}) = \begin{pmatrix} k - 2 \\ 3 \end{pmatrix}$$

$$(\mathbf{A} - \mathbf{P})^{\top} (\mathbf{A} - \mathbf{P}) = (3 \quad k - 4) \begin{pmatrix} 3 \\ k - 4 \end{pmatrix} = 9 + (k - 4)^2 = k^2 - 8k + 25$$

$$(\mathbf{B} - \mathbf{P})^{\top} (\mathbf{B} - \mathbf{P}) = (k - 2 \quad 3) \begin{pmatrix} k - 2 \\ 3 \end{pmatrix} = (k - 2)^2 + 9 = k^2 - 4k + 13$$

$$\text{From above, } k^2 - 8k + 25 = k^2 - 4k + 13$$

$$\implies 4k = 12 \implies k = 3$$

Final Answer

Hence, the final answer is $k = 3$

