

4.7.13

INDHIRESH S - EE25BTECH11027

13 September, 2025

Question

Find the distance between the lines l_1 and l_2 given by

$$\begin{aligned}\vec{r} &= \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k}) \\ \vec{r} &= 3\hat{i} + 3\hat{j} - 5\hat{k} + \mu(2\hat{i} + 3\hat{j} + 6\hat{k})\end{aligned}$$

Given equation:

$$\vec{r} = \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k}) \quad (1)$$

$$\vec{r} = 3\hat{i} + 3\hat{j} - 5\hat{k} + \mu(2\hat{i} + 3\hat{j} + 6\hat{k}) \quad (2)$$

$$(3)$$

Theoretical Solution

The given lines are in the form

$$\mathbf{r} = \mathbf{a}_1 + \lambda \mathbf{b} \quad (4)$$

$$\mathbf{r} = \mathbf{a}_2 + \mu \mathbf{b} \quad (5)$$

Where,

$$\mathbf{a}_1 = \begin{pmatrix} 1 \\ 2 \\ -4 \end{pmatrix} \quad \mathbf{a}_2 = \begin{pmatrix} 3 \\ 3 \\ -5 \end{pmatrix} \quad \mathbf{b} = \begin{pmatrix} 2 \\ 3 \\ 6 \end{pmatrix} \quad (6)$$

The given two lines are parallel. The distance between two parallel lines is given by:

$$d = \frac{\|(\mathbf{a}_2 - \mathbf{a}_1) \times \mathbf{b}\|}{\|\mathbf{b}\|} \quad (7)$$

$$\mathbf{a}_2 - \mathbf{a}_1 = \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} 2 \\ 3 \\ 6 \end{pmatrix} \quad (8)$$

Theoretical solution

Let,

$$\mathbf{a}_2 - \mathbf{a}_1 = \mathbf{a} \quad (9)$$

Now finding:

$$(\mathbf{a}_2 - \mathbf{a}_1) \times \mathbf{b} = \mathbf{a} \times \mathbf{b} \quad (10)$$

$$|\mathbf{A}_{23} \ \mathbf{B}_{23}| = \begin{vmatrix} 1 & 3 \\ -1 & 6 \end{vmatrix} = 9 \quad (11)$$

$$|\mathbf{A}_{31} \ \mathbf{B}_{31}| = \begin{vmatrix} 2 & 2 \\ -1 & 6 \end{vmatrix} = 14 \quad (12)$$

$$|\mathbf{A}_{12} \ \mathbf{B}_{12}| = \begin{vmatrix} 2 & 2 \\ 1 & 3 \end{vmatrix} = 4 \quad (13)$$

Theoretical Solution

$$\|\mathbf{a} \times \mathbf{b}\| = \left\| \begin{pmatrix} |\mathbf{A}_{23} \ \mathbf{B}_{23}| \\ |\mathbf{A}_{31} \ \mathbf{B}_{31}| \\ |\mathbf{A}_{12} \ \mathbf{B}_{12}| \end{pmatrix} \right\| \quad (14)$$

$$\|\mathbf{a} \times \mathbf{b}\| = \left\| \begin{pmatrix} 9 \\ 14 \\ 4 \end{pmatrix} \right\| \quad (15)$$

$$\|\mathbf{a} \times \mathbf{b}\| = \sqrt{293} \quad (16)$$

$$\|(\mathbf{a}_2 - \mathbf{a}_1) \times \mathbf{b}\| = \sqrt{293} \quad (17)$$

Theoretical Solution

$$\|\mathbf{b}\| = \sqrt{\mathbf{b}^T \mathbf{b}} \quad (18)$$

$$\|\mathbf{b}\| = \sqrt{4 + 9 + 36} = \sqrt{49} \quad (19)$$

$$\|\mathbf{b}\| = 7 \quad (20)$$

Substituting the values in Eq.7:

$$d = \frac{\sqrt{293}}{7} \quad (21)$$

Therefore the distance between the lines l_1 and l_2 is $\frac{\sqrt{293}}{7}$

```
#include <stdio.h>
#include <math.h>

int main() {
    double A[3] = {1,2,-4};
    double B[3] = {3,3,-5};
    double d[3] = {2,3,6};
    double AB[3], cross[3];
    double dist;

    // Compute B - A
    for(int i=0;i<3;i++) AB[i] = B[i] - A[i];

    // Cross product (AB x d)
    cross[0] = AB[1]*d[2] - AB[2]*d[1];
    cross[1] = AB[2]*d[0] - AB[0]*d[2];
    cross[2] = AB[0]*d[1] - AB[1]*d[0];
```



```
// Norms
double num = sqrt(cross[0]*cross[0] + cross[1]*cross[1] +
    cross[2]*cross[2]);
double den = sqrt(d[0]*d[0] + d[1]*d[1] + d[2]*d[2]);

dist = num / den;
printf(Distance between lines = %lf\n, dist);

// Output A, B, direction vector for plotting
printf(A: %lf %lf %lf\n, A[0], A[1], A[2]);
printf(B: %lf %lf %lf\n, B[0], B[1], B[2]);
printf(d: %lf %lf %lf\n, d[0], d[1], d[2]);

return 0;
}
```

Python Code

```
import ctypes
import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d.art3d import Poly3DCollection

# Load shared C library
lib = ctypes.CDLL('./distance.so')
lib.main()

# Define points and direction vector (same as in C)
A = np.array([1,2,-4])
B = np.array([3,3,-5])
d = np.array([2,3,6])

# Generate points for both lines
t = np.linspace(-2,2,10)
line1 = A[:,None] + d[:,None]*t
line2 = B[:,None] + d[:,None]*t
```

Python Code

```
# Plot
fig = plt.figure()
ax = fig.add_subplot(111, projection='3d')

# Lines
ax.plot(line1[0], line1[1], line1[2], 'b-', label='Line 11')
ax.plot(line2[0], line2[1], line2[2], 'g-', label='Line 12')

# Points
ax.scatter(*A,color='r',s=50)
ax.text(*A,A(1,2,-4),color='red')
ax.scatter(*B,color='orange',s=50)
ax.text(*B,B(3,3,-5),color='orange')

# Dotted line AB
ax.plot([A[0],B[0]], [A[1],B[1]], [A[2],B[2]], 'k--', label='
    Connecting AB')
```

```
ax.set_title(Figure)
ax.set_xlabel(X-axis)
ax.set_ylabel(Y-axis)
ax.set_zlabel(Z-axis)
ax.legend()
plt.savefig(/media/indhiresh-s/New Volume/Matrix/ee1030-2025/
ee25btech11027/MATGEO/4.7.13/figs/figure1.png)
plt.show()
```

Figure

