

# 2.10.55

EE25BTECH11026-Harsha

## Question:

The edges of a parallelopiped are of unit length and are parallel to non-coplanar unit vectors  $\hat{a}, \hat{b}, \hat{c}$  such that  $\hat{a} \cdot \hat{b} = \hat{b} \cdot \hat{c} = \hat{c} \cdot \hat{a} = \frac{1}{2}$ . Then, the volume of the parallelopiped is

1)  $\frac{1}{\sqrt{2}}$

2)  $\frac{1}{2\sqrt{2}}$

3)  $\frac{\sqrt{3}}{2}$

4)  $\frac{1}{\sqrt{3}}$

## Solution:

Let us solve the given equation theoretically and then verify the solution computationally.

According to the question, the edges of the parallelopiped are parallel to the unit vectors  $\hat{a}, \hat{b}, \hat{c}$  and

$$\hat{a}^T \hat{b} = \hat{b}^T \hat{c} = \hat{c}^T \hat{a} = \frac{1}{2}$$

As we know that the volume of parallelopiped is given by

$$V = [\mathbf{a} \ \mathbf{b} \ \mathbf{c}]$$

and

$$[\mathbf{a} \ \mathbf{b} \ \mathbf{c}][\mathbf{a} \ \mathbf{b} \ \mathbf{c}]^T = \mathbf{G}$$

where  $\mathbf{G}$  is the Gram Matrix.

$$\therefore \mathbf{G} = \begin{pmatrix} \hat{a}^T \hat{a} & \hat{a}^T \hat{b} & \hat{a}^T \hat{c} \\ \hat{b}^T \hat{a} & \hat{b}^T \hat{b} & \hat{b}^T \hat{c} \\ \hat{c}^T \hat{a} & \hat{c}^T \hat{b} & \hat{c}^T \hat{c} \end{pmatrix} = \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 1 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 1 \end{pmatrix}$$

For calculating the  $\det(\mathbf{G})$ , we can use the concept of eigen values.

Eigen values are those scalars which satisfies the following condition, For any non-zero eigen-vector  $\mathbf{v}$  and coefficient matrix  $\mathbf{M}$ ,

$$\mathbf{M}\mathbf{v} = \lambda\mathbf{v}, \text{ where } \lambda \text{ is an eigen value.}$$

$$\mathbf{G} = (1 - \rho)\mathbf{I} + \rho \mathbf{1}\mathbf{1}^T, \text{ where } \rho = \frac{1}{2} \text{ and } \mathbf{1} = \begin{pmatrix} 1 & 1 & 1 \end{pmatrix}$$

Let  $\mathbf{1}\mathbf{1}^T = \mathbf{J}$ . As we could see that the eigen-vector of  $\mathbf{J}$  is  $\mathbf{1}$  and by the rule,

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 3 \end{pmatrix} = 3\mathbf{1}$$

So, 3 is a eigen value of  $\mathbf{J}$ . Also, we know that the sum of eigen values is equal to trace of a matrix, we can say that the sum of the other eigen values would be 0. Also, we know that any orthogonal vector to  $\mathbf{1}$ , say  $\begin{pmatrix} -1 & 1 & 0 \end{pmatrix}^T$ ,

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} = \mathbf{0}$$

yields 0. Thus we can say that 0 is also one of the eigen value of  $\mathbf{J}$ . As sum of the other eigen values other than 3 is zero, the other eigen value must be zero.

$\therefore$  eigen values of  $\mathbf{J}$  are  $\{3, 0, 0\}$

Modifying the above equation on  $\mathbf{G}$ ,

$$\therefore \mathbf{G}\mathbf{v} = \frac{1}{2}\mathbf{I}\mathbf{v} + \frac{1}{2}\mathbf{J}\mathbf{v}$$

$$\implies \mathbf{G}\mathbf{v} = \frac{(1 + \mu)}{2}\mathbf{v}$$

where  $\mu$  is the eigen value of  $\mathbf{J}$ . Here the eigen value of  $\mathbf{G}$  is  $\frac{1+\mu}{2}$  and substituting the obtained eigen values of  $\mathbf{J}$  in this equation, we get the eigen values of  $\mathbf{G}$  to be  $\{2, \frac{1}{2}, \frac{1}{2}\}$

As we know that for eigen values of  $\mathbf{G}$  being  $\{\mu_1, \mu_2, \mu_3\}$ ,

$$\det(\mathbf{G}) = \mu_1\mu_2\mu_3$$

$$\therefore \det(\mathbf{G}) = 2 \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{2}$$

$$\implies V = \sqrt{\det(\mathbf{G})} = \frac{1}{\sqrt{2}} \text{ units}$$

From the figure, taking an example of vectors  $\mathbf{a}$  and  $\mathbf{b}$ , it is clearly verified that the theoretical solution matches with the computational solution.

Parallelepiped formed by unit vectors  $a$ ,  $b$ ,  $c$

