

12.144

EE25BTECH11026-Harsha

Question:

Let $\mathbf{A} = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 6 & 2 \\ 0 & 2 & 6 \end{pmatrix}$ and let $\lambda_1 \geq \lambda_2 \geq \lambda_3$ be the eigen values of \mathbf{A} .

(a) The triple $(\lambda_1, \lambda_2, \lambda_3)$ equals

1) (9, 4, 2)

2) (8, 4, 3)

3) (9, 3, 3)

4) (7, 5, 3)

(b) The Matrix \mathbf{P} such that

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$$

is

1) $\begin{pmatrix} \frac{1}{\sqrt{3}} & 0 & \frac{-2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{-2}{\sqrt{2}} & \frac{1}{\sqrt{6}} \end{pmatrix}$

2) $\begin{pmatrix} \frac{1}{\sqrt{3}} & \frac{-2}{\sqrt{6}} & 0 \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} & \frac{-1}{\sqrt{2}} \end{pmatrix}$

3) $\begin{pmatrix} 0 & 0 & 1 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} & 0 \end{pmatrix}$

4) $\begin{pmatrix} 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & 0 & \frac{-1}{\sqrt{2}} \end{pmatrix}$

Solution:

Let us solve the given question theoretically and then verify the solution computationally.

(a) The eigen values of \mathbf{A} is obtained using characteristic polynomial, which is given by,

$$\det|\mathbf{A} - \lambda\mathbf{I}| = 0 \quad (4.1)$$

$$\therefore \begin{vmatrix} 3 - \lambda & 0 & 0 \\ 0 & 6 - \lambda & 2 \\ 0 & 2 & 6 - \lambda \end{vmatrix} = 0 \quad (4.2)$$

$$\therefore (3 - \lambda)((6 - \lambda)^2 - 4) = 0 \quad (4.3)$$

$$\implies (\lambda - 3)(\lambda - 4)(\lambda - 8) = 0 \quad (4.4)$$

$$\therefore (\lambda_1, \lambda_2, \lambda_3) = (8, 4, 3) \quad (4.5)$$

(b) The given relation can be computed as ,

$$\mathbf{A} = \mathbf{PDP}^{-1} \quad (4.6)$$

where \mathbf{D} is the diagonal matrix of eigenvalues of \mathbf{A} .

From (4.6), we can infer that it is the Eigen-value decomposition of matrix \mathbf{A} .

Therefore, \mathbf{P} is the ortho-normalized matrix of collection of eigen vectors of \mathbf{A} .

$$\mathbf{P} = (\mathbf{v}_1 \quad \mathbf{v}_2 \quad \mathbf{v}_3) \quad (4.7)$$

where $\mathbf{v}_1, \mathbf{v}_2$ and \mathbf{v}_3 are the normalized eigen vectors of \mathbf{A} .

Eigenvectors \mathbf{v} for any square matrix \mathbf{A} is defined as

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v} \quad (4.8)$$

where λ is a scalar and is called the eigen value of \mathbf{A} .

$$\therefore (\mathbf{A} - \lambda\mathbf{I})\mathbf{v} = \mathbf{0} \quad (4.9)$$

For $\lambda = 3$,

$$\left(\begin{pmatrix} 3 & 0 & 0 \\ 0 & 6 & 2 \\ 0 & 2 & 6 \end{pmatrix} - 3 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) \mathbf{v} = \mathbf{0} \quad (4.10)$$

$$\text{Let } \mathbf{v} = \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix}.$$

$$\therefore \begin{pmatrix} 0 & 0 & 0 \\ 0 & 3 & 2 \\ 0 & 2 & 3 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (4.11)$$

$$\therefore 3\beta + 2\gamma = 0 \quad \text{and} \quad 2\beta + 3\gamma = 0 \quad (4.12)$$

$$\therefore \beta = \gamma = 0 \quad (4.13)$$

$$\mathbf{e}_1 = \begin{pmatrix} \alpha \\ 0 \\ 0 \end{pmatrix} \quad (4.14)$$

For $\lambda = 4$,

$$\left(\begin{pmatrix} 3 & 0 & 0 \\ 0 & 6 & 2 \\ 0 & 2 & 6 \end{pmatrix} - 4 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) \mathbf{v} = 0 \quad (4.15)$$

$$\text{Let } \mathbf{v} = \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix}.$$

$$\therefore \begin{pmatrix} -1 & 0 & 0 \\ 0 & 2 & 2 \\ 0 & 2 & 2 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (4.16)$$

$$\therefore 2\beta + 2\gamma = 0 \quad \text{and} \quad \alpha = 0 \quad (4.17)$$

$$\therefore \beta = -\gamma \quad (4.18)$$

$$\mathbf{e}_2 = \beta \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} \quad (4.19)$$

For $\lambda = 8$,

$$\left(\begin{pmatrix} 3 & 0 & 0 \\ 0 & 6 & 2 \\ 0 & 2 & 6 \end{pmatrix} - 8 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right) \mathbf{v} = 0 \quad (4.20)$$

$$\text{Let } \mathbf{v} = \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix}.$$

$$\therefore \begin{pmatrix} -5 & 0 & 0 \\ 0 & -2 & 2 \\ 0 & 2 & -2 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \\ \gamma \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad (4.21)$$

$$\therefore 2\beta - 2\gamma = 0 \quad \text{and} \quad \alpha = 0 \quad (4.22)$$

$$\therefore \beta = \gamma \quad (4.23)$$

$$\mathbf{e}_3 = \gamma \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \quad (4.24)$$

As we require unit eigen-vectors,

$$\Rightarrow \mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad \mathbf{v}_2 = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} \quad \mathbf{v}_3 = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} \end{pmatrix} \quad (4.25)$$

$$\therefore \mathbf{P} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & \frac{1}{\sqrt{2}} & \frac{-1}{\sqrt{2}} \end{pmatrix} \quad (4.26)$$