Question 1.4.15

AI25BTECH11040 - Vivaan Parashar

September 12, 2025

Question:

Find the area of the triangle formed by joining the midpoints of the sides of the triangle ABC, whose vertices are A(0,-1), B(2,1), and C(0,3)

Solution:

Let us start by finding the midpoints, let's call them D, E and F. The midpoint formula is: (Here the vectors represent position vectors of the points from the origin)

$$\mathbf{D} = \frac{\mathbf{A} + \mathbf{B}}{2} \tag{1}$$

$$\mathbf{E} = \frac{\mathbf{B} + \mathbf{C}}{2} \tag{2}$$

$$\mathbf{F} = \frac{\mathbf{C} + \mathbf{A}}{2} \tag{3}$$

$$\therefore \mathbf{D} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \mathbf{E} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \mathbf{F} = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \tag{4}$$

Now the area formula for a triangle with vertices at ${\bf P},\,{\bf Q}$ and ${\bf R}$ is given by:

Area =
$$\frac{1}{2} |(\mathbf{P} - \mathbf{Q}) \times (\mathbf{P} - \mathbf{R})|$$
 (5)

∴ Area of
$$\triangle DEF = \frac{1}{2} |(\mathbf{D} - \mathbf{E}) \times (\mathbf{D} - \mathbf{F})|$$
 (6)

$$= \frac{1}{2} \left| \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right) \times \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} - \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) \right| \tag{7}$$

$$= \frac{1}{2} \left| \begin{pmatrix} 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right| \tag{8}$$

$$=\frac{1}{2}|0-2|=1 \tag{9}$$

Diagram:




Figure: Diagram showing the triangle ABC and the triangle DEF.