

Matgeo Presentation - Problem 10.7.104

ee25btech11056 - Suraj.N

September 29, 2025

Problem Statement

Let a , b and λ be positive real numbers. Suppose P is an end point of the latus rectum of the parabola $y^2 = 4\lambda x$, and suppose the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ passes through the point P . If the tangents to the parabola and the ellipse at the point P are perpendicular to each other, then the eccentricity of ellipse is

Name	Value
Parabola	$\mathbf{x}^\top \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} -2\lambda \\ 0 \end{pmatrix}^\top \mathbf{x} = 0$
Ellipse	$\mathbf{x}^\top \begin{pmatrix} \frac{1}{a^2} & 0 \\ 0 & \frac{1}{b^2} \end{pmatrix} \mathbf{x} - 1 = 0$

Table : Parabola and Ellipse

Solution

The parameters of the parabola are :

$$\mathbf{v}_1 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad \mathbf{u}_1 = \begin{pmatrix} -2\lambda \\ 0 \end{pmatrix} \quad f_1 = 0 \quad (0.1)$$

The parameters of the ellipse are :

$$\mathbf{v}_2 = \begin{pmatrix} \frac{1}{a^2} & 0 \\ 0 & \frac{1}{b^2} \end{pmatrix} \quad \mathbf{u}_2 = \mathbf{0} \quad f_2 = -1 \quad (0.2)$$

The end point of the latus rectum of parabola is

$$\mathbf{P} = \begin{pmatrix} \lambda \\ 2\lambda \end{pmatrix} \quad (0.3)$$

Solution

The equation of tangent to the parabola at \mathbf{P} is given as :

$$(\mathbf{V}_1\mathbf{P} + \mathbf{u}_1)^\top \mathbf{x} + \mathbf{u}_1^\top \mathbf{P} + f_1 = 0 \quad \mathbf{n}_1 = \mathbf{V}_1\mathbf{P} + \mathbf{u}_1 \quad (0.4)$$

The equation of tangent to the ellipse at \mathbf{P} is given as :

$$(\mathbf{V}_2\mathbf{P} + \mathbf{u}_2)^\top \mathbf{x} + \mathbf{u}_2^\top \mathbf{P} + f_2 = 0 \quad \mathbf{n}_2 = \mathbf{V}_2\mathbf{P} + \mathbf{u}_2 \quad (0.5)$$

As the tangents at \mathbf{P} are perpendicular , the normal vectors of the tangents are also perpendicular

$$\mathbf{n}_1^\top \mathbf{n}_2 = 0 \quad (0.6)$$

$$(\mathbf{V}_1\mathbf{P} + \mathbf{u}_1)^\top \mathbf{V}_2\mathbf{P} = 0 \quad (0.7)$$

$$(\mathbf{P}^\top \mathbf{V}_1^\top + \mathbf{u}_1^\top) \mathbf{V}_2\mathbf{P} = 0 \quad (0.8)$$

$$\begin{pmatrix} -1 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{a^2} & 0 \\ 0 & \frac{1}{b^2} \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = 0 \quad (0.9)$$

$$\frac{a^2}{b^2} = \frac{1}{2} \quad (0.10)$$

Solution

From (0.2) , the eigen values of \mathbf{V}_2 are the diagonal entries as it is an upper triangular matrix and also $a < b$

$$\lambda_1 = \frac{1}{b^2} \qquad \lambda_2 = \frac{1}{a^2} \qquad (0.11)$$

The eccentricity e of ellipse is given as

$$e = \sqrt{1 - \frac{\lambda_1}{\lambda_2}} \qquad (0.12)$$

$$e = \sqrt{1 - \frac{a^2}{b^2}} \qquad (0.13)$$

From (0.10) , we get

$$e = \frac{1}{\sqrt{2}} \qquad (0.14)$$

Plot




Fig : Parabola and Ellipse