

12.18

Aditya Appana - EE25BTECH11004

October 5,2025

Question

X is 1 km northeast of **Y**. **Y** is 1 km southeast of **Z**. **W** is 1 km west of **Z**. **P** is 1 km south of **W**. **Q** is 1 km east of **P**. What is the distance between **X** and **Q** in km?

① 1

② $\sqrt{2}$

③ $\sqrt{3}$

④ 2

Let \mathbf{X} be $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$. Every subsequent vector can be expressed as a rotation by a particular angle θ with respect to the previous vector.
The matrix to rotate a vector by angle θ about the origin is:

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \quad (1)$$

Solution

X is 1km north-east of **Y**, so **Y** is 1km south-west of **X**. Therefore:

$$\mathbf{Y} - \mathbf{X} = \begin{pmatrix} \cos 225 & -\sin 225 \\ \sin 225 & \cos 225 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (2)$$

$$\mathbf{Y} - \mathbf{X} = \begin{pmatrix} \frac{-1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} \end{pmatrix} \quad (3)$$

Y is 1km south-east of **Z**, so **Z** is 1km north-west of **Y**. Therefore:

$$\mathbf{Z} - \mathbf{Y} = \begin{pmatrix} \cos 135 & -\sin 135 \\ \sin 135 & \cos 135 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (4)$$

$$\mathbf{Z} - \mathbf{Y} = \begin{pmatrix} \frac{-1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} \quad (5)$$

Solution

W is 1km west of **Z**. Therefore:

$$\mathbf{W} - \mathbf{Z} = \begin{pmatrix} \cos 180 & -\sin 180 \\ \sin 180 & \cos 180 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (6)$$

$$\mathbf{W} - \mathbf{Z} = \begin{pmatrix} -1 \\ 0 \end{pmatrix} \quad (7)$$

P is 1km south of **W**. Therefore:

$$\mathbf{P} - \mathbf{W} = \begin{pmatrix} \cos 270 & -\sin 270 \\ \sin 270 & \cos 270 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (8)$$

$$\mathbf{P} - \mathbf{W} = \begin{pmatrix} 0 \\ -1 \end{pmatrix} \quad (9)$$

Solution

Q is 1km east of **P**. Therefore:

$$\mathbf{Q} - \mathbf{P} = \begin{pmatrix} \cos 0 & -\sin 0 \\ \sin 0 & \cos 0 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (10)$$

$$\mathbf{Q} - \mathbf{P} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (11)$$

Adding (3), (5), (7), (9), (11) together, we get $\mathbf{Q} - \mathbf{X} = \mathbf{Q} = \begin{pmatrix} -\sqrt{2} \\ -1 \end{pmatrix}$.

Solution

The distance between $\mathbf{X} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ and $\mathbf{Q} = \begin{pmatrix} -\sqrt{2} \\ -1 \end{pmatrix}$ is:

$$\|\mathbf{X} - \mathbf{Q}\| = \left\| \begin{pmatrix} \sqrt{2} \\ 1 \end{pmatrix} \right\| \quad (12)$$

$$\sqrt{(\sqrt{2})^2 + 1^2} = \sqrt{3} \quad (13)$$

Therefore the correct option is **C**.

Codes Permalink

Figure




Figure: Plot