

## 4.11.6

J.NAVYASRI- EE25BTECH11028

september 2025

## Question:

Find the equation of the plane passing through the intersection of the planes

$$(r) \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$$

and

$$(r) \cdot (2\hat{i} + 3\hat{j} - \hat{k}) + 4 = 0$$

and parallel to the  $X$ -axis. Hence, find the distance of the plane from the  $X$ -axis.

## Solution:

The two given planes are

$$\Pi_1 : x + y + z = 1 \quad (1)$$

$$\Pi_2 : 2x + 3y - z + 4 = 0 \quad (2)$$

A plane passing through their intersection can be written as

$$\Pi : (x + y + z - 1) + \lambda(2x + 3y - z + 4) = 0 \quad (3)$$

Expanding:

$$x + y + z - 1 + \lambda(2x + 3y - z + 4) = 0 \quad (4)$$

$$x(1 + 2\lambda) + y(1 + 3\lambda) + z(1 - \lambda) + (-1 + 4\lambda) = 0 \quad (5)$$

A plane is parallel to the X-axis if the coefficient of x is zero:

$$1 + 2\lambda = 0 \Rightarrow \lambda = -\frac{1}{2} \quad (6)$$

$$y \left( 1 + 3 \left( -\frac{1}{2} \right) \right) + z \left( 1 - \left( -\frac{1}{2} \right) \right) + \left( -1 + 4 \left( -\frac{1}{2} \right) \right) = 0 \quad (7)$$

$$y \left( -\frac{1}{2} \right) + z \left( \frac{3}{2} \right) - 3 = 0 \quad (8)$$

Multiply through by 2:

$$-y + 3z - 6 = 0 \Rightarrow y - 3z + 6 = 0 \quad (9)$$

So, the required plane is

$$\boxed{y - 3z + 6 = 0} \quad (10)$$

The  $X$ -axis is the line  $y = 0, z = 0$ .

Distance from a plane  $y - 3z + 6 = 0$  to a point  $(0, 0, 0)$  on the  $X$ -axis is

$$d = \frac{|y_0 - 3z_0 + 6|}{\sqrt{1^2 + (-3)^2}} = \frac{|0 - 0 + 6|}{\sqrt{1 + 9}} = \frac{6}{\sqrt{10}} \quad (11)$$

$$\boxed{\text{Distance} = \frac{6}{\sqrt{10}}} \quad (12)$$

# Python Code

```
import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D

# Plane:  $y - 3z + 6 = 0 \Rightarrow y = 3z - 6$ 
# Create grid for plane
z = np.linspace(-5, 5, 20)
x = np.linspace(-5, 5, 20)
X, Z = np.meshgrid(x, z)
Y = 3*Z - 6

# Create figure
fig = plt.figure(figsize=(10,7))
ax = fig.add_subplot(111, projection='3d')
ax.plot_surface(X, Y, Z, alpha=0.5, color='cyan', rstride=1,
               cstride=1, edgecolor='k')
```

# Python Code

```
# Plot X-axis (y=0, z=0)
ax.plot([-5,5],[0,0],[0,0], color='red', linewidth=2, label='X-
axis')

# Point on X-axis (origin)
origin = np.array([0,0,0])

# Distance point on plane from origin (perpendicular foot)
# Plane: 0*x + 1*y - 3*z + 6 = 0 => normal = (0,1,-3)
normal = np.array([0,1,-3])
d = 6

# Formula for projection of origin onto plane
t = -(np.dot(normal, origin) + d) / np.dot(normal, normal)
foot = origin + t*normal

# Plot perpendicular distance line
ax.plot([origin[0], foot[0]], [origin[1], foot[1]], [origin[2],
foot[2]],
```

```
        color='green', linewidth=2, label='Shortest Distance')
# Plot origin and foot point
ax.scatter(*origin, color='red', s=50, label='Origin (on X-axis)'
)
ax.scatter(*foot, color='blue', s=50, label='Foot of
perpendicular')

# Labels
ax.set_xlabel('X-axis')
ax.set_ylabel('Y-axis')
ax.set_zlabel('Z-axis')
ax.set_title('Plane  $y - 3z + 6 = 0$  and Distance from X-axis')
ax.legend()
plt.savefig ("fig8.png")
plt.show()
```



# C Code

```
#include <stdio.h>
#include <math.h>

int main() {
    // Coefficients of the final plane:  $y - 3z + 6 = 0$ 
    double A = 0, B = 1, C = -3, D = 6;

    double x0 = 0, y0 = 0, z0 = 0;

    double numerator = fabs(A*x0 + B*y0 + C*z0 + D);
    double denominator = sqrt(A*A + B*B + C*C);
    double distance = numerator / denominator;

    printf("Equation of the plane:  $y - 3z + 6 = 0$ \n");
    printf("Distance of plane from X-axis: %.4f\n", distance);

    return 0;
}
```

```
import ctypes
import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D

lib = ctypes.CDLL("./libplane.so")

# Variables to store plane coefficients
a = ctypes.c_double()
b = ctypes.c_double()
c = ctypes.c_double()
d = ctypes.c_double()
```

```
# Call the C function to get coefficients
lib.get_plane(ctypes.byref(a), ctypes.byref(b), ctypes.byref(c),
             ctypes.byref(d))

# Extract values
a, b, c, d = a.value, b.value, c.value, d.value
print(f"Plane equation: {a}x + {b}y + {c}z + {d} = 0")

x = np.linspace(-5, 5, 30) # X range
z = np.linspace(-5, 5, 30) # Z range
X, Z = np.meshgrid(x, z)

# Solve plane eqn for Y:  $y = (-ax - cz - d)/b$ 
Y = (-a*X - c*Z - d) / b
```

# Python and C Code

```
fig = plt.figure(figsize=(8,6))
ax = fig.add_subplot(111, projection="3d")

ax.plot_surface(X, Y, Z, alpha=0.5, color="cyan", label="Plane")

ax.plot([-5, 5], [0, 0], [0, 0], color="red", linewidth=3, label=
        "X-axis")

# Take a point on X-axis (0,0,0)
P = np.array([0,0,0])

# Formula: foot of perpendicular from point to plane
num = a*P[0] + b*P[1] + c*P[2] + d
den = a*a + b*b + c*c

Px = P[0] - a*num/den
Py = P[1] - b*num/den
Pz = P[2] - c*num/den
```

```
# Plot distance line
ax.plot([P[0], Px], [P[1], Py], [P[2], Pz],
        color="black", linewidth=3, label="Shortest distance")

# Mark points
ax.scatter(P[0], P[1], P[2], color="red", s=50, label="Origin (on
X-axis)")
ax.scatter(Px, Py, Pz, color="blue", s=50, label="Foot on Plane")

ax.set_xlabel("X-axis")
ax.set_ylabel("Y-axis")
ax.set_zlabel("Z-axis")
ax.set_title("Plane  $y - 3z + 6 = 0$  and Distance from X-axis")
ax.legend()

plt.show()
```