

## 1.7.12

Rathlavath Jeevan-AI25BTECH11026

August 27,2025

# Question

Find the value of  $k$ , if the points  $P(5, 4)$ ,  $Q(7, k)$  and  $R(9, -2)$  are collinear.

*Hint:* Three points  $P(x_1, y_1)$ ,  $Q(x_2, y_2)$ ,  $R(x_3, y_3)$  are collinear if the area of the triangle formed by them is zero.

# Theoretical Solution

## Solution:

$$\mathbf{P} = \begin{pmatrix} 5 \\ 4 \end{pmatrix}, \quad \mathbf{Q} = \begin{pmatrix} 7 \\ k \end{pmatrix}, \quad \mathbf{R} = \begin{pmatrix} 9 \\ -2 \end{pmatrix} \quad (1)$$

**Collinearity via rank** Three points  $P, Q, R$  are collinear iff

$$\text{rank}(\mathbf{Q} - \mathbf{P} \quad \mathbf{R} - \mathbf{P}) = 1. \quad (2)$$

Compute the direction columns:

$$\mathbf{Q} - \mathbf{P} = \begin{pmatrix} 7 - 5 \\ k - 4 \end{pmatrix} = \begin{pmatrix} 2 \\ k - 4 \end{pmatrix}, \quad \mathbf{R} - \mathbf{P} = \begin{pmatrix} 9 - 5 \\ -2 - 4 \end{pmatrix} = \begin{pmatrix} 4 \\ -6 \end{pmatrix}. \quad (3)$$

Hence the collinearity matrix is

$$M = \begin{pmatrix} 2 & 4 \\ k - 4 & -6 \end{pmatrix}. \quad (4)$$

**Row reduction (rank = 1)**

# Theoretical Solution

$$\begin{pmatrix} 2 & 4 \\ k-4 & -6 \end{pmatrix} \xrightarrow{R_1 \leftarrow \frac{1}{2}R_1} \begin{pmatrix} 1 & 2 \\ k-4 & -6 \end{pmatrix} \xrightarrow{R_2 \leftarrow R_2 - (k-4)R_1} \begin{pmatrix} 1 & 2 \\ 0 & 2(1-k) \end{pmatrix}. \quad (5)$$

For  $\text{rank}(M) = 1$ , the second row must be the zero row:

$$2(1-k) = 0 \Rightarrow k = 1. \quad (6)$$

**Conclusion** For  $k = \boxed{1}$ , the three points  $P(5, 4)$ ,  $Q(7, k)$ ,  $R(9, -2)$  are collinear.

```
#include <stdio.h>

int main() {
    int x1 = 5, y1 = 4;
    int x2 = 7, y2; // y2 = k
    int x3 = 9, y3 = -2;
    int k;

    // Equation:  $x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2) = 0$ 
    // Substituting values
    //  $5(k - (-2)) + 7((-2) - 4) + 9(4 - k) = 0$ 
    // Solve manually inside program:

    // Simplified form:  $-4k + 4 = 0 \Rightarrow k = 1$ 
    k = 1;
```

```
printf("The value of k is: %d\n", k);  
  
return 0;  
}
```

```
import numpy as np
import matplotlib.pyplot as plt
from mpl_toolkits.mplot3d import Axes3D

# Points
x1, y1 = 5, 4
x2, y2 = 7, 1 # k = 1 (solution)
x3, y3 = 9, -2

# Create figure
fig = plt.figure()
ax = fig.add_subplot(111, projection='3d')
```

# Python Code

```
# Plot the points in 3D (z = 0 for 2D points)
ax.scatter([x1, x2, x3], [y1, y2, y3], [0, 0, 0], c='r', s=100,
           label='Points')

# Draw line through the points
xs = np.array([x1, x2, x3])
ys = np.array([y1, y2, y3])
zs = np.array([0, 0, 0])
ax.plot(xs, ys, zs, label='Collinear Line')

# Labels and title
ax.set_xlabel('X axis')
ax.set_ylabel('Y axis')
ax.set_zlabel('Z axis')
ax.set_title('3D Visualization of Collinear Points')
ax.legend()
```



```
# Save plot as picture
plt.savefig("collinear_points.png", dpi=300)

# Show the plot
plt.show()

print("Graph saved as collinear_points.png")
```

## 3D Visualization of Collinear Points

