

MatGeo Assignment 1.11.9

AI25BTECH11007

September 1, 2025

Question

If

$$\mathbf{a} = \hat{i} - 7\hat{j} + 7\hat{k} \quad \text{and} \quad \mathbf{b} = 3\hat{i} - 2\hat{j} + 2\hat{k},$$

find a unit vector perpendicular to both the vectors \mathbf{a} and \mathbf{b} .

Solution

We need a vector \mathbf{n} such that

$$\mathbf{n} \cdot \mathbf{a} = 0, \quad \mathbf{n} \cdot \mathbf{b} = 0 \quad (1)$$

where

$$\mathbf{a} = \begin{pmatrix} 1 \\ -7 \\ 7 \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} 3 \\ -2 \\ 2 \end{pmatrix}. \quad (2)$$

Let

$$\mathbf{n} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}. \quad (3)$$

Orthogonality conditions,

$$\mathbf{n} \cdot \mathbf{a} = x - 7y + 7z = 0 \quad (4)$$

$$\mathbf{n} \cdot \mathbf{b} = 3x - 2y + 2z = 0 \quad (5)$$

Solve equations,
From (4),

$$x = 7y - 7z. \quad (6)$$

Substitute (6) into (5):

$$3(7y - 7z) - 2y + 2z = 0 \quad (7)$$

$$21y - 21z - 2y + 2z = 0 \quad (8)$$

$$19y - 19z = 0 \quad (9)$$

$$y = z. \quad (10)$$

From (6) and (10):

$$x = 7y - 7y = 0. \quad (11)$$

Thus,

$$\mathbf{n} = \begin{pmatrix} 0 \\ y \\ y \end{pmatrix} = y \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}. \quad (12)$$

Normalize,


$$\hat{n} = \frac{\begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}}{\sqrt{0^2 + 1^2 + 1^2}} = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (13)$$

Hence, a unit vector perpendicular to both **a** and **b** is

$$\hat{n} = \frac{1}{\sqrt{2}}(\hat{j} + \hat{k}), \quad (14)$$

or its negative.

Vectors a (red), b (blue), and unit normal \hat{n} (green)



Conclusion

Therefore, a unit vector perpendicular to both **a** and **b** is

$$\hat{n} = \frac{1}{\sqrt{2}}(\hat{j} + \hat{k}),$$

or its negative.