

# Linear Regression II

---

Nipun Batra and the teaching staff

January 24, 2024

IIT Gandhinagar

## Relation between #instances and # Variables

If  $N < M$ , then it is an under-determined system

## Relation between #instances and # Variables

If  $N < M$ , then it is an under-determined system

Example:  $N=2$ ;  $M=3$

## Relation between #instances and # Variables

If  $N < M$ , then it is an under-determined system

Example:  $N=2$ ;  $M=3$

$$\begin{bmatrix} 30 \\ 40 \end{bmatrix} = \begin{bmatrix} 1 & 6 & 30 \\ 1 & 5 & 20 \end{bmatrix} \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix}$$

## Relation between #instances and # Variables

If  $N < M$ , then it is an under-determined system

Example:  $N=2$ ;  $M=3$

$$\begin{bmatrix} 30 \\ 40 \end{bmatrix} = \begin{bmatrix} 1 & 6 & 30 \\ 1 & 5 & 20 \end{bmatrix} \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix}$$

$$\begin{array}{rcl} 30 & = & \theta_0 + 6\theta_1 + 30\theta_2 \\ 40 & = & \theta_0 + 5\theta_1 + 20\theta_2 \\ \hline -10 & = & -1\theta_1 - 10\theta_2 \end{array} \quad (1)$$

The above equation can have infinitely many solutions.

Under-determined system:  $\epsilon_i = 0$  for all  $i$

## Relation between #instances and # Variables

What if  $N > M$

## Relation between #instances and # Variables

What if  $N > M$

Then it is an over determined system. So, the sum of squared residuals  $> 0$ .

## Class Exercise

Solve the linear system below using normal equation method

$x_1$	$x_2$	$y$
1	2	4
2	4	6
3	6	8



# Multi-collinearity

There can be situations where  $X^T X$  is not computable.

## Multi-collinearity

There can be situations where  $X^T X$  is not computable.

This condition arises when the  $|X^T X| = 0$ .

$$X = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 4 \\ 1 & 3 & 6 \end{bmatrix} \quad (2)$$

## Multi-collinearity

There can be situations where  $X^T X$  is not computable.

This condition arises when the  $|X^T X| = 0$ .

$$X = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 4 \\ 1 & 3 & 6 \end{bmatrix} \quad (2)$$

The matrix  $X$  is not full rank.

# Multi-collinearity

It arises when one or more predictor variable/feature in  $X$  can be expressed as a linear combinations of others

How to tackle it?

- Regularize

# Multi-collinearity

It arises when one or more predictor variable/feature in  $X$  can be expressed as a linear combinations of others

How to tackle it?

- Regularize
- Drop variables

# Multi-collinearity

It arises when one or more predictor variable/feature in  $X$  can be expressed as a linear combinations of others

How to tackle it?

- Regularize
- Drop variables
- Use different subsets of data

# Multi-collinearity

It arises when one or more predictor variable/feature in  $X$  can be expressed as a linear combinations of others

How to tackle it?

- Regularize
- Drop variables
- Use different subsets of data
- Avoid dummy variable trap

## Dummy variables

Say Pollution in Delhi =  $P$



## Dummy variables

Say Pollution in Delhi =  $P$

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * Wind\ speed + \theta_3 * Wind\ Direction$$

## Dummy variables

Say Pollution in Delhi =  $P$

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_2 * Wind\ speed + \theta_3 * Wind\ Direction$$

But, wind direction is a categorical variable.

# Dummy variables

Say Pollution in Delhi =  $P$

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_1 * \text{Wind speed} + \theta_3 * \text{Wind Direction}$$

But, wind direction is a categorical variable.

It is denoted as follows  $\{N:0, E:1, W:2, S:3\}$

# Dummy variables

Say Pollution in Delhi =  $P$

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_1 * Wind\ speed + \theta_3 * Wind\ Direction$$

But, wind direction is a categorical variable.

It is denoted as follows {N:0, E:1, W:2, S:3 }

Can we use the direct encoding?

# Dummy variables

Say Pollution in Delhi =  $P$

$$P = \theta_0 + \theta_1 * \#Vehicles + \theta_1 * Wind\ speed + \theta_3 * Wind\ Direction$$

But, wind direction is a categorical variable.

It is denoted as follows {N:0, E:1, W:2, S:3 }

Can we use the direct encoding?

Then this implies that  $S > W > E > N$

# Dummy Variables

N-1 Variable encoding

	Is it N?	Is it E?	Is it W?
N	1	0	0
E	0	1	0
W	0	0	1
S	0	0	0

# Dummy Variables

N Variable encoding

	Is it N?	Is it E?	Is it W?	Is it S?
N	1	0	0	0
E	0	1	0	0
W	0	0	1	0
S	0	0	0	1

Which is better  $N$  variable encoding or  $N-1$  variable encoding?



# Dummy Variables

Which is better N variable encoding or N-1 variable encoding?

The N-1 variable encoding is better because the N variable encoding can cause multi-collinearity.

# Dummy Variables

Which is better N variable encoding or N-1 variable encoding?

The N-1 variable encoding is better because the N variable encoding can cause multi-collinearity.

Is it  $S = 1 - (\text{Is it N} + \text{Is it W} + \text{Is it E})$

# Binary Encoding

N	00
E	01
W	10
S	11

## Binary Encoding

N	00
E	01
W	10
S	11

W and S are related by one bit.

## Binary Encoding

N	00
E	01
W	10
S	11

W and S are related by one bit.

This introduces dependencies between them, and this can cause confusion in classifiers.

## Interpreting Dummy variables

Gender	height
F	...
F	...
F	...
M	...
M	...

## Interpreting Dummy variables

Gender	height
F	...
F	...
F	...
M	...
M	...

Encoding

# Interpreting Dummy variables

Gender	height
F	...
F	...
F	...
M	...
M	...

Encoding

Is Female	height
1	...
1	...
1	...
0	...
0	...



## Interpreting Dummy Variables

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

$$height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$$

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

$$height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$$

We get  $\theta_0 = 5.8$  and  $\theta_0 = 6$

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

$$\text{height}_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$$

We get  $\theta_0 = 5.8$  and  $\theta_0 = 6$

$\theta_0 = \text{Avg height of Male} = 5.9$

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

$$height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$$

We get  $\theta_0 = 5.8$  and  $\theta_0 = 6$

$\theta_0 = \text{Avg height of Male} = 5.9$

$\theta_0 + \theta_1$  is chosen based (equal to) on 5, 5.2, 5.4 (for three records).

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

$$height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$$

We get  $\theta_0 = 5.8$  and  $\theta_0 = 6$

$\theta_0 = \text{Avg height of Male} = 5.9$

$\theta_0 + \theta_1$  is chosen based (equal to) on 5, 5.2, 5.4 (for three records).

$\theta_1$  is chosen based on 5-5.9, 5.2-5.9, 5.4-5.9

## Interpreting Dummy Variables

Is Female	height
1	5
1	5.2
1	5.4
0	5.8
0	6

$$height_i = \theta_0 + \theta_1 * (\text{Is Female}) + \epsilon_i$$

We get  $\theta_0 = 5.8$  and  $\theta_0 = 6$

$\theta_0 = \text{Avg height of Male} = 5.9$

$\theta_0 + \theta_1$  is chosen based (equal to) on 5, 5.2, 5.4 (for three records).

$\theta_1$  is chosen based on  $5-5.9$ ,  $5.2-5.9$ ,  $5.4-5.9$   $\theta_1 = \text{Avg. female height } (5+5.2+5.4)/3 - \text{Avg. male height}(5.9)$



## Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

## Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

$$x_i = \begin{cases} 1 & \text{if } i \text{ th person is female} \\ -1 & \text{if } i \text{ th person is male} \end{cases}$$

## Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

$$x_i = \begin{cases} 1 & \text{if } i \text{ th person is female} \\ -1 & \text{if } i \text{ th person is male} \end{cases}$$

$$y_i = \theta_0 + \theta_1 x_i + \epsilon_i = \begin{cases} \theta_0 + \theta_1 + \epsilon_i & \text{if } i \text{ th person is female} \\ \theta_0 - \theta_1 + \epsilon_i & \text{if } i \text{ th person is male.} \end{cases}$$

## Interpreting Dummy Variables

Alternatively, instead of a 0/1 coding scheme, we could create a dummy variable

$$x_i = \begin{cases} 1 & \text{if } i \text{ th person is female} \\ -1 & \text{if } i \text{ th person is male} \end{cases}$$

$$y_i = \theta_0 + \theta_1 x_i + \epsilon_i = \begin{cases} \theta_0 + \theta_1 + \epsilon_i & \text{if } i \text{ th person is female} \\ \theta_0 - \theta_1 + \epsilon_i & \text{if } i \text{ th person is male.} \end{cases}$$

Now,  $\theta_0$  can be interpreted as average person height.  $\theta_1$  as the amount that female height is above average and male height is below average.

## Alternative parameter estimation

$$\hat{y}_i = \theta_0 + \theta_1 x_i$$

## Alternative parameter estimation

$$\hat{y}_i = \theta_0 + \theta_1 x_i$$

$$\epsilon_i = y_i - \hat{y}_i$$

## Alternative parameter estimation

$$\hat{y}_i = \theta_0 + \theta_1 x_i$$

$$\epsilon_i = y_i - \hat{y}_i$$

$$\sum \epsilon_i^2 = \sum (y_i - \theta_0 - \theta_1 x_i)^2$$

## Alternative parameter estimation

$$\hat{y}_i = \theta_0 + \theta_1 x_i$$

$$\epsilon_i = y_i - \hat{y}_i$$

$$\sum \epsilon_i^2 = \sum (y_i - \theta_0 - \theta_1 x_i)^2$$

Now, we compute the derivative of it with all the  $\theta_j$ . Let us solve for  $x$  being a scalar.



## Alternative parameter estimation

$$\begin{aligned}\frac{\partial}{\partial \theta_0} \sum \epsilon_i^2 &= 2 \sum (y_i - \theta_0 - \theta_1 x_i)(-1) = 0 \\ 0 &= \sum y_i - N\theta_0 - \sum \theta_1 x_i \\ \theta_0 &= \frac{\sum y_i - \theta_1 \sum x_i}{N}\end{aligned}\tag{3}$$

$$\theta_0 = \bar{y} - \theta_1 \bar{x}$$

## Alternative parameter estimation

$$\frac{\partial}{\partial \theta_1} \sum \epsilon_i^2 = 0$$

$$\Rightarrow 2 \sum_{i=1}^N (y_i - \theta_0 - \theta_1 x_i)(-x_i) = 0$$

$$\Rightarrow \sum_{i=1}^N (x_i y_i - \theta_0 x_i - \theta_1 x_i^2) = 0$$

$$\Rightarrow \sum \theta_1 x_i^2 = \sum x_i y_i - \sum \theta_0 x_i$$

$$\Rightarrow \sum \theta_1 x_i^2 = \sum x_i y_i - \sum (\bar{y} - \theta_1 \bar{x}) x_i$$

## Alternative parameter estimation

$$\Rightarrow \sum \theta_1 x_i^2 = \sum x_i y_i - \bar{y} \sum x_i + \theta_1 \bar{x} \sum x_i$$

$$\Rightarrow \sum x_i y_i - \sum x_i \bar{y} = \theta_1 (-\bar{x} \sum x_i + \sum x_i^2)$$

$$\theta_1 = \frac{\sum x_i y_i - \sum x_i \bar{y}}{\sum x_i^2 - \bar{x} \sum x_i}$$

## Alternative parameter estimation

$$\theta_1 = \frac{\frac{1}{N} \sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y})}{\frac{1}{N} \sum_{i=1}^N (x_i - \bar{x})^2}$$

$$\theta_1 = \frac{\text{Cov}(x, y)}{\text{variance}(x)}$$