



Graphs and tangents

Sarang S. Sane




Recall : the graph of a function



Recall : the graph of a function

Let $f : X \rightarrow Y$ be a function. Then the graph of f is the subset

$$\Gamma(f) = \{(x, f(x)) | x \in X\} \subseteq X \times Y.$$




Recall : the graph of a function

Let $f : X \rightarrow Y$ be a function. Then the graph of f is the subset

$$\Gamma(f) = \{(x, f(x)) | x \in X\} \subseteq X \times Y.$$

Let $f : D \rightarrow \mathbb{R}$ be a function where $D \subseteq \mathbb{R}$. Then the graph of f can be drawn as a in \mathbb{R}^2 by considering points $\{(x, y) | y = f(x)\}$.




Recall : the graph of a function

Let $f : X \rightarrow Y$ be a function. Then the graph of f is the subset


$$\Gamma(f) = \{(x, f(x)) | x \in X\} \subseteq X \times Y.$$

Let $f : D \rightarrow \mathbb{R}$ be a function where $D \subseteq \mathbb{R}$. Then the graph of f can be drawn as a in \mathbb{R}^2 by considering points $\{(x, y) | y = f(x)\}$.

Example :




Graphs of functions : recall the trigonometric functions




Graph of $y = \sin(x)$

Graphs of functions : recall the trigonometric functions




Graph of $y = \sin(x)$

Graph of $y = \cos(x)$




Graphs of functions : more examples



$$f(x) = x^3 - 3x^2 + x + 2$$


Graphs of functions : more examples



$$\Gamma(x^3 - 3x^2 + x + 2)$$


$$\{(x, x^3 - 3x^2 + x + 2) \mid x \in \mathbb{R}\}$$

Graph of $y = e^x$




$$\{(x, e^x) \mid x \in \mathbb{R}\}$$

Graphs of functions : exponential decay



Graph of $y = e^{-x}$


Graphs of functions : the normal distribution




$$\text{Graph of } y = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

Curves

A **curve** is a figure that is obtained as the path of a moving point.



Curves : Visual example



Curves (contd.)

A curve can be thought of as a figure obtained by bending a line at various places.




Curves (contd.)

A curve can be thought of as a figure obtained by bending a line at various places.






Curves (contd.)

A curve can be thought of as a figure obtained by bending a line at various places.




Curves (contd.)

A curve can be thought of as a figure obtained by bending a line at various places.



The intuition of a tangent line to a curve



The intuition of a tangent line to a curve


A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .



The intuition of a tangent line to a curve

A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .


Traditionally, it was thought of as a line which *just touches* the curve at that point.



The intuition of a tangent line to a curve

A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .


Traditionally, it was thought of as a line which *just touches* the curve at that point.



The intuition of a tangent line to a curve

A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .


Traditionally, it was thought of as a line which *just touches* the curve at that point.



The intuition of a tangent line to a curve


A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .

Traditionally, it was thought of as a line which *just touches* the curve at that point.




Tangent lines : some means of identification

Often, a **tangent line** to a curve C at a point p (on C) has the property that it passes through the point p but does not intersect the curve C in any other point *close to* the point p , and lines parallel and close to it either do not intersect C *close to* p , or intersect the curve C in two (or more) points close to p .




Tangent lines : some means of identification

Often, a **tangent line** to a curve C at a point p (on C) has the property that it passes through the point p but does not intersect the curve C in any other point *close to* the point p , and lines parallel and close to it either do not intersect C *close to* p , or intersect the curve C in two (or more) points *close to* p .




What is a tangent (line) to a function?

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} .



What is a tangent (line) to a function?

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} . Assume that $\Gamma(f)$, the graph of f is a curve. Let $x \in D$.



What is a tangent (line) to a function?

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} . Assume that $\Gamma(f)$, the graph of f is a curve. Let $x \in D$.

Then a **tangent (line)** to f at x is a tangent (line) to $\Gamma(f)$ at the point $(x, f(x))$.



What is a tangent (line) to a function?

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} . Assume that $\Gamma(f)$, the graph of f is a curve. Let $x \in D$.

Then a **tangent (line)** to f at x is a tangent (line) to $\Gamma(f)$ at the point $(x, f(x))$.






Figure: Tangent lines for $y = x^3$

Tangent line for $y = 2^x$




Examples advising caution




$$\Gamma(\lfloor x \rfloor)$$

Examples advising caution



$\Gamma(\lfloor x \rfloor)$

Graph of $y = |x|$



Thank you

