

Multivariable Calculus

(Lecture-2)

Department of Mathematics
Bennett University
India

17th October, 2018

Learning Outcome of the Lecture

We learn

- **First Topic:** Visualizing functions
 - $F : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where $n = 2, 3$ - Curves
 - $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$: Contour Lines, Level Curves, Contour Plots
 - $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$: Level Surfaces

Learning Outcome of the Lecture

We learn

- **First Topic:** Visualizing functions
 - $F : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where $n = 2, 3$ - Curves
 - $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$: Contour Lines, Level Curves, Contour Plots
 - $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$: Level Surfaces
- **Second Topic:** Coordinates Systems
 - \mathbb{R}^2 : Cartesian Coordinates, Polar Coordinates
 - \mathbb{R}^3 : Cartesian Coordinates, Cylindrical Coordinates, Spherical Coordinates

Learning Outcome of the Lecture

We learn

- **First Topic:** Visualizing functions
 - $F : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where $n = 2, 3$ - Curves
 - $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$: Contour Lines, Level Curves, Contour Plots
 - $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$: Level Surfaces
- **Second Topic:** Coordinates Systems
 - \mathbb{R}^2 : Cartesian Coordinates, Polar Coordinates
 - \mathbb{R}^3 : Cartesian Coordinates, Cylindrical Coordinates, Spherical Coordinates
- **Third Topic:** Topology of Sets in \mathbb{R}^n :
 - Open Set, Closed Set, Bounded Set, Compact Set, Connected Set, Convex Set

First Topic

Visualizing Functions

Curves as Vector Valued Functions of Real Variable

Definition

Plane Curve:

A curve in the $2D$ -plane (\mathbb{R}^2) is defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^2$ where I is an interval in \mathbb{R} .



Curves as Vector Valued Functions of Real Variable

Definition

Plane Curve:

A curve in the 2D-plane (\mathbb{R}^2) is defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^2$ where I is an interval in \mathbb{R} .

Space Curve:

A curve in the Euclidean space \mathbb{R}^n can be defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where I is an interval in \mathbb{R} .

Examples:

Curves as Vector Valued Functions of Real Variable

Definition

Plane Curve:

A curve in the 2D-plane (\mathbb{R}^2) is defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^2$ where I is an interval in \mathbb{R} .

Space Curve:

A curve in the Euclidean space \mathbb{R}^n can be defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where I is an interval in \mathbb{R} .

Examples:

- Circle in \mathbb{R}^2 : $f(t) = (2 \cos t, 2 \sin t)$ for $t \in [0, 2\pi]$.



Curves as Vector Valued Functions of Real Variable

Definition

Plane Curve:

A curve in the 2D-plane (\mathbb{R}^2) is defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^2$ where I is an interval in \mathbb{R} .

Space Curve:

A curve in the Euclidean space \mathbb{R}^n can be defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where I is an interval in \mathbb{R} .

Examples:

- Circle in \mathbb{R}^2 : $f(t) = (2 \cos t, 2 \sin t)$ for $t \in [0, 2\pi]$.
- Parabola in \mathbb{R}^2 : $f(t) = (t, t^2)$ for $t \in \mathbb{R}$.



Curves as Vector Valued Functions of Real Variable

Definition

Plane Curve:

A curve in the 2D-plane (\mathbb{R}^2) is defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^2$ where I is an interval in \mathbb{R} .

Space Curve:

A curve in the Euclidean space \mathbb{R}^n can be defined as a continuous function $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$ where I is an interval in \mathbb{R} .

Examples:

- Circle in \mathbb{R}^2 : $f(t) = (2 \cos t, 2 \sin t)$ for $t \in [0, 2\pi]$.
- Parabola in \mathbb{R}^2 : $f(t) = (t, t^2)$ for $t \in \mathbb{R}$.
- Straight Line Segment joining the point A and B in \mathbb{R}^n :
 $f(t) = tB + (1 - t)A$ for $t \in [0, 1]$.

Curves in $2D$ or $3D$

Curves in \mathbb{R}^2 can be visualized by drawing the curve on the two-dimensional coordinates system ($2D$ Plane).

Curves in $2D$ or $3D$

Curves in \mathbb{R}^2 can be visualized by drawing the curve on the two-dimensional coordinates system ($2D$ Plane).



Curves in 2D or 3D

Curves in \mathbb{R}^2 can be visualized by drawing the curve on the two-dimensional coordinates system (2D Plane).



Curves in \mathbb{R}^3 can be visualized by drawing the curve in the three-dimensional coordinates system (3D Space).

Curves in 2D or 3D

Curves in \mathbb{R}^2 can be visualized by drawing the curve on the two-dimensional coordinates system (2D Plane).



Curves in \mathbb{R}^3 can be visualized by drawing the curve in the three-dimensional coordinates system (3D Space).

Example: Circular helix $f(t) = (\cos t, \sin t, t)$ for $t \geq 0$.


Curves in 2D or 3D

Curves in \mathbb{R}^2 can be visualized by drawing the curve on the two-dimensional coordinates system (2D Plane).



Curves in \mathbb{R}^3 can be visualized by drawing the curve in the three-dimensional coordinates system (3D Space).

Example: Circular helix $f(t) = (\cos t, \sin t, t)$ for $t \geq 0$.



Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

To visualize a function $z = f(x, y)$:

Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

To visualize a function $z = f(x, y)$:

- Draw the graph of f as a surface $z = f(x, y)$ in space, the surface being the set of points $(x, y, f(x, y))$. We can think the graph of f as the surface of a mountain.

Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

To visualize a function $z = f(x, y)$:

- Draw the graph of f as a surface $z = f(x, y)$ in space, the surface being the set of points $(x, y, f(x, y))$. We can think the graph of f as the surface of a mountain.
- If we think of the graph of f as the surface of a mountain, then the trace in a plane $z = c$ is a curve of constant altitude (which is called as **contour line**). A mountain climber walking along such a trace would neither ascend nor descend.



Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

To visualize a function $z = f(x, y)$:

- Draw the graph of f as a surface $z = f(x, y)$ in space, the surface being the set of points $(x, y, f(x, y))$. We can think the graph of f as the surface of a mountain.
- If we think of the graph of f as the surface of a mountain, then the trace in a plane $z = c$ is a curve of constant altitude (which is called as **contour line**). A mountain climber walking along such a trace would neither ascend nor descend.
- If we project a contour line on the xy -plane, we get the corresponding **level curve**.

Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

To visualize a function $z = f(x, y)$:

- Draw the graph of f as a surface $z = f(x, y)$ in space, the surface being the set of points $(x, y, f(x, y))$. We can think the graph of f as the surface of a mountain.
- If we think of the graph of f as the surface of a mountain, then the trace in a plane $z = c$ is a curve of constant altitude (which is called as **contour line**). A mountain climber walking along such a trace would neither ascend nor descend.
- If we project a contour line on the xy -plane, we get the corresponding **level curve**.
- A collection of level curves is called a **contour plot**.

Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $z = f(x, y)$ be a real valued function.

Definition

Contour Lines:

The **contour line** is the curve in which the surface $z = f(x, y)$ is cut by the plane $z = c$. We also call the contour line as the **trace of the graph of f in the plane $z = c$** .



Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $z = f(x, y)$ be a real valued function.

Definition

Contour Lines:

The **contour line** is the curve in which the surface $z = f(x, y)$ is cut by the plane $z = c$. We also call the contour line as the **trace of the graph of f in the plane $z = c$** .

Level Curves:

The **level curve** is a curve in the domain of f along which f has some constant value $f(x, y) = c$.



Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Let $z = f(x, y)$ be a real valued function.

Definition

Contour Lines:

The **contour line** is the curve in which the surface $z = f(x, y)$ is cut by the plane $z = c$. We also call the contour line as the **trace of the graph of f in the plane $z = c$** .

Level Curves:

The **level curve** is a curve in the domain of f along which f has some constant value $f(x, y) = c$.

Contour Plot or Contour Map:

A set/collection of level curves for $z = f(x, y)$ is called a **contour plot** or **contour map** of f .



Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Contour Line ($z = c$) = Collection of points (x, y, c) such that $f(x, y) = c$.

Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Contour Line ($z = c$) = Collection of points (x, y, c) such that $f(x, y) = c$.

Level curve (of height c) = $\{(x, y) \in \mathbb{R}^2 : f(x, y) = c\}$.



Visualizing Functions $f : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Contour Line ($z = c$) = Collection of points (x, y, c) such that $f(x, y) = c$.


Level curve (of height c) = $\{(x, y) \in \mathbb{R}^2 : f(x, y) = c\}$.

The idea behind Contour Plot is to provide three dimensional information in a two-dimensional setting.


Picture Example: Contour Lines, Level Curves, Contour Plot



Picture Example: Contour Lines and Level Curves



Picture Example: Some Contour Plot



Example: Contour Lines, Level Curves, Contour Plot

Let $f(x, y) = x^2 + y^2$.

Let $z = f(x, y)$ be the surface which is given by the set of points $\{(x, y, z) \in \mathbb{R}^3 : z = x^2 + y^2\}$.

Example: Contour Lines, Level Curves, Contour Plot

Let $f(x, y) = x^2 + y^2$.

Let $z = f(x, y)$ be the surface which is given by the set of points $\{(x, y, z) \in \mathbb{R}^3 : z = x^2 + y^2\}$.

Example:

Find the contour line and level curve of $f(x, y)$ for the value $c = 25$.

Example: Contour Lines, Level Curves, Contour Plot

Let $f(x, y) = x^2 + y^2$.

Let $z = f(x, y)$ be the surface which is given by the set of points $\{(x, y, z) \in \mathbb{R}^3 : z = x^2 + y^2\}$.

Example:

Find the contour line and level curve of $f(x, y)$ for the value $c = 25$.

- Contour line $f(x, y) = 25$ is:
 $x^2 + y^2 = 25$ on the plane $z = 25$.

Example: Contour Lines, Level Curves, Contour Plot

Let $f(x, y) = x^2 + y^2$.

Let $z = f(x, y)$ be the surface which is given by the set of points $\{(x, y, z) \in \mathbb{R}^3 : z = x^2 + y^2\}$.

Example:

Find the contour line and level curve of $f(x, y)$ for the value $c = 25$.

- Contour line $f(x, y) = 25$ is:
 $x^2 + y^2 = 25$ on the plane $z = 25$.
- Level curve $f(x, y) = 25$ is:
 $x^2 + y^2 = 25$ on the xy -plane.

Example: Contour Lines, Level Curves, Contour Plot

Let $f(x, y) = x^2 + y^2$.

Let $z = f(x, y)$ be the surface which is given by the set of points $\{(x, y, z) \in \mathbb{R}^3 : z = x^2 + y^2\}$.

Example:

Find the contour line and level curve of $f(x, y)$ for the value $c = 25$.

- Contour line $f(x, y) = 25$ is:
 $x^2 + y^2 = 25$ on the plane $z = 25$.
- Level curve $f(x, y) = 25$ is:
 $x^2 + y^2 = 25$ on the xy -plane.

Find the contour line and level curve of $f(x, y)$ for the value $c = 0$?

Confusion in terminologies

Because of the close association of contour lines with level curves there is no firm agreement about which word to use for which kind of curve. The convention in our course is that

Level curves lie in the domain of f
(in the plane $z = 0$)

while

Contour lines lie on the surface defined by f
(in the appropriate plane $z = c$).

Visualizing Functions $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$

To draw the graph of a function $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$, we need four dimensional space. So, we visualize/analyze the function f by drawing its **level surfaces**.

Visualizing Functions $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$

To draw the graph of a function $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$, we need four dimensional space. So, we visualize/analyze the function f by drawing its **level surfaces**.

Definition

Level Surfaces:



Visualizing Functions $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$

To draw the graph of a function $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$, we need four dimensional space. So, we visualize/analyze the function f by drawing its **level surfaces**.

Definition

Level Surfaces:

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a function given by $w = f(x, y, z)$. The point (x, y, z) in \mathbb{R}^3 at which the function $w = f(x, y, z)$ takes on a constant value $f(x, y, z) = c$ usually make up a surface called a **level surface** of the function f .



Visualizing Functions $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$

To draw the graph of a function $f : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$, we need four dimensional space. So, we visualize/analyze the function f by drawing its **level surfaces**.

Definition

Level Surfaces:

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a function given by $w = f(x, y, z)$. The point (x, y, z) in \mathbb{R}^3 at which the function $w = f(x, y, z)$ takes on a constant value $f(x, y, z) = c$ usually make up a surface called a **level surface** of the function f .

That is, for any c , the set of points (x, y, z) for which $f(x, y, z) = c$ (form a surface and) is called a **level surface** of the function.



Example of Level Surfaces

Example:

Describe the level surfaces of the function $w = \sqrt{x^2 + y^2 + z^2}$.

Example of Level Surfaces

Example:

Describe the level surfaces of the function $w = \sqrt{x^2 + y^2 + z^2}$.

The value of function w (in this particular example) can be interpreted as the distance from the origin to the point (x, y, z) in rectangular coordinates.

Example of Level Surfaces

Example:

Describe the level surfaces of the function $w = \sqrt{x^2 + y^2 + z^2}$.

The value of function w (in this particular example) can be interpreted as the distance from the origin to the point (x, y, z) in rectangular coordinates.

- $c > 0$:
- $c = 0$:
- $c < 0$:

Example of Level Surfaces

Example:

Describe the level surfaces of the function $w = \sqrt{x^2 + y^2 + z^2}$.

The value of function w (in this particular example) can be interpreted as the distance from the origin to the point (x, y, z) in rectangular coordinates.

- $c > 0$: Each level surface $\sqrt{x^2 + y^2 + z^2} = c$ where $c > 0$ is a sphere of radius c centered at the origin.
- $c = 0$:
- $c < 0$:

Example of Level Surfaces

Example:

Describe the level surfaces of the function $w = \sqrt{x^2 + y^2 + z^2}$.

The value of function w (in this particular example) can be interpreted as the distance from the origin to the point (x, y, z) in rectangular coordinates.

- $c > 0$: Each level surface $\sqrt{x^2 + y^2 + z^2} = c$ where $c > 0$ is a sphere of radius c centered at the origin.
- $c = 0$: The level surface $\sqrt{x^2 + y^2 + z^2} = 0$ contains the origin only.
- $c < 0$:

Example of Level Surfaces


Example:

Describe the level surfaces of the function $w = \sqrt{x^2 + y^2 + z^2}$.

The value of function w (in this particular example) can be interpreted as the distance from the origin to the point (x, y, z) in rectangular coordinates.

- $c > 0$: Each level surface $\sqrt{x^2 + y^2 + z^2} = c$ where $c > 0$ is a sphere of radius c centered at the origin.
- $c = 0$: The level surface $\sqrt{x^2 + y^2 + z^2} = 0$ contains the origin only.
- $c < 0$: There is no level surface in this case.

Example: Level Surfaces




Coordinate Systems





\mathbb{R}^2 : Cartesian Coordinates/ Rectangular Coordinates

Any point P in the plane (in 2D) can be assigned coordinates in the rectangular (or cartesian) coordinates system as (x, y) .



\mathbb{R}^2 : Polar Coordinates





- For each nonzero point $P = (x, y) \neq (0, 0)$ the polar coordinates (r, θ) of P are given by the equations

$$x = r \cos \theta, \quad y = r \sin \theta, \quad \text{or} \quad x^2 + y^2 = r^2, \quad \frac{y}{x} = \tan \theta.$$



\mathbb{R}^2 : Polar Coordinates





- For each nonzero point $P = (x, y) \neq (0, 0)$ the polar coordinates (r, θ) of P are given by the equations

$$x = r \cos \theta, \quad y = r \sin \theta, \quad \text{or} \quad x^2 + y^2 = r^2, \quad \frac{y}{x} = \tan \theta.$$

- The points (r, θ) and $(r, \theta + 2n\pi)$ where n is any integer denote the same (geometrical) point.



\mathbb{R}^2 : Polar Coordinates



- For each nonzero point $P = (x, y) \neq (0, 0)$ the polar coordinates (r, θ) of P are given by the equations

$$x = r \cos \theta, \quad y = r \sin \theta, \quad \text{or} \quad x^2 + y^2 = r^2, \quad \frac{y}{x} = \tan \theta.$$

- The points (r, θ) and $(r, \theta + 2n\pi)$ where n is any integer denote the same (geometrical) point.
- When P is origin, then $r = 0$ but θ is undetermined, so nothing can be done to remove the ambiguity (in defining θ) there.


\mathbb{R}^3 : Cartesian Coordinates/ Rectangular Coordinates

Select a point O to be the origin of the coordinate system, and choose three mutually perpendicular straight lines at O to be x , y , and z coordinate axes.

\mathbb{R}^3 : Cartesian Coordinates/ Rectangular Coordinates


Select a point O to be the origin of the coordinate system, and choose three mutually perpendicular straight lines at O to be x , y , and z coordinate axes.

For example: The y and z axes are in the plane, and the x -axis is perpendicular to this plane and points toward the reader.



\mathbb{R}^3 : Cylindrical Coordinates

A cylindrical coordinate system consists of polar coordinates (r, θ) in a plane together with a third coordinate z measured along an axis perpendicular to the $r\theta$ -plane which is the xy -plane. This means that the z -coordinate in the cylindrical coordinate system is the same as the z -coordinate in the cartesian system.



Relation between Cylindrical and Rectangular Coordinates

Cylindrical and Rectangular coordinates are related by the following equations:

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$z = z$$

where $r^2 = x^2 + y^2$ and $\tan \theta = \frac{y}{x}$.

Example

Problem: Describe the set of point $P(r, \theta, z)$ whose cylindrical coordinates satisfy:

(a) $r = \text{constant}$, (b) $\theta = \text{constant}$, (c) $z = \text{constant}$.

Example

Problem: Describe the set of point $P(r, \theta, z)$ whose cylindrical coordinates satisfy:

(a) $r = \text{constant}$, (b) $\theta = \text{constant}$, (c) $z = \text{constant}$.

Solution:

- The equation $r = a$ ($a > 0$) describes a **cylinder** about the z -axis. The equation $r = 0$ represents the z -axis.

Example

Problem: Describe the set of point $P(r, \theta, z)$ whose cylindrical coordinates satisfy:

(a) $r = \text{constant}$, (b) $\theta = \text{constant}$, (c) $z = \text{constant}$.

Solution:

- The equation $r = a$ ($a > 0$) describes a **cylinder** about the z -axis. The equation $r = 0$ represents the z -axis.
- The equation $\theta = \theta_0$ describes the **plane** that contains the z -axis and makes an angle θ_0 with the positive x -axis.

Example

Problem: Describe the set of point $P(r, \theta, z)$ whose cylindrical coordinates satisfy:


(a) $r = \text{constant}$, (b) $\theta = \text{constant}$, (c) $z = \text{constant}$.

Solution:

- The equation $r = a$ ($a > 0$) describes a **cylinder** about the z -axis. The equation $r = 0$ represents the z -axis.
- The equation $\theta = \theta_0$ describes the **plane** that contains the z -axis and makes an angle θ_0 with the positive x -axis.
- The equation $z = k$ describes a **plane** perpendicular to the z -axis.

\mathbb{R}^3 : Spherical Coordinates

Spherical coordinates are useful when there is a center of symmetry that we can take as the origin. The spherical coordinates (ρ, ϕ, θ) of a given point P are shown in the following Figure



Here $\rho = \sqrt{x^2 + y^2 + z^2}$.

- The **radial coordinate** ρ is the distance from the origin O to P ; ρ is always non-negative and is zero only if P coincides with the origin.



- The **radial coordinate** ρ is the distance from the origin O to P ; ρ is always non-negative and is zero only if P coincides with the origin.
- The **latitude angle**(second spherical coordinate) ϕ is the angle measured down from the z -axis to the line OP (That is, the angle from the positive z -axis to the line OP ; so, it is restricted to the interval $0 \leq \phi \leq \pi$).


- The **radial coordinate** ρ is the distance from the origin O to P ; ρ is always non-negative and is zero only if P coincides with the origin.
- The **latitude angle**(second spherical coordinate) ϕ is the angle measured down from the z -axis to the line OP (That is, the angle from the positive z -axis to the line OP ; so, it is restricted to the interval $0 \leq \phi \leq \pi$.
- The **longitude angle**(third spherical coordinate) θ is the same as in polar and cylindrical coordinates; it is the angle between the positive x -axis and the projection OQ of OP into the xy -plane. Usually, we will restrict θ to the interval $0 \leq \theta < 2\pi$.




- The **radial coordinate** ρ is the distance from the origin O to P ; ρ is always non-negative and is zero only if P coincides with the origin.
- The **latitude angle** (second spherical coordinate) ϕ is the angle measured down from the z -axis to the line OP (That is, the angle from the positive z -axis to the line OP ; so, it is restricted to the interval $0 \leq \phi \leq \pi$).
- The **longitude angle** (third spherical coordinate) θ is the same as in polar and cylindrical coordinates; it is the angle between the positive x -axis and the projection OQ of OP into the xy -plane. Usually, we will restrict θ to the interval $0 \leq \theta < 2\pi$.

Note: Incidentally, some books give spherical coordinates in the order (ρ, θ, ϕ) with the ϕ and θ reversed, and you should watch out for this when you read elsewhere.







ρ varies from
0 to ρ_0
with θ and ϕ fixed.



ϕ varies from
0 to π
with θ fixed.



θ varies from
0 to 2π .



Example

Problem: Describe the set of point $P(\rho, \phi, \theta)$ whose spherical coordinates satisfy:

(a) $\rho = \text{constant}$, (b) $\phi = \text{constant}$, (c) $\theta = \text{constant}$.

Example

Problem: Describe the set of point $P(\rho, \phi, \theta)$ whose spherical coordinates satisfy:

(a) $\rho = \text{constant}$, (b) $\phi = \text{constant}$, (c) $\theta = \text{constant}$.

Solution:

- The equation $\rho = k$ describes the **sphere** of radius k centered at the origin.

Example

Problem: Describe the set of point $P(\rho, \phi, \theta)$ whose spherical coordinates satisfy:

(a) $\rho = \text{constant}$, (b) $\phi = \text{constant}$, (c) $\theta = \text{constant}$.

Solution:

- The equation $\rho = k$ describes the **sphere** of radius k centered at the origin.
- The equation $\phi = \phi_0$ describes a single **cone** whose vertex lies at the origin and whose axis lies along the z -axis. If ϕ_0 is greater than $\frac{\pi}{2}$, the cone $\phi = \phi_0$ opens downward.

Example

Problem: Describe the set of point $P(\rho, \phi, \theta)$ whose spherical coordinates satisfy:

(a) $\rho = \text{constant}$, (b) $\phi = \text{constant}$, (c) $\theta = \text{constant}$.

Solution:

- The equation $\rho = k$ describes the **sphere** of radius k centered at the origin.
- The equation $\phi = \phi_0$ describes a single **cone** whose vertex lies at the origin and whose axis lies along the z -axis. If ϕ_0 is greater than $\frac{\pi}{2}$, the cone $\phi = \phi_0$ opens downward.
- The equation $\theta = \theta_0$ describes a **half-plane**, containing the z -axis whose trace in the xy -plane is the ray that makes an angle θ_0 with the positive x -axis.

- The relation between spherical coordinates and cylindrical coordinates are given by

$$r = \rho \sin \phi$$

$$z = \rho \cos \phi$$

$$\theta = \theta$$

- The relation between spherical coordinates and cylindrical coordinates are given by

$$r = \rho \sin \phi$$

$$z = \rho \cos \phi$$

$$\theta = \theta$$

- The relation between spherical coordinates and cartesian coordinates are given by

$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

$$z = \rho \cos \phi$$

*Topology of Sets
in
the Euclidean Space \mathbb{R}^n*

The Euclidean Space \mathbb{R}^n

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for } 1 \leq i \leq n\}.$$



The Euclidean Space \mathbb{R}^n

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for } 1 \leq i \leq n\}.$$

- Let $X = (x_1, x_2, \dots, x_n)$ and $Y = (y_1, y_2, \dots, y_n)$ be any two points in \mathbb{R}^n . Then,

The Euclidean Space \mathbb{R}^n

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for } 1 \leq i \leq n\}.$$

- Let $X = (x_1, x_2, \dots, x_n)$ and $Y = (y_1, y_2, \dots, y_n)$ be any two points in \mathbb{R}^n . Then,

$$d(X, Y) = |X - Y| = \|X - Y\| = \sqrt{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}.$$



The Euclidean Space \mathbb{R}^n

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for } 1 \leq i \leq n\}.$$

- Let $X = (x_1, x_2, \dots, x_n)$ and $Y = (y_1, y_2, \dots, y_n)$ be any two points in \mathbb{R}^n . Then,

$$d(X, Y) = |X - Y| = \|X - Y\| = \sqrt{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}.$$

- An **(open) ball** with center $A = (a_1, \dots, a_n)$ and radius r is denoted by $B(A, r)$ and is defined by

$$B(A, r) = \{X = (x_1, \dots, x_n) : d(X, A) = |X - A| < r\}.$$



The Euclidean Space \mathbb{R}^n

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{R} \text{ for } 1 \leq i \leq n\}.$$

- Let $X = (x_1, x_2, \dots, x_n)$ and $Y = (y_1, y_2, \dots, y_n)$ be any two points in \mathbb{R}^n . Then,

$$d(X, Y) = |X - Y| = \|X - Y\| = \sqrt{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}.$$

- An **(open) ball** with center $A = (a_1, \dots, a_n)$ and radius r is denoted by $B(A, r)$ and is defined by

$$B(A, r) = \{X = (x_1, \dots, x_n) : d(X, A) = |X - A| < r\}.$$

It is also denoted by $B_r(A)$.

It is also called an **(open) neighborhood** of A .

it is also denoted by $N(A, r)$ or $N(A)$ or $N_r(A)$.

Interior Point and Limit Point

Let S be a subset of \mathbb{R}^n .

Definition

A point X_0 is said to be an **interior point** of S if there is an open neighborhood $N(X_0)$ of X_0 such that $N(X_0) \subseteq S$.

Interior Point and Limit Point

Let S be a subset of \mathbb{R}^n .

Definition

A point X_0 is said to be an **interior point** of S if there is an open neighborhood $N(X_0)$ of X_0 such that $N(X_0) \subseteq S$.

The set of all interior points of S is called the **interior set** of S and is denoted by S^o or $Int(S)$.

Interior Point and Limit Point

Let S be a subset of \mathbb{R}^n .

Definition

A point X_0 is said to be an **interior point** of S if there is an open neighborhood $N(X_0)$ of X_0 such that $N(X_0) \subseteq S$.

The set of all interior points of S is called the **interior set** of S and is denoted by S^o or $Int(S)$.

Definition

A point X_0 is said to be a **limit point** (or accumulation point or cluster point) of S if every neighborhood $N(X_0)$ of X_0 contains at least one point from S which is not equal to X_0 .



Interior Point and Limit Point

Let S be a subset of \mathbb{R}^n .

Definition

A point X_0 is said to be an **interior point** of S if there is an open neighborhood $N(X_0)$ of X_0 such that $N(X_0) \subseteq S$.

The set of all interior points of S is called the **interior set** of S and is denoted by S^o or $Int(S)$.

Definition

A point X_0 is said to be a **limit point** (or accumulation point or cluster point) of S if every neighborhood $N(X_0)$ of X_0 contains at least one point from S which is not equal to X_0 .

The set of all limit points of S is called the **derived set** of S and is denoted by S' or $Der(S)$.



Boundary Point and Exterior Point

Let S be a subset of \mathbb{R}^n . The complement of the set S in \mathbb{R}^n is defined as $S^c = \{X \in \mathbb{R}^n, : X \notin S\} = \mathbb{R}^n \setminus S$.

Boundary Point and Exterior Point

Let S be a subset of \mathbb{R}^n . The complement of the set S in \mathbb{R}^n is defined as $S^c = \{X \in \mathbb{R}^n, : X \notin S\} = \mathbb{R}^n \setminus S$.

Definition

A point X_0 is said to be a **boundary point** of S if every neighborhood $N(X_0)$ of X_0 contains points from S as well as points from the complement of S .

Boundary Point and Exterior Point

Let S be a subset of \mathbb{R}^n . The complement of the set S in \mathbb{R}^n is defined as $S^c = \{X \in \mathbb{R}^n, : X \notin S\} = \mathbb{R}^n \setminus S$.

Definition

A point X_0 is said to be a **boundary point** of S if every neighborhood $N(X_0)$ of X_0 contains points from S as well as points from the complement of S .

The set of all boundary points of S is called the **boundary set** of S and is denoted by ∂S or $Bd(S)$.

Boundary Point and Exterior Point

Let S be a subset of \mathbb{R}^n . The complement of the set S in \mathbb{R}^n is defined as $S^c = \{X \in \mathbb{R}^n, : X \notin S\} = \mathbb{R}^n \setminus S$.

Definition

A point X_0 is said to be a **boundary point** of S if every neighborhood $N(X_0)$ of X_0 contains points from S as well as points from the complement of S .

The set of all boundary points of S is called the **boundary set** of S and is denoted by ∂S or $Bd(S)$.

Definition

A point X_0 is said to be an **exterior point** of S if there is an open neighborhood $N(X_0)$ of X_0 such that $N(X_0) \subseteq S^c$.
That is, X_0 is the interior point of S^c .

Boundary Point and Exterior Point

Let S be a subset of \mathbb{R}^n . The complement of the set S in \mathbb{R}^n is defined as $S^c = \{X \in \mathbb{R}^n, : X \notin S\} = \mathbb{R}^n \setminus S$.

Definition

A point X_0 is said to be a **boundary point** of S if every neighborhood $N(X_0)$ of X_0 contains points from S as well as points from the complement of S .

The set of all boundary points of S is called the **boundary set** of S and is denoted by ∂S or $Bd(S)$.

Definition

A point X_0 is said to be an **exterior point** of S if there is an open neighborhood $N(X_0)$ of X_0 such that $N(X_0) \subseteq S^c$.
That is, X_0 is the interior point of S^c .

The set of all exterior points of S is called the **exterior set** of S and is denoted by $Ext(S)$.

Open Set, Closed Set

Let S be a subset of \mathbb{R}^n .

Open Set, Closed Set

Let S be a subset of \mathbb{R}^n .

Definition

A set S is said to be **open** in \mathbb{R}^n if **each** point of S is an interior point of S .



Open Set, Closed Set

Let S be a subset of \mathbb{R}^n .

Definition

A set S is said to be **open** in \mathbb{R}^n if **each** point of S is an interior point of S .

That is, for each point X in S , there exists an open neighborhood $N(X)$ of X such that $N(X) \subseteq S$.

Open Set, Closed Set

Let S be a subset of \mathbb{R}^n .

Definition

A set S is said to be **open** in \mathbb{R}^n if **each** point of S is an interior point of S .

That is, for each point X in S , there exists an open neighborhood $N(X)$ of X such that $N(X) \subseteq S$.

Definition

A set S is said to be **closed** in \mathbb{R}^n if its complement set S^c is an open set in \mathbb{R}^n .



Open Set, Closed Set

Let S be a subset of \mathbb{R}^n .

Definition

A set S is said to be **open** in \mathbb{R}^n if **each** point of S is an interior point of S .

That is, for each point X in S , there exists an open neighborhood $N(X)$ of X such that $N(X) \subseteq S$.

Definition

A set S is said to be **closed** in \mathbb{R}^n if its complement set S^c is an open set in \mathbb{R}^n .

Basically a closed set is a set together with its boundary points. It can be characterized as: **A set S is closed if and only if it contains all its limit points.**



Bounded Set and Compact Set

Let S be a nonempty subset of \mathbb{R}^n .

Bounded Set and Compact Set

Let S be a nonempty subset of \mathbb{R}^n .

Definition

A nonempty set S is said to be a **bounded set** if **there exists** an open ball $B(X_0, r_0)$ for some $X_0 \in \mathbb{R}^n$ with $r_0 > 0$ such that $S \subseteq B(X_0, r_0)$.



Bounded Set and Compact Set

Let S be a nonempty subset of \mathbb{R}^n .

Definition

A nonempty set S is said to be a **bounded set** if **there exists** an open ball $B(X_0, r_0)$ for some $X_0 \in \mathbb{R}^n$ with $r_0 > 0$ such that $S \subseteq B(X_0, r_0)$. That is, the set S can be put inside an open ball with some center and a finite radius.



Bounded Set and Compact Set

Let S be a nonempty subset of \mathbb{R}^n .

Definition

A nonempty set S is said to be a **bounded set** if **there exists** an open ball $B(X_0, r_0)$ for some $X_0 \in \mathbb{R}^n$ with $r_0 > 0$ such that $S \subseteq B(X_0, r_0)$. That is, the set S can be put inside an open ball with some center and a finite radius.

Definition

A non-empty subset S of \mathbb{R}^n is said to be **compact** if it is closed and bounded.



Bounded Set and Compact Set

Let S be a nonempty subset of \mathbb{R}^n .

Definition

A nonempty set S is said to be a **bounded set** if **there exists** an open ball $B(X_0, r_0)$ for some $X_0 \in \mathbb{R}^n$ with $r_0 > 0$ such that $S \subseteq B(X_0, r_0)$. That is, the set S can be put inside an open ball with some center and a finite radius.

Definition

A non-empty subset S of \mathbb{R}^n is said to be **compact** if it is closed and bounded.

The above definition of compact set is true only in the Euclidean spaces.



Connected Set and Convex Set

Let S be a subset of \mathbb{R}^n .

Connected Set and Convex Set

Let S be a subset of \mathbb{R}^n .

Definition

An open set S is said to be **connected** in the Euclidean space \mathbb{R}^n if **any two** points of S can be joined by a path γ (continuous curve) such that the path γ lies **entirely** inside the set S .



Connected Set and Convex Set

Let S be a subset of \mathbb{R}^n .

Definition

An open set S is said to be **connected** in the Euclidean space \mathbb{R}^n if **any two** points of S can be joined by a path γ (continuous curve) such that the path γ lies **entirely** inside the set S .

A set S is said to be **disconnected** if it is **not connected**.

Connected Set and Convex Set

Let S be a subset of \mathbb{R}^n .

Definition

An open set S is said to be **connected** in the Euclidean space \mathbb{R}^n if **any two** points of S can be joined by a path γ (continuous curve) such that the path γ lies **entirely** inside the set S .

A set S is said to be **disconnected** if it is **not connected**.

Definition

A set S is said to be **convex** if **every** straight line segment L joining **any two** points of S lies **entirely** inside the set S .



Connected Set and Convex Set

Let S be a subset of \mathbb{R}^n .

Definition

An open set S is said to be **connected** in the Euclidean space \mathbb{R}^n if **any two** points of S can be joined by a path γ (continuous curve) such that the path γ lies **entirely** inside the set S .

A set S is said to be **disconnected** if it is **not connected**.

Definition

A set S is said to be **convex** if **every** straight line segment L joining **any two** points of S lies **entirely** inside the set S .

Note: Every convex set is connected, but converse is not true.