

Multivariable Calculus

(Lecture-3)

Department of Mathematics
Bennett University
India

23rd October, 2018

Learning Outcome of the Lecture

We learn

- Limits of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$
- Algebra of Limits
- How to show the limit exists?
- How to show the limit does not exist?
- Limit and Iterated Limits
- Continuity of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$
- Properties of continuous functions

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When both n and m are equal to 1, such a function is called real-valued function of a real variable,

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When both n and m are equal to 1, such a function is called **real-valued function** of a **real variable**, or more briefly, a **real function**.

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When both n and m are equal to 1, such a function is called **real-valued function** of a **real variable**, or more briefly, a real function.
- When $n = 1$ and $m > 1$, it is called a **vector-valued function** of a **real variable**,

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When both n and m are equal to 1, such a function is called a real-valued function of a real variable, or more briefly, a real function.
- When $n = 1$ and $m > 1$, it is called a vector-valued function of a real variable, or more briefly, a vector function.

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When both n and m are equal to 1, such a function is called a **real-valued function** of a **real variable**, or more briefly, a **real function**.
- When $n = 1$ and $m > 1$, it is called a **vector-valued function** of a **real variable**, or more briefly, a **vector function**.
- When $n > 1$ and $m = 1$, the function is called a **real-valued function** of a **vector variable**,



Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When **both n and m are equal to 1**, such a function is called **real-valued function** of a **real variable**, or more briefly, a real function.
- When **$n = 1$ and $m > 1$** , it is called a **vector-valued function** of a **real variable**, or more briefly, a vector function.
- When **$n > 1$ and $m = 1$** , the function is called a **real-valued function** of a **vector variable**, or more briefly, a scalar field.

Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When **both n and m are equal to 1**, such a function is called **real-valued function** of a **real variable**, or more briefly, a real function.
- When **$n = 1$ and $m > 1$** , it is called a **vector-valued function** of a **real variable**, or more briefly, a vector function.
- When **$n > 1$ and $m = 1$** , the function is called a **real-valued function** of a **vector variable**, or more briefly, a scalar field.
- When **$n > 1$ and $m > 1$** , it is called a **vector-valued function** of a **vector variable**,



Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m .

- When **both n and m are equal to 1**, such a function is called a **real-valued function** of a **real variable**, or more briefly, a **real function**.
- When **$n = 1$ and $m > 1$** , it is called a **vector-valued function** of a **real variable**, or more briefly, a **vector function**.
- When **$n > 1$ and $m = 1$** , the function is called a **real-valued function** of a **vector variable**, or more briefly, a **scalar field**.
- When **$n > 1$ and $m > 1$** , it is called a **vector-valued function** of a **vector variable**, or simply a **vector field**.



Examples of Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

- **Real function:** $f(x) = x^2 + \sin(x)$ for $x \in \mathbb{R}$.

Examples of Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

- **Real function:** $f(x) = x^2 + \sin(x)$ for $x \in \mathbb{R}$.
- **Vector function:** Motion of a Particle moving in the space

$$R : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^3, \quad R(t) = (t, 2t, e^{-t}) \quad \text{for } t > 0.$$

Examples of Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

- **Real function:** $f(x) = x^2 + \sin(x)$ for $x \in \mathbb{R}$.
- **Vector function:** Motion of a Particle moving in the space

$$R : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^3, \quad R(t) = (t, 2t, e^{-t}) \quad \text{for } t > 0.$$

- **Scalar field:** Temperature at any point in a cube $T : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$.

$$T(x, y, z) = x^2 - 2y + 5z^3 \quad \text{for } |x| < 1, |y| < 1, |z| < 1.$$

Examples of Functions $F : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$

- **Real function:** $f(x) = x^2 + \sin(x)$ for $x \in \mathbb{R}$.
- **Vector function:** Motion of a Particle moving in the space

$$R : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^3, \quad R(t) = (t, 2t, e^{-t}) \quad \text{for } t > 0.$$

- **Scalar field:** Temperature at any point in a cube $T : S \subseteq \mathbb{R}^3 \rightarrow \mathbb{R}$.

$$T(x, y, z) = x^2 - 2y + 5z^3 \quad \text{for } |x| < 1, |y| < 1, |z| < 1.$$

- **Vector field:** Vector Potential Function

$$V(x, y) = (x^2 + 2, e^x + 3y)$$

for $|x| \leq 1, |y| \leq 1$.

Component Functions / Coordinate Functions

Definition

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m where $n \geq 1$ and $m \geq 1$.
Let $X = (x_1, x_2, \dots, x_n)$ be a point in S .



Component Functions / Coordinate Functions

Definition

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m where $n \geq 1$ and $m \geq 1$.

Let $X = (x_1, x_2, \dots, x_n)$ be a point in S .

Then, $F(X)$ can be written as

$$F(X) = (f_1(X), f_2(X), \dots, f_m(X)),$$



Component Functions / Coordinate Functions

Definition

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m where $n \geq 1$ and $m \geq 1$.

Let $X = (x_1, x_2, \dots, x_n)$ be a point in S .

Then, $F(X)$ can be written as

$$F(X) = (f_1(X), f_2(X), \dots, f_m(X)),$$

where $f_i (1 \leq i \leq m)$ is a real valued function defined by

$$f_i(X) = \langle F(X), e_i \rangle \quad \text{for } X \in S,$$



Component Functions / Coordinate Functions

Definition

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m where $n \geq 1$ and $m \geq 1$.

Let $X = (x_1, x_2, \dots, x_n)$ be a point in S .

Then, $F(X)$ can be written as

$$F(X) = (f_1(X), f_2(X), \dots, f_m(X)),$$

where $f_i (1 \leq i \leq m)$ is a real valued function defined by

$$f_i(X) = \langle F(X), e_i \rangle \quad \text{for } X \in S,$$

and it is called a **component** (or **coordinate**) function of F .



Component Functions / Coordinate Functions

Definition

Let F be a function from $S \subseteq \mathbb{R}^n$ in to \mathbb{R}^m where $n \geq 1$ and $m \geq 1$.

Let $X = (x_1, x_2, \dots, x_n)$ be a point in S .

Then, $F(X)$ can be written as

$$F(X) = (f_1(X), f_2(X), \dots, f_m(X)),$$

where $f_i (1 \leq i \leq m)$ is a real valued function defined by

$$f_i(X) = \langle F(X), e_i \rangle \quad \text{for } X \in S,$$

and it is called a **component** (or **coordinate**) function of F .

- If $m = 1$, then the component function of F and the function F are same.

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$
- $f_2(x_1, x_2, x_3, x_4) = x_1^2,$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$
- $f_2(x_1, x_2, x_3, x_4) = x_1^2,$
- $f_3(x_1, x_2, x_3, x_4) = x_3 e^{x_1},$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$
- $f_2(x_1, x_2, x_3, x_4) = x_1^2,$
- $f_3(x_1, x_2, x_3, x_4) = x_3 e^{x_1},$
- $f_4(x_1, x_2, x_3, x_4) = 5x_2,$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$
- $f_2(x_1, x_2, x_3, x_4) = x_1^2,$
- $f_3(x_1, x_2, x_3, x_4) = x_3 e^{x_1},$
- $f_4(x_1, x_2, x_3, x_4) = 5x_2,$
- $f_5(x_1, x_2, x_3, x_4) = x_4.$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$
- $f_2(x_1, x_2, x_3, x_4) = x_1^2,$
- $f_3(x_1, x_2, x_3, x_4) = x_3 e^{x_1},$
- $f_4(x_1, x_2, x_3, x_4) = 5x_2,$
- $f_5(x_1, x_2, x_3, x_4) = x_4.$

Then

$$F(X) = (f_1(X), f_2(X), f_3(X), f_4(X), f_5(X)) \quad \text{for} \quad X = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4.$$

Example

Let $F : \mathbb{R}^4 \rightarrow \mathbb{R}^5$ be the function defined by

$$F(x_1, x_2, x_3, x_4) = (x_1 + x_2, x_1^2, x_3 e^{x_1}, 5x_2, x_4)$$

Define $f_i : \mathbb{R}^4 \rightarrow \mathbb{R}$ ($1 \leq i \leq 5$) as

- $f_1(x_1, x_2, x_3, x_4) = x_1 + x_2,$
- $f_2(x_1, x_2, x_3, x_4) = x_1^2,$
- $f_3(x_1, x_2, x_3, x_4) = x_3 e^{x_1},$
- $f_4(x_1, x_2, x_3, x_4) = 5x_2,$
- $f_5(x_1, x_2, x_3, x_4) = x_4.$

Then

$$F(X) = (f_1(X), f_2(X), f_3(X), f_4(X), f_5(X)) \quad \text{for } X = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4.$$

The functions f_i ($1 \leq i \leq 5$) are called the **component functions** or simply **components** of F .

Recall: Limits of Real Functions

Definition

Let $f : S \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a real function. Let a be a limit point of S .



Recall: Limits of Real Functions

Definition

Let $f : S \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a real function. Let a be a limit point of S . A point b is said to be the **limit of f at a** if



Recall: Limits of Real Functions

Definition

Let $f : S \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a real function. Let a be a limit point of S . A point b is said to be the **limit of f at a** if for **every** $\epsilon > 0$ there **exists** a $\delta > 0$ such that

$$x \in S \quad \text{and} \quad 0 < |x - a| < \delta \quad \Rightarrow \quad |f(x) - b| < \epsilon.$$



Recall: Limits of Real Functions

Definition

Let $f : S \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a real function. Let a be a limit point of S . A point b is said to be the **limit of f at a** if for **every** $\epsilon > 0$ there **exists** a $\delta > 0$ such that

$$x \in S \quad \text{and} \quad 0 < |x - a| < \delta \quad \Rightarrow \quad |f(x) - b| < \epsilon.$$

In this case we write

$$\lim_{x \rightarrow a} f(x) = b,$$

and say that $\lim_{x \rightarrow a} f(x)$ exists and is equal to b .



Recall: Limits of Real Functions

Definition

Let $f : S \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a real function. Let a be a limit point of S . A point b is said to be the **limit of f at a** if for **every** $\epsilon > 0$ there **exists** a $\delta > 0$ such that

$$x \in S \quad \text{and} \quad 0 < |x - a| < \delta \quad \Rightarrow \quad |f(x) - b| < \epsilon.$$

In this case we write

$$\lim_{x \rightarrow a} f(x) = b,$$

and say that $\lim_{x \rightarrow a} f(x)$ exists and is equal to b .

In other words, we say that **$f(x)$ approaches b as x approaches a** or **$f(x)$ has the limit b as x tends to a .**



Limits of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Definition

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. Let $A = (x_0, y_0)$ be a limit point of S .



Limits of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Definition

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. Let $A = (x_0, y_0)$ be a limit point of S . A point L is said to be the **limit of F at A** if



Limits of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Definition

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. Let $A = (x_0, y_0)$ be a limit point of S . A point L is said to be the **limit of F at A** if for **every** $\epsilon > 0$ there **exists** a $\delta > 0$ such that

$$X \in S \quad \text{and} \quad 0 < \|X - A\| < \delta \quad \Rightarrow \quad |F(X) - L| < \epsilon.$$



Limits of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Definition

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. Let $A = (x_0, y_0)$ be a limit point of S . A point L is said to be the **limit of F at A** if for **every** $\epsilon > 0$ there **exists** a $\delta > 0$ such that

$$X \in S \quad \text{and} \quad 0 < \|X - A\| < \delta \quad \Rightarrow \quad |F(X) - L| < \epsilon.$$

In this case we write

$$\lim_{X \rightarrow A} F(X) = L,$$

and say that $\lim_{X \rightarrow A} F(X)$ exists and is equal to L .



Limits of Functions $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$

Definition

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. Let $A = (x_0, y_0)$ be a limit point of S . A point L is said to be the **limit of F at A** if for **every** $\epsilon > 0$ there **exists** a $\delta > 0$ such that

$$X \in S \quad \text{and} \quad 0 < \|X - A\| < \delta \quad \Rightarrow \quad |F(X) - L| < \epsilon.$$

In this case we write

$$\lim_{X \rightarrow A} F(X) = L,$$

and say that $\lim_{X \rightarrow A} F(X)$ exists and is equal to L .

In other words, we say that **$F(X)$ approaches L as X approaches A** or **$F(X)$ has the limit L as X tends to A .**



Concept of the limit of functions in multivariable calculus
is essentially same as that of

Concept of the limit of functions in single variable calculus.

Concept of Limit is same, but it has a deeper insight

In multivariable calculus, the definition of limit of F has the following deeper insight:

- X approaches the point A means, “ X can approach A along any path/curve”.

Concept of Limit is same, but it has a deeper insight

In multivariable calculus, the definition of limit of F has the following deeper insight:

- X approaches the point A means, “ X can approach A along any path/curve”.
- $F(X)$ is approaching always the same point/value L , irrespective of the path on which X approaches A .

Concept of Limit is same, but it has a deeper insight

In multivariable calculus, the definition of limit of F has the following deeper insight:

- X approaches the point A means, “ X can approach A along any path/curve”.
- $F(X)$ is approaching always the same point/value L , irrespective of the path on which X approaches A .

In such case only, we are saying that the limit of $F(X)$ exists as X approaches A .

Concept of Limit is same, but it has a deeper insight

In multivariable calculus, the definition of limit of F has the following deeper insight:

- X approaches the point A means, “ X can approach A along any path/curve”.
- $F(X)$ is approaching always the same point/value L , irrespective of the path on which X approaches A .

In such case only, we are saying that the limit of $F(X)$ exists as X approaches A .

- That is, Along each path on which X approaches A , the function $F(X)$ is always approaching the same point/value L .

Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1:** Deciding to show that the limit exists Guessing the limiting value L .

Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit exists** Guessing the limiting value L . In this case, we expect that the limit will exist and the limiting value is guessed as $L = 0$.

Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1:** Deciding to show that the limit exists Guessing the limiting value L . In this case, we expect that the limit will exist and the limiting value is guessed as $L = 0$.
- **Step 2:** Estimating $|f(x, y) - L|$ in the neighborhood of $N(0, 0) \setminus (0, 0)$.



Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit exists** Guessing the limiting value L . In this case, we expect that the limit will exist and the limiting value is guessed as $L = 0$.
- **Step 2: Estimating $|f(x, y) - L|$ in the neighborhood of $N(0, 0) \setminus (0, 0)$.**

$$|f(x, y) - 0| \leq |x| + |y| \quad \text{for } (x, y) \in N(0, 0) \setminus (0, 0).$$

- **Step 3: Choosing δ :**

Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit exists** Guessing the limiting value L . In this case, we expect that the limit will exist and the limiting value is guessed as $L = 0$.
- **Step 2: Estimating $|f(x, y) - L|$ in the neighborhood of $N(0, 0) \setminus (0, 0)$.**

$$|f(x, y) - 0| \leq |x| + |y| \quad \text{for } (x, y) \in N(0, 0) \setminus (0, 0).$$

- **Step 3: Choosing δ :** Let $\epsilon > 0$ be given. Choose $\delta = \epsilon/2$. Then

$$0 < \|(x, y) - (0, 0)\| = \sqrt{x^2 + y^2} < \delta = \epsilon/2$$

Example

Let $f(x, y) = x \sin(1/y) + y \sin(1/x)$ if $xy \neq 0$, and $f(x, y) = 0$ if $xy = 0$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit exists** Guessing the limiting value L . In this case, we expect that the limit will exist and the limiting value is guessed as $L = 0$.
- **Step 2: Estimating $|f(x, y) - L|$ in the neighborhood of $N(0, 0) \setminus (0, 0)$.**

$$|f(x, y) - 0| \leq |x| + |y| \quad \text{for } (x, y) \in N(0, 0) \setminus (0, 0).$$

- **Step 3: Choosing δ :** Let $\epsilon > 0$ be given. Choose $\delta = \epsilon/2$. Then

$$0 < \|(x, y) - (0, 0)\| = \sqrt{x^2 + y^2} < \delta = \epsilon/2$$

$$\implies |f(x, y) - 0| \leq |x| + |y| \leq 2\sqrt{x^2 + y^2} < 2 \times \frac{\epsilon}{2} = \epsilon.$$



How to show Limit Does Not Exist

If we want to show $\lim_{X \rightarrow A} F(X)$ **DOES NOT** exist, then it is enough to show one of the following:

How to show Limit Does Not Exist

If we want to show $\lim_{X \rightarrow A} F(X)$ **DOES NOT** exist, then it is enough to show one of the following:

- $F(X)$ is approaching two different points/values L_1 and L_2 ($L_1 \neq L_2$) while X approaches A along two different paths,

OR

How to show Limit Does Not Exist

If we want to show $\lim_{X \rightarrow A} F(X)$ **DOES NOT** exist, then it is enough to show one of the following:

- $F(X)$ is approaching two different points/values L_1 and L_2 ($L_1 \neq L_2$) while X approaches A along two different paths,

OR

- there exists a path on which the function $F(X)$ is not approaching any point/value as X approaches A ,

OR

How to show Limit Does Not Exist

If we want to show $\lim_{X \rightarrow A} F(X)$ **DOES NOT** exist, then it is enough to show one of the following:

- $F(X)$ is approaching two different points/values L_1 and L_2 ($L_1 \neq L_2$) while X approaches A along two different paths,

OR

- there exists a path on which the function $F(X)$ is not approaching any point/value as X approaches A ,

OR

- there exists a path on which the function $|F(X)|$ is approaching ∞ as X approaches A .

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1:** Deciding to show that the limit does not exist.

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

- **Step 2: Path I**

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

- **Step 2: Path I**

Consider the path $\gamma_1 : y = x, x \neq 0$ and $x \rightarrow 0$.

As $(x, y) \rightarrow (0, 0)$ along γ_1 , $f(x, y) = \frac{x^2}{x^2+x^2} = \frac{1}{2} \rightarrow \frac{1}{2}$.



Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

- **Step 2: Path I**

Consider the path $\gamma_1 : y = x, x \neq 0$ and $x \rightarrow 0$.

As $(x, y) \rightarrow (0, 0)$ along γ_1 , $f(x, y) = \frac{x^2}{x^2+x^2} = \frac{1}{2} \rightarrow \frac{1}{2}$.

- **Step 3: Path II**

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

- **Step 2: Path I**

Consider the path $\gamma_1 : y = x, x \neq 0$ and $x \rightarrow 0$.

As $(x, y) \rightarrow (0, 0)$ along γ_1 , $f(x, y) = \frac{x^2}{x^2+x^2} = \frac{1}{2} \rightarrow \frac{1}{2}$.

- **Step 3: Path II**

Consider the path $\gamma_2 : y = 2x, x \neq 0$ and $x \rightarrow 0$

As $(x, y) \rightarrow (0, 0)$ along γ_2 , $f(x, y) = \frac{2x^2}{x^2+4x^2} = \frac{2}{5} \rightarrow \frac{2}{5}$.



Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

- **Step 2: Path I**

Consider the path $\gamma_1 : y = x, x \neq 0$ and $x \rightarrow 0$.

As $(x, y) \rightarrow (0, 0)$ along γ_1 , $f(x, y) = \frac{x^2}{x^2+x^2} = \frac{1}{2} \rightarrow \frac{1}{2}$.

- **Step 3: Path II**

Consider the path $\gamma_2 : y = 2x, x \neq 0$ and $x \rightarrow 0$

As $(x, y) \rightarrow (0, 0)$ along γ_2 , $f(x, y) = \frac{2x^2}{x^2+4x^2} = \frac{2}{5} \rightarrow \frac{2}{5}$.

- **Step 4: Conclusion**

Example

Let $f(x, y) = \frac{xy}{x^2+y^2}$ for $(x, y) \neq (0, 0)$. Examine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$.

- **Step 1: Deciding to show that the limit does not exist.**

In this case, we expect that the limit will not exist. We shall find two paths along which f approaches two different values.

- **Step 2: Path I**

Consider the path $\gamma_1 : y = x, x \neq 0$ and $x \rightarrow 0$.

As $(x, y) \rightarrow (0, 0)$ along γ_1 , $f(x, y) = \frac{x^2}{x^2+x^2} = \frac{1}{2} \rightarrow \frac{1}{2}$.

- **Step 3: Path II**

Consider the path $\gamma_2 : y = 2x, x \neq 0$ and $x \rightarrow 0$

As $(x, y) \rightarrow (0, 0)$ along γ_2 , $f(x, y) = \frac{2x^2}{x^2+4x^2} = \frac{2}{5} \rightarrow \frac{2}{5}$.

- **Step 4: Conclusion**

Since f approaches two different values as $(x, y) \rightarrow (0, 0)$ along two different paths, we conclude that $\lim_{(x,y) \rightarrow (0,0)} f(x, y)$ does **NOT** exist.



Algebra of Limits

Theorem

Algebra of Limits

Theorem

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ and $G : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be two functions. Let X_0 be a limit point of S .

Assume that $\lim_{X \rightarrow X_0} F(X)$ and $\lim_{X \rightarrow X_0} G(X)$ exist. Then,

Algebra of Limits

Theorem

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ and $G : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be two functions. Let X_0 be a limit point of S .

Assume that $\lim_{X \rightarrow X_0} F(X)$ and $\lim_{X \rightarrow X_0} G(X)$ exist. Then,

- $\lim_{X \rightarrow X_0} (F(X) + G(X)) = \lim_{X \rightarrow X_0} F(X) + \lim_{X \rightarrow X_0} G(X).$

Algebra of Limits

Theorem

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ and $G : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be two functions. Let X_0 be a limit point of S .

Assume that $\lim_{X \rightarrow X_0} F(X)$ and $\lim_{X \rightarrow X_0} G(X)$ exist. Then,

- $\lim_{X \rightarrow X_0} (F(X) + G(X)) = \lim_{X \rightarrow X_0} F(X) + \lim_{X \rightarrow X_0} G(X)$.
- $\lim_{X \rightarrow X_0} \lambda F(X) = \lambda \lim_{X \rightarrow X_0} F(X)$ for every scalar λ .

Algebra of Limits

Theorem

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ and $G : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be two functions. Let X_0 be a limit point of S .

Assume that $\lim_{X \rightarrow X_0} F(X)$ and $\lim_{X \rightarrow X_0} G(X)$ exist. Then,

- $\lim_{X \rightarrow X_0} (F(X) + G(X)) = \lim_{X \rightarrow X_0} F(X) + \lim_{X \rightarrow X_0} G(X)$.
- $\lim_{X \rightarrow X_0} \lambda F(X) = \lambda \lim_{X \rightarrow X_0} F(X)$ for every scalar λ .
- $\lim_{X \rightarrow X_0} |F(X)| = |\lim_{X \rightarrow X_0} F(X)|$, provided the limit in the R.H.S. exists.

Algebra of Limits

Theorem

Let $F : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ and $G : S \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ be two functions. Let X_0 be a limit point of S .

Assume that $\lim_{X \rightarrow X_0} F(X)$ and $\lim_{X \rightarrow X_0} G(X)$ exist. Then,

- $\lim_{X \rightarrow X_0} (F(X) + G(X)) = \lim_{X \rightarrow X_0} F(X) + \lim_{X \rightarrow X_0} G(X)$.
- $\lim_{X \rightarrow X_0} \lambda F(X) = \lambda \lim_{X \rightarrow X_0} F(X)$ for every scalar λ .
- $\lim_{X \rightarrow X_0} |F(X)| = |\lim_{X \rightarrow X_0} F(X)|$, provided the limit in the R.H.S. exists.
- If $\lim_{X \rightarrow X_0} G(X) \neq 0$, then

$$\lim_{X \rightarrow X_0} \left(\frac{F(X)}{G(X)} \right) = \frac{\lim_{X \rightarrow X_0} F(X)}{\lim_{X \rightarrow X_0} G(X)}.$$