

AutoML: Evaluation

Background: Statistical Hypotheses Tests

Bernd Bischl Frank Hutter Lars Kotthoff
Marius Lindauer Joaquin Vanschoren

Background: statistical hypothesis tests

- When we have a lot of data, we need to summarize it
 - ▶ But we already saw that summarization hides a lot of data
 - ▶ Ideally, we want to draw high-level conclusions (e.g., “A outperforms B on datasets of type X”)

Background: statistical hypothesis tests

- When we have a lot of data, we need to summarize it
 - ▶ But we already saw that summarization hides a lot of data
 - ▶ Ideally, we want to draw high-level conclusions (e.g., “A outperforms B on datasets of type X”)
 - Problem: we only have a finite number of observations
 - ▶ Can we attribute observed performance differences to chance?
 - ▶ Are we reasonably sure that a claim we make is reproducible?
- ~> Statistical tests can help

Statistical hypothesis testing

- 1 Define initial research hypothesis

Statistical hypothesis testing

- 1 Define initial research hypothesis
- 2 Derive null H_0 and alternative H_1 hypothesis
 - ▶ Alternative hypothesis should be your research hypothesis

First example: Courtroom Trial

- A prosecutor tries to prove the guilt of the defendant
- H_0 : The defendant is not guilty
 - ▶ Accepted for the moment
(“not guilty as long as their guilt is not proven”)
- H_1 : The defendant is guilty
 - ▶ prosecutor hopes to support that

First example: Courtroom Trial

- A prosecutor tries to prove the guilt of the defendant
- H_0 : The defendant is not guilty
 - ▶ Accepted for the moment
(“not guilty as long as their guilt is not proven”)
- H_1 : The defendant is guilty
 - ▶ prosecutor hopes to support that

First example: Courtroom Trial

- A prosecutor tries to prove the guilt of the defendant
- H_0 : The defendant is not guilty
 - ▶ Accepted for the moment
(“not guilty as long as their guilt is not proven”)
- H_1 : The defendant is guilty
 - ▶ prosecutor hopes to support that

	Truly not guilty	Truly guilty
Found not guilty	Acquittal	Type II Error
Found guilty	Type I Error	Conviction

⇒ We want to minimize Type I error!

Statistical hypothesis testing (cont'd)

- 1 Define initial research hypothesis
- 2 Derive null H_0 and alternative H_1 hypothesis
 - ▶ Alternative hypothesis should be your research hypothesis
- 3 Consider statistical assumptions
 - ▶ E.g., is your data Gaussian distributed?

Statistical hypothesis testing (cont'd)

- ① Define initial research hypothesis
- ② Derive null H_0 and alternative H_1 hypothesis
 - ▶ Alternative hypothesis should be your research hypothesis
- ③ Consider statistical assumptions
 - ▶ E.g., is your data Gaussian distributed?
- ④ Decide test and test statistic T
 - ▶ The correct test depends on your statistical assumptions.
 - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)

Statistical hypothesis testing (cont'd)

- ① Define initial research hypothesis
- ② Derive null H_0 and alternative H_1 hypothesis
 - ▶ Alternative hypothesis should be your research hypothesis
- ③ Consider statistical assumptions
 - ▶ E.g., is your data Gaussian distributed?
- ④ Decide test and test statistic T
 - ▶ The correct test depends on your statistical assumptions.
 - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)
- ⑤ Decide significance level α
(i.e., acceptable Type-I error to reject null hypothesis)

Statistical hypothesis testing (cont'd)

- ① Define initial research hypothesis
- ② Derive null H_0 and alternative H_1 hypothesis
 - ▶ Alternative hypothesis should be your research hypothesis
- ③ Consider statistical assumptions
 - ▶ E.g., is your data Gaussian distributed?
- ④ Decide test and test statistic T
 - ▶ The correct test depends on your statistical assumptions.
 - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)
- ⑤ Decide significance level α
(i.e., acceptable Type-I error to reject null hypothesis)
- ⑥ Compute observed t_{obs} of test statistic T
- ⑦ Calculate p -value given t_{obs}
 - ▶ i.e., probability under the null hypothesis of sampling a test statistic as extreme as observed (probability of Type-I error)

Statistical hypothesis testing (cont'd)

- 1 Define initial research hypothesis
- 2 Derive null H_0 and alternative H_1 hypothesis
 - ▶ Alternative hypothesis should be your research hypothesis
- 3 Consider statistical assumptions
 - ▶ E.g., is your data Gaussian distributed?
- 4 Decide test and test statistic T
 - ▶ The correct test depends on your statistical assumptions.
 - ▶ Typically: if you use more assumptions, the test is more powerful (i.e., less Type-I error)
- 5 Decide significance level α
(i.e., acceptable Type-I error to reject null hypothesis)
- 6 Compute observed t_{obs} of test statistic T
- 7 Calculate p -value given t_{obs}
 - ▶ i.e., probability under the null hypothesis of sampling a test statistic as extreme as observed (probability of Type-I error)
- 8 If $p < \alpha$, reject null hypothesis in favor of alternative hypothesis
 - ▶ If $p > \alpha$, it doesn't tell you anything about the null hypothesis!

Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”

Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”
- Null Hypothesis H_0 : $\mu = 100$ (μ is the population mean of this class)
- Alternative Hypothesis H_1 : $\mu > 100$ (one-sided hypothesis)

Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”
- Null Hypothesis H_0 : $\mu = 100$ (μ is the population mean of this class)
- Alternative Hypothesis H_1 : $\mu > 100$ (one-sided hypothesis)
- IQ values are known to be normally distributed with $X \sim \mathcal{N}(100, 15)$
 - ▶ \rightarrow statistical assumption

Second example for a statistical test

- Claim: “the students in this course are more intelligent than average”
- Null Hypothesis H_0 : $\mu = 100$ (μ is the population mean of this class)
- Alternative Hypothesis H_1 : $\mu > 100$ (one-sided hypothesis)
- IQ values are known to be normally distributed with $X \sim \mathcal{N}(100, 15)$
 - ▶ \rightarrow statistical assumption
- Let's say we observed IQ values x_i of 9 students in the class:
 - ▶ $\{x_1, \dots, x_9\} = \{116, 128, 125, 119, 89, 99, 105, 116, 118\}$.
 - ▶ The sample mean is $\bar{x} = 112.8$
 - ▶ Does this data support the claim?

Example continued


- Distribution of the test statistic

- ▶ Under H_0 , we know that each $x_i \sim \mathcal{N}(100, 15)$
- ▶ The test statistic that we measure is the sample mean $\bar{x} = \frac{1}{9} \sum_{i=1}^9 x_i$


Example continued

- Distribution of the test statistic

- ▶ Under H_0 , we know that each $x_i \sim \mathcal{N}(100, 15)$
- ▶ The test statistic that we measure is the sample mean $\bar{x} = \frac{1}{9} \sum_{i=1}^9 x_i$
- ▶ Under H_0 , the distribution of \bar{x} is $\mathcal{N}(100, 15/\sqrt{9})$
 - Our observation $\bar{x} = 112.8$ is quite extreme under that distribution




General principle




- Compare the test statistic (here: \bar{x}) to its sampling distribution under H_0

General principle




- Compare the test statistic (here: \bar{x}) to its sampling distribution under H_0
- **P-value:** probability p of observing values **at least as extreme as \bar{x}**

General principle



- Compare the test statistic (here: \bar{x}) to its sampling distribution under H_0
- **P-value:** probability p of observing values **at least as extreme as \bar{x}**
- Compare p to pre-defined confidence level α (usually $\alpha = 0.05$);
if $p < \alpha$, **reject H_0**

General principle



- Compare the test statistic (here: \bar{x}) to its sampling distribution under H_0
- **P-value:** probability p of observing values **at least as extreme as \bar{x}**
- Compare p to pre-defined confidence level α (usually $\alpha = 0.05$);
if $p < \alpha$, **reject H_0**
- With $\alpha = 0.01$, would we reject H_0 in this case?

Summary of example

- We just used a so-called *Z-test*
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions: $X \sim \mathcal{N}(\mu, \sigma^2)$, with known μ and σ^2

Summary of example

- We just used a so-called *Z-test*
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions: $X \sim \mathcal{N}(\mu, \sigma^2)$, with known μ and σ^2
- *Test statistic*: sample mean \bar{x} ; evaluate under $\mathcal{N}(\mu = \mu_0, s = \sigma^2/\sqrt{n})$


Summary of example

- We just used a so-called **Z-test**
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions: $X \sim \mathcal{N}(\mu, \sigma^2)$, with known μ and σ^2
- **Test statistic**: sample mean \bar{x} ; evaluate under $\mathcal{N}(\mu = \mu_0, s = \sigma^2/\sqrt{n})$
- Equivalent: compute the **Z-statistic**: $Z = (\bar{x} - \mu_0)/s$ and evaluate cumulative distribution $\Phi(Z)$ of Z under $\mathcal{N}(0, 1)$

Summary of example


- We just used a so-called *Z-test*
- $H_0: \mu = \mu_0, H_1: \mu > \mu_0$
- Assumptions: $X \sim \mathcal{N}(\mu, \sigma^2)$, with known μ and σ^2
- *Test statistic*: sample mean \bar{x} ; evaluate under $\mathcal{N}(\mu = \mu_0, s = \sigma^2/\sqrt{n})$
- Equivalent: compute the *Z-statistic*: $Z = (\bar{x} - \mu_0)/s$ and evaluate cumulative distribution $\Phi(Z)$ of Z under $\mathcal{N}(0, 1)$
 - ▶ There are standard tables to look up $\Phi(Z)$ for different values of Z
 - ▶ Nowadays, there are standard libraries to compute $\Phi(Z)$

Two-sided tests




- Similar to one-sided tests, but testing for extreme values in both tails
- Example Z-test: two-sided alternative hypothesis $H_1: \mu \neq \mu_0$

Two-sided tests



- Similar to one-sided tests, but testing for extreme values in both tails
- Example Z-test: two-sided alternative hypothesis $H_1: \mu \neq \mu_0$
- Compute $Z = (\bar{x} - \mu_0)/s$ as before
- Compute p -value as $p = 2\Phi(Z)$, to account for both tails

Two-sided tests



- Similar to one-sided tests, but testing for extreme values in both tails
- Example Z-test: two-sided alternative hypothesis $H_1: \mu \neq \mu_0$
- Compute $Z = (\bar{x} - \mu_0)/s$ as before
- Compute p -value as $p = 2\Phi(Z)$, to account for both tails
- With $\alpha = 0.01$, would we reject H_0 in this case?

General points about statistical hypothesis tests

- What if $p > \alpha$?
 - ▶ Failure to reject H_0
 - ▶ We cannot conclude that we can accept H_0 !

General points about statistical hypothesis tests

- What if $p > \alpha$?
 - ▶ Failure to reject H_0
 - ▶ We cannot conclude that we can accept H_0 !
- Beware (i): most tests make some assumptions
 - ▶ E.g., Z -test and popular t -test assume **normality**
 - ▶ Our data is often far from normally-distributed
 - ↪ E.g., exponential runtime distributions of optimizers
 - ↪ E.g., distribution of fitting a neural network with different random seeds is not well studied

General points about statistical hypothesis tests

- What if $p > \alpha$?
 - ▶ Failure to reject H_0
 - ▶ We cannot conclude that we can accept H_0 !
- Beware (i): most tests make some assumptions
 - ▶ E.g., Z -test and popular t -test assume normality
 - ▶ Our data is often far from normally-distributed
 - ↪ E.g., exponential runtime distributions of optimizers
 - ↪ E.g., distribution of fitting a neural network with different random seeds is not well studied
- Beware (ii): if you use cross-validation, observations are not independent (you cannot apply statistical tests that assume independence)