JEE–MAIN EXAMINATION – JANUARY 2026

(HELD ON THURSDAY 22nd JANUARY 2026) **TIME : 3:00 PM TO 6:00 PM**

CHEMISTRY — TEST PAPER WITH ANSWER

SECTION-A

51. At T(K), 100 g of 98% H_2SO_4 (w/w) aqueous solution is mixed with 100 g of 49% H_2SO_4 (w/w) aqueous solution. What is the mole fraction of H_2SO_4 in the resultant solution ?

(Given : Atomic mass H = 1 u ; S = 32 u; O = 16 u)


(Assume that temperature after mixing remains constant)

(1) 0.9 (2) 0.1

(3) 0.337 (4) 0.663

Ans. (4)

52. Consider the following reaction :



The product Y formed is :

(1) 2-methylhex-2-yne

(2) 5-methylhex-2-yne

(3) 2-methylhex-3-yne

(4) Isopropylbut-1-yne

Ans. (3)

53. $A + 2B \longrightarrow AB_2$

36.0 g of 'A' (Molar mass : 60 g mol^{-1}) and 56.0 g of 'B' (Molar mass : 80 g mol^{-1}) are allowed to react. Which of the following statements are **correct** ?

(A) 'A' is the limting reagent

(B) 77.0 g of AB_2 is formed

(C) Molar mass of AB_2 is 140 g mol^{-1}

(D) 15.0 g of A is left unreacted after the completion of reaction.

Choose the **correct** answer from the options given below :

(1) C and D only (2) A and C only

(3) B and D only (4) A and B only

Ans. (3)

54. Given below are two statements :

Statement-I L: The first ionization enthalpy of Cr is lower than that of Mn.

Statement-II : The second and third ionization enthalpies of Cr are higher than those of Mn.

In the light of the above statements, choose the **correct** answer from the options given below :


(1) Both Statement-I and Statement-II are false.

(2) Statement-I is true but Statement-II is false.


(3) Both Statement-I and Statement-II are true.


(4) Statement-I is false but Statement-II is true.


Ans. (2)


55. 

The final product [B] is :


(1) 

(2) 

(3) 

(4) 

Ans. (3)



56. When 1 g of compound (X) is subjected to Kjeldahl's method for estimation of nitrogen, 15 mL, 1M H_2SO_4 was neutralized by ammonia evolved. The percentage of nitrogen in compound (X) is :

(1) 21

(2) 0.42

(3) 42

(4) 0.21

Ans. (3)

57. Correct statements regarding Arrhenius equation among the following are :

(A) Factor $e^{-Ea/RT}$ corresponds to fraction of molecules having kinetic energy less than Ea.

(B) At a given temperature, lower the Ea, faster is the reaction.

(C) Increase in temperature by about 10°C doubles the rate of reaction.

(D) Plot of log k vs $\frac{1}{T}$ gives a straight line with slope $= -\frac{Ea}{R}$.

Choose the **correct** answer from the options given below :

(1) B and D only


(2) A and B only

(3) A and C only

(4) B and C only

Ans. (4)

58. The IUPAC name of the following comopound is :



(1) n-propyl-2-bromo-5-methylheptanoate

(2) 2-bromo-5-methylhexylpropanoate

(3) 2-bromo-5-methylpropanoate

(4) n-propyl-1-bromo-4-methylhexanoate

Ans. (1)

59. Given below are two statements :

Statement-I : Element 'X' and 'Y' are the most and least electronegative elements, respectively among N, As, Sb and P. The nature of the oxides X_2O_3 and Y_2O_3 is acidic and amphoteric, respectively.

Statement-II : BCl_3 is covalent in nature and gets hydrolysed in water. It produces $[B(OH)_4]^-$ and $[B(H_2O)_6]^{3+}$ in aqueous medium.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both Statement-I and Statement-II are true.

(2) Statement-I is true but Statement-II is false.

(3) Both Statement-I and Statement-II are false.

(4) Statement-I is false but Statement-II is true.

Ans. (2)

60. Match List-I with List-II.

List-I Reaction of glucose with	**List-II Product formed**
A. Hydroxylamine	I. Glouconic acid
B. Br_2 water	II. Glucose pentacetate
C. Excess acetic anhydride	III. Saccharic acid
D. Concentrated HNO_3	IV. Glucoxime

Choose the **correct** answer from the options given below :

(1) A-I, B-III, C-IV, D-II

(2) A-IV, B-I, C-II, D-III

(3) A-III, B-I, C-IV, D-II

(4) A-IV, B-III, C-II, D-I

Ans. (2)

61. Among H_2S, H_2O, NF_3, NH_3 and $CHCl_3$, identify the molecule (X) with lowest dipole moment value. The number of lone pairs of electrons present on the central atom of the molecule (X) is :

(1) 2 (2) 0

(3) 1 (4) 3

Ans. (3)



62. Given below are two statements :

Statement-I : C < O < N < F is the correct order in terms of first ionization enthalpy values.

Statement-II : S > Se > Te > Po > O is the correct order in terms of the magnitude of electron gain enthalpy values.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Statement-I is false but Statement-II is true

(2) Both Statement-I and Statement-II are true.

(3) Both Statement-I and Statement-II are false.

(4) Statement-I is true but Statement-II is false.

Ans. (2)

63. Which of the following mixture gives a buffer solution with pH = 9.25 ?

Given : pK_b (NH_4OH) = 4.75

(1) 0.2M NH_4OH (0.4 L) + 0.1M HCl (1L)

(2) 0.2M NH_4OH (0.5 L) + 0.1M HCl (0.5 L)

(3) 0.5M NH_4OH (0.2 L) + 0.2M HCl (0.5 L)

(4) 0.4M NH_4OH (1 L) + 0.1M HCl (1L)

Ans. (2)

64. The energy of first (lowest) Balmer line of H atom is x J. The energy (in J) of second Balmer line of H atom is :

(1) x^2 (2) $\frac{x}{1.35}$

(3) 2x (4) 1.35 x

Ans. (4)

65. Identify the **correct** statements :

A. Hydrated salts can be used as primary standard.

B. Primary standard should not undergo any reaction with air.

C. Reactions of primary standard with another substance should be instantaneous and stoichiometric.

D. Primary standard should not be soluble in water.

E. Primary standard should have low relative molar mass.

Choose the **correct** answer from the options given below :

(1) A, B, C and E only

(2) A, B, and C only

(3) A, B and E only

(4) D and E only

Ans. (2)

66. $[Ni(PPh_3)_2Cl_2]$ is a paramagnetic complex. Identify the **INCORRECT** statements about this complex.

A. The complex exhibits geometrical isomerism.

B. The complex is white in colour.

C. The calculated spin-only magnetic moment of the complex is 2.84 BM.

D. The calculated CFSE (Crystal Field Stabilization Energy) of Ni in this complex is $-0.8\Delta_0$.

E. The geometrical arrangement of ligands in this complex is similar to that in $Ni(CO)_4$.

Choose the **correct** answer from the options given below :

(1) A and B only

(2) A, B and D only

(3) C and D only

(4) C, D and E only

Ans. (2)

67. Consider the following reduction processes :

$Al^{3+} + 3e^- \rightarrow Al(s)$, $E^o = -1.66$ V

$Fe^{3+} + e^- \rightarrow Fe^{2+}$, $E^o = +0.77$ V

$Co^{3+} + e^- \rightarrow Co^{2+}$, $E^o = +1.81$ V

$Cr^{3+} + 3e^- \rightarrow Cr(s)$, $E^o = -0.74$ V

The tendency to act as reducing agent decreases in the order :


(1) Al > Cr > Fe^{2+} > Co^{2+}

(2) Al > Fe^{2+} > Cr > Co^{2+}

(3) Al > Cr > Co^{2+} > Fe^{2+}

(4) Cr > Fe^{2+} > Al > Co^{2+}

Ans. (1)





68. The compound A, $C_8H_8O_2$ reacts with acetophenone to form a single product via cross-Aldol condensation. The compound A on reaction with conc. NaOH forms a substituted benzyl alcohol as


(1) 2-hydroxy acetophenone
(2) 4-methyoxy benzaldehyde
(3) 4-hydroxy benzylaldehyde
(4) 4-methyl benzoic acid


Ans. (2)


69. 3, 3-Dimenthyl-2-butanol **cannot** be prepared by :

A. 

B. 

C. 

D. 


E. 


Choose the **correct** answer from the options given below :


(1) B only (2) B and E only
(3) B and C only (4) B, C and E only


Ans. (2)

70. The dibromo compound [P] (molecular formula : $C_9H_{10}Br_2$) when heated with excess sodamide followed by treatment with dilute HCl gives [Q]. On warming [Q] with mercuric sulphate and dilute sulphuric acid yield [R] which gives positive Iodoform test but negative Tollen's test. The compound [P] is :

(1) 

(2) 

(3) 

(4) 

Ans. (3)

SECTION-B

71. Consider the following electrochemical cell :

$Pt \mid O_2(g)\ (1bar) \mid HCl\ (aq) \parallel M^{2+}(aq,\ 1.0\ M) \mid M(s)$

The pH above which, oxygen gas would start to evolve at anode is _____ (nearest integer).

$$\left[\text{Given : } \left.\begin{matrix} E^0_{M^{2+}/M} = 0.994V \\ E^0_{O_2/H_2O} = 1.23V \end{matrix}\right\} \text{standard reduction potential} \quad \text{and } \frac{RT}{F}(2.303) = 0.059\ V \text{ at the given condition}\right]$$

Ans. (4)

72. If the enthalpy of sublimation of Li is 155 kJ mol^{-1}, enthalpy of dissociation of F_2 is 150 kJ mol^{-1}, ionization enthalpy of Li is 520 kJ mol^{-1}, electron gain enthalpy of F is –313 kJ mol^{-1}, standard enthalpy of formation of LiF is –594 kJ mol^{-1}. The magnitude of lattice enthalpy of LiF is _____ kJ mol^{-1} (Nearest integer).

Ans. (1031)

73. Among the following oxides of 3d elements, the number of mixed oxides are_____.

Ti_2O_3, V_2O_4, Cr_2O_3, Mn_3O_4, Fe_3O_4, Fe_2O_3, Co_3O_4

Ans. (3)

74. The mass of benzanilide obtained from the benzoylation reaction of 5.8 g of aniline, if yield of product is 82%, is_____g (nearest integer).

(Given molar mass in g mol^{-1} H:1, C:12, N:14, O:16)

Ans. (10)

75. Consider A $\xrightarrow{k_1}$ B and C $\xrightarrow{k_2}$ D are two reactions. If the rate constant (k_1) of the A $\rightarrow$ B reaction can be expressed by the following equation $\log_{10} k = 14.34 - \frac{1.5\times10^4}{T/K}$ and activation energy of C $\rightarrow$ D reaction (Ea_2) is $\frac{1}{5}$th of the A $\rightarrow$ B reaction (Ea_1), then the value of (Ea_2) is _____kJ mol^{-1}. (Nearest Integer)

Ans. (57)



ALLEN

For Class 12th Pass Students

RISE. REPEAT.
RANK UP IN JEE



JOIN LEADER COURSE
JEE (Main+Adv.) 2027

26th Mar & 15th Apr

Know more

ALLEN ONLINE

Think JEE 2027 will be your best shot?

Join the **Leader Online Course!**

Win up to **90%** scholarship via **ASAT**



Enrol Now

ALLEN ONLINE

Get REAL exam practice for JEE Main 2026

with the

Major Online Test Series!

- 13 full-syllabus tests
- 100+ additional mock tests
- 50,000+ teacher-recommended Qs. & more



Buy Now

ALLEN

Get The Latest

IIT-JEE Special Books


at Your Door Steps...!!

JOIN THE JOURNEY OF LEARNING

with

HANDBOOKS | ADVANCED-QB | SCORE PAPERS

PHYSICS EXPRESS | MAIN PYQ's | Adv. PYQ's



SHOP NOW

Available in
HINDI & ENGLISH