JEE–MAIN EXAMINATION – APRIL 2026

(HELD ON SUNDAY 05th APRIL 2026) **TIME : 9:00 AM TO 12:00 NOON**

MATHEMATICS — TEST PAPER WITH SOLUTION

SECTION-A

1. Let a, b ∈ C. Let α, β be the roots of the equation $x^2 + ax + b = 0$. If $\beta-\alpha=\sqrt{11}$ and $\beta^2-\alpha^2=3i\sqrt{11}$, then $(\beta^3-\alpha^3)^2$ is equal to :

(1) 160 (2) 176

(3) 194 (4) 187

Ans. (2)

Sol. $\beta-\alpha=\sqrt{11}$ …(1)

and $\beta^2-\alpha^2=3i\sqrt{11}$

$\Rightarrow (\beta+\alpha)(\beta-\alpha)=3i\sqrt{11}$

$\Rightarrow \beta+\alpha=3i$ …(2)

from (1) and (2)

$$\beta=\frac{\sqrt{11}+3i}{2},\quad \alpha=\frac{3i-\sqrt{11}}{2}$$

$$\Rightarrow \alpha\beta=\frac{-9-11}{4} \Rightarrow \alpha\beta=-5$$

Now $(\beta^3-\alpha^3)^2=(\beta-\alpha)^2(\beta^2+\alpha^2+\alpha\beta)^2$

$=(\beta-\alpha)^2((\beta+\alpha)^2-\alpha\beta)^2$

$=(11)(-9+5)^2=176$

2. Let the sum of the first n terms of an A.P. be $3n^2 + 5n$. Then the sum of squares of the first 10 terms of the A.P. is :

(1) 10220 (2) 12860

(3) 15220 (4) 19780

Ans. (3)

Sol. $S_n = 3n^2 + 5n$

$T_n = S_n - S_{n-1}$

$\Rightarrow T_n = (3n^2 + 5n) - (3(n-1)^2) + 5(n-1))$

$\Rightarrow T_n = 6n + 2$

$$\sum_{n=1}^{10} T_n^2 = \sum_{n=1}^{10}(6n+2)^2$$

$$= 36\sum_{n=1}^{10} n^2 + 24\sum_{n=1}^{10} n + \sum_{n=1}^{10} 4$$

$= 15220$

3. Let A be a 3 × 3 matrix such that

$$A^T\begin{bmatrix}1\\0\\1\end{bmatrix}=\begin{bmatrix}5\\2\\2\end{bmatrix},\quad A^T\begin{bmatrix}0\\0\\1\end{bmatrix}=\begin{bmatrix}3\\1\\1\end{bmatrix},\quad A\begin{bmatrix}1\\0\\1\end{bmatrix}=\begin{bmatrix}3\\4\\4\end{bmatrix}$$ and

$$A\begin{bmatrix}0\\0\\1\end{bmatrix}=\begin{bmatrix}1\\3\\1\end{bmatrix}$$

If det(A) = 1, then det(adj(A^2 + A)) is equal to :

(1) 16 (2) 25

(3) 49 (4) 64

Ans. (4)

Sol. Let $A=\begin{pmatrix}\alpha_1 & \alpha_2 & \alpha_3\\ \beta_1 & \beta_2 & \beta_3\\ \gamma_1 & \gamma_2 & \gamma_3\end{pmatrix}$

Now $A=\begin{pmatrix}\alpha_1 & \beta_1 & \gamma_1\\ \alpha_2 & \beta_2 & \gamma_2\\ \alpha_3 & \beta_3 & \gamma_3\end{pmatrix}\begin{pmatrix}0\\0\\1\end{pmatrix}=\begin{pmatrix}3\\1\\1\end{pmatrix}$

$$\Rightarrow \begin{pmatrix}\gamma_1\\ \gamma_2\\ \gamma_3\end{pmatrix}=\begin{pmatrix}3\\1\\1\end{pmatrix} \quad \ldots(1)$$

Now $\begin{pmatrix}\alpha_1 & \alpha_2 & \alpha_3\\ \beta_1 & \beta_2 & \beta_3\\ \gamma_1 & \gamma_2 & \gamma_3\end{pmatrix}\begin{pmatrix}0\\0\\1\end{pmatrix}=\begin{pmatrix}1\\3\\1\end{pmatrix}$

$$\Rightarrow \begin{pmatrix}\alpha_3\\ \beta_3\\ \gamma_3\end{pmatrix}=\begin{pmatrix}1\\3\\1\end{pmatrix} \quad \ldots(2)$$

$$A^T\begin{pmatrix}1\\0\\1\end{pmatrix}-A^T\begin{pmatrix}0\\0\\1\end{pmatrix}=\begin{pmatrix}5\\2\\2\end{pmatrix}-\begin{pmatrix}3\\1\\1\end{pmatrix}$$

$$\Rightarrow A^T\begin{pmatrix}1\\0\\0\end{pmatrix}=\begin{pmatrix}2\\1\\1\end{pmatrix}$$

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

$$\Rightarrow \begin{pmatrix}\alpha_1\\ \alpha_2\\ \alpha_3\end{pmatrix}=\begin{pmatrix}2\\1\\1\end{pmatrix} \quad \ldots(3)$$

$$\text{Now } A\begin{pmatrix}1\\0\\1\end{pmatrix}-A\begin{pmatrix}0\\0\\1\end{pmatrix}=\begin{pmatrix}3\\4\\4\end{pmatrix}-\begin{pmatrix}1\\3\\1\end{pmatrix}$$

$$\Rightarrow A\begin{pmatrix}1\\0\\0\end{pmatrix}=\begin{pmatrix}2\\1\\3\end{pmatrix}$$

$$\Rightarrow \begin{pmatrix}\alpha_1\\ \beta_1\\ \gamma_1\end{pmatrix}=\begin{pmatrix}2\\1\\3\end{pmatrix} \quad \ldots(4)$$

$$A=\begin{pmatrix}2&1&1\\1&\beta_2&3\\3&1&1\end{pmatrix}$$

$|A| = 2(\beta_2 - 3) - 1(1-9) + 1(1-3\beta_2) = 1$

$\Rightarrow 2\beta_2 - 6 + 8 + 1 - 3\beta_2 = 1$

$\Rightarrow \beta_2 = 2$

$$\text{So } A=\begin{pmatrix}2&1&1\\1&2&3\\3&1&1\end{pmatrix}$$

$$A^2=\begin{pmatrix}2&1&1\\1&2&3\\3&1&1\end{pmatrix}\begin{pmatrix}2&1&1\\1&2&3\\3&1&1\end{pmatrix}=\begin{pmatrix}8&5&6\\13&8&10\\10&6&7\end{pmatrix}$$

$$A^2+A=\begin{pmatrix}10&6&7\\14&10&13\\13&7&8\end{pmatrix} \Rightarrow \det(A^2+A)=8$$

$\Rightarrow \text{Det}(\text{adj}(A^2+A)) = 8^2 = 64$

4. Consider the system of linear equations in x, y, z :

$x + 2y + tz = 0,$

$6x + y + 5tz = 0,$

$3x + t^2y + f(t)z = 0,$

Where $f : R \to R$ is a differentiable function. If this system has infinitely many solutions for all $t \in R$, then f

(1) is a constant function

(2) is strictly increasing on R

(3) is strictly decreasing on R

(4) has two critical points

Ans. (2)

Sol. $$D=\begin{vmatrix}1&2&t\\6&1&5t\\3&t^2&f(t)\end{vmatrix}=0$$

$\Rightarrow 1(f(t) - 5t^3) - 2(6f(t) - 15t) + t(6t^2 - 3) = 0$

$$f(t)=\frac{t^3+12t}{11}$$

$$f'(t)=\frac{1}{11}(3t^2+12)>0 \quad \forall\, t \in R$$

$\Rightarrow f(t)$ is strictly increasing on R

5. $\displaystyle\sum_{n=1}^{10}\left(\frac{528}{n(n+1)(n+2)}\right)$ is equal to :

(1) 65 (2) 130

(3) 220 (4) 440

Ans. (2)

Sol. Let $\displaystyle\sum_{n=1}^{10}\frac{528}{n(n+1)(n+2)}=\lambda$

Let $T_n=\dfrac{1}{n(n+1)(n+2)}$

$$T_n=\frac{1}{2}\left\{\frac{(n+2)-n}{n(n+1)(n+2)}\right\}$$

$$T_n=\frac{1}{2}\left\{\frac{1}{n(n+1)}-\frac{1}{(n+1)(n+2)}\right\}$$

$$T_1=\frac{1}{2}\left\{\frac{1}{1.2}-\frac{1}{2.3}\right\}$$

$$T_2=\frac{1}{2}\left\{\frac{1}{2.3}-\frac{1}{3.4}\right\}$$

$\vdots$

$$T_{10}=\frac{1}{2}\left\{\frac{1}{10.11}-\frac{1}{11.12}\right\}$$

$$S_n=\frac{1}{2}\left[\frac{1}{2}-\frac{1}{132}\right]$$

$$S_n=\frac{1}{2}\left[\frac{66-1}{132}\right]=\frac{65}{264}$$

$$\text{So, } \lambda=\frac{65}{264}\times 528$$

$\lambda = 130$

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

6. Let tan A, tan B, where $A, B \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, be the roots of the quadratic equation $x^2 - 2x - 5 = 0$. Then $20\sin^2\left(\frac{A+B}{2}\right)$ is equal to :

(1) $10+\sqrt{10}$ (2) $10-2\sqrt{10}$

(3) $10-3\sqrt{10}$ (4) $10-\sqrt{10}$

Ans. **(3)**

Sol. $x^2 - 2x - 5 = 0$

$\tan A + \tan B = 2; \quad \tan A \tan B = -5$

$\therefore \tan(A+B) = \frac{2}{1-(-5)} = \frac{1}{3}$

$\Rightarrow \cos(A+B) = \frac{3}{\sqrt{10}}$

$\therefore 20\left(\sin^2\left(\frac{A+B}{2}\right)\right) = \frac{10}{2}(1-\cos(A+B))$

$= 10\left(1-\frac{3}{\sqrt{10}}\right) = \left(10-3\sqrt{10}\right)$

7. A letter is known to have arrived by post either from KANPUR or from ANANTPUR. On the the envelope just two consecutive letters AN are visible. The probability, that the letter came from ANANTPUR, is :

(1) $\frac{7}{10}$ (2) $\frac{10}{17}$

(3) $\frac{12}{19}$ (4) $\frac{7}{19}$

Ans. **(2)**

Sol. $P(I) : \frac{1}{2}\ P(A/I) \to \frac{1}{5}$

$P(II) : \frac{1}{2}\ P(A/II) \to \frac{2}{7}$

$\therefore \text{Reqd. Prob} = \frac{\frac{1}{2}\times\frac{2}{7}}{\frac{1}{2}\times\frac{1}{5}+\frac{1}{2}\times\frac{2}{7}} = \frac{\frac{2}{7}\times 5}{\frac{17}{7}} = \frac{10}{17}$

8. The mean deviation about the mean for the data

x_i	5	7	9	10	12	15
f_i	8	6	2	2	2	6

is equal to :

(1) $\frac{40}{13}$ (2) $\frac{42}{13}$

(3) $\frac{44}{13}$ (4) $\frac{46}{13}$

Ans. **(3)**

Sol. $\text{M.D.} = \frac{\sum f_i |x_i - \bar{x}|}{\sum f_i}$

$\bar{x} = \frac{\sum f_i x_i}{\sum f_i} = \frac{40+42+18+20+24+90}{26}$

$= \frac{234}{26} = 9$

$\text{M.D.} = \frac{8(4)+6(2)+2(0)+2(1)+2(3)+6(6)}{26}$

$= \frac{84}{26} = \frac{44}{13}$

9. Let a focus of the ellipse E : $\frac{x^2}{a^2}+\frac{y^2}{b^2}=1$ be S(4, 0) and its eccentricity be $\frac{4}{5}$. If the point P(3, α) lies on E and O is the origin, then the area of ΔPOS is equal to :


(1) 12/5 (2) 14/5

(3) 24/5 (4) 48/5

Ans. **(3)**

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol.



$a > b$

Focus $(ae, 0) = (4, 0)$

$\therefore ae = 4$

$a\left(\frac{4}{5}\right) = 4 \Rightarrow a = 5$

$\because b^2 = a^2(1 - e^2)$

$b^2 = 25\left(1 - \frac{16}{25}\right)$

$\therefore E : \frac{x^2}{25} + \frac{y^2}{9} = 1$

$\because$ P lies on the ellipse E

$\therefore \frac{9}{25} + \frac{\alpha^2}{9} = 1$

$\therefore \alpha = \pm\frac{12}{5}$

$\therefore$ Area of $\Delta_{POS} = \frac{1}{2}(OS)(PN)$

$= \frac{1}{2} \times 4 \times \frac{12}{5} = \frac{24}{5}$

10. Let P be a moving point on the circle

$$x^2 + y^2 - 6x - 8y + 21 = 0$$

Then, the maximum distance of P from the vertex of the parabola $x^2 + 6x + y + 13 = 0$ is equal to :

(1) 8 (2) 10


(3) 12 (4) 9

Ans. (3)

Sol. Centre : C(3,4) & r = 2

Parabola $(x + 3)^2 = -(y + 4)$

Vertex is A(–3,–4)



$AP_{max} = AC + r = \sqrt{36 + 64} + 2 = 12$


11. In an equilateral triangle PQR, let the vertex P be at (3, 5) and the side QR be along the line $x + y = 4$. If the orthocentre of the triangle PQR is (α, β), then $9(\alpha + \beta)$ is equal to :

(1) 16 (2) 27

(3) 36 (4) 48

Ans. (4)

Sol.



$Q : \frac{x-3}{1} = \frac{y-5}{1} = -\frac{(3+5-4)}{1+1}$

Q(1, 3)

So orthocentre is $\left(\frac{2+3}{3}, \frac{6+5}{3}\right)$

$\left(\frac{5}{3}, \frac{11}{3}\right)$

So $9(\alpha + \beta) = 3(16) = 48$

12. The sum of all the integral values of p such that the equation $3\sin^2 x + 12\cos x - 3 = p$, $x \in R$, has at least one solution, is :

(1) –54 (2) –60

(3) –75 (4) –84

Ans. (3)

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. $P = 12\cos x - 3\cos^2 x$

$P = -3(\cos^2 x - 4\cos x)$

$P = -3((\cos x - 2)^2 - 2)$

put $\cos x = -1 \Rightarrow P = -15$

put $\cos x = 1 \Rightarrow P = 9$

$-15 \le P \le 9$

Sum of all integers $= -(10 + 11 + ... + 15) = -75$

13. The square of the distance of the point P(5, 6, 7) from the line $\frac{x-2}{2} = \frac{y-5}{3} = \frac{z-2}{4}$ is equal to :

(1) 3 (2) 5


(3) 6 (4) 8

Ans. **(3)**

Sol. $\overrightarrow{PQ} \cdot \vec{A} = 0$

$\Rightarrow (2\lambda - 3)2 + (3\lambda - 1)3 + (4\lambda - 5)4 = 0$

$\Rightarrow \lambda = 1$



Point Q = (4, 8, 6)

$PQ^2 = 6$

14. Let $\vec{a} = \sqrt{7}\hat{i} + \hat{j} - \hat{k}$ and $\vec{b} = \hat{j} + 2\hat{k}$. If $\vec{r}$ is a vector such that $\vec{r} \times \vec{a} + \vec{a} \times \vec{b} = \vec{0}$ and $\vec{r} \cdot \vec{a} = 0$, then $|3\vec{r}|^2$ is equal to :

(1) 44 (2) 54

(3) 86 (4) 132

Ans. **(1)**

Sol. $\vec{r} \times \vec{a} - \vec{b} \times \vec{a} = \vec{0}$

$(\vec{r} - \vec{b}) \times \vec{a} = \vec{0}$

$\vec{r} - \vec{b} = \lambda\vec{a}$

$\vec{r} = \vec{b} + \lambda\vec{a}$

$\vec{r}.\vec{a} = 0 \Rightarrow \vec{a}.\vec{b} + \lambda|\vec{a}|^2 = 0$

$\lambda = -\frac{\vec{a}.\vec{b}}{|\vec{a}|^2} = -\frac{(1-2)}{9} = \frac{1}{9}$

$\vec{r} = \vec{b} + \frac{\vec{a}}{9}$

$|3\vec{r}|^2 = 9|\vec{r}|^2 = 9\left(b^2 + \frac{a^2}{81} + \frac{2(\vec{a}.\vec{b})}{9}\right) = 44$

15. The square of the distance of the point of intersection of the lines $\vec{r} = \left(\hat{i} + \hat{j} - \hat{k}\right) + \lambda(a\hat{i} - \hat{j})$, $a \neq 0$ and $\vec{r} = \left(4\hat{i} - \hat{k}\right) + \mu(2\hat{i} + a\hat{k})$ from the origin is :

(1) 5 (2) 10

(3) 17 (4) 26

Ans. **(3)**

Sol. Equating both the equations

$\left(\hat{i} + \hat{j} - \hat{k}\right) + \lambda\left(a\hat{i} - \hat{j}\right) = \left(4\hat{i} - \hat{k}\right) + \mu\left(2\hat{i} + a\hat{k}\right)$

$1 + a\lambda = 4 + 2\mu$

$1 - \lambda = 0 \Rightarrow \lambda = 1$

$-1 = -1 + a\mu$

$a\mu = 0 \Rightarrow \mu = 0$

$a = 3$

point of intersection $= \hat{i} + \hat{j} - \hat{k} + 3\hat{i} - \hat{j} = 4\hat{i} - \hat{k}$

Ans. 17


16. The area of the region

$R = \{(x,y): xy \le 27, 1 \le y \le x^2\}$ is equal to :

(1) $78\log_e 3 - \frac{52}{3}$ (2) $54\log_e 3 - \frac{52}{3}$

(3) $54\log_e 3 - \frac{26}{3}$ (4) $54\log_e 3 + \frac{26}{3}$

Ans. **(2)**



Area $= \int_1^3 (x^2 - 1)dx + \int_3^{27}\left(\frac{27}{x} - 1\right)dx$

$= \left[\frac{x^3}{3} - x\right]_1^3 + [27\ell nx - x]_3^{27}$

$= \frac{26}{3} - 2 + 27\,\ell n\,9 - 24$

$= 27\,\ell n\,9 - \frac{26 \times 2}{3}$

$= 54\,\ell n\,3 - \frac{52}{3}$

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

17. The product of all possible values of α, for which

$$\lim_{x\to 0}\left(\frac{1-\cos(\alpha x)\cos((\alpha+1)x)\cos((\alpha+2)x)}{\sin^2((\alpha+1)x)}\right)=2,$$

is :

(1) –2 (2) 1

(3) –1 (4) $\frac{5}{4}$

Ans. (3)

Sol. $\lim_{x\to 0}\frac{1-\cos\alpha x.\cos((\alpha+1)x)\cos((\alpha+2)x)}{(\alpha+1)^2x^2}$

By using L.H. Rule we get

$\Rightarrow \frac{1}{2}(\alpha+1)^2[\alpha^2+(\alpha+1)^2+(\alpha+2)^2]=2$

$\alpha^2+(\alpha+1)^2+(\alpha+2)^2=4(\alpha+1)^2$

$\alpha^2+(\alpha+2)^2=3(\alpha+1)^2$

$\Rightarrow \alpha^2+2\alpha-1=0$

Product = –1

18. The value of the integral $\int_0^\infty \frac{\log_e(x)}{x^2+4}dx$ is:

(1) $\frac{\pi\log_e(2)}{2}$ (2) $\frac{\pi\log_e(2)}{4}$

(3) $1+\pi\log_e(2)$ (4) $2+\pi\log_e(2)$

Ans. (2)

Sol. Put $x=2t \Rightarrow dx=2dt$

$$I=\int_0^\infty \frac{\ell n2t}{4t^2+4}(2dt)=\frac{1}{2}\int_0^\infty \frac{\ell n2+\ell nt}{t^2+1}dt$$

$$=\frac{1}{2}\int_0^\infty \frac{\ell n2}{t^2+1}dt+\frac{1}{2}\int_0^\infty \frac{\ell nt}{t^2+1}dt$$

$$\left[\frac{\ln 2}{2}\tan^{-1}t\right]_0^\infty+I_1$$

$$=\frac{\ell n2}{2}\cdot\frac{\pi}{2}+I_1$$

$$I_2=\frac{1}{2}\int_0^\infty \frac{\ell nt}{t^2+1}dt \qquad t=\frac{1}{u}\Rightarrow dt=-\frac{du}{u^2}$$

$$=\frac{1}{2}\int_\infty^0 \frac{\ell n(1/u)}{\frac{1}{u^2}+1}\left(-\frac{du}{u^2}\right)$$

Add $\Rightarrow I_1=0$

$$I=\frac{\pi\ell n2}{4}$$

19. Let $f: R \to R$ be a differentiable function such that $f\left(\frac{x+y}{3}\right)=\frac{f(x)+f(y)}{3}$ for all $x, y \in R$ and $f'(0) = 3$. Then the minimum value of the function $g(x) = 3 + e^x f(x)$, is

(1) $3\left(\frac{e+1}{e}\right)$ (2) $3\left(\frac{e-1}{e}\right)$

(3) $\frac{3-e}{e}$ (4) 3e

Ans. (2)

Sol. $f\left(\frac{x+y}{3}\right)=\frac{f(x)+f(y)}{3}$, put $x = y = 0$

$f(0)=\frac{2f(0)}{3}$

$\Rightarrow f(0) = 0$.....(1)

$f'\left(\frac{x+y}{3}\right)\cdot\frac{1}{3}=\frac{1}{3}f'(x)$

put $x = 0$

$f'\left(\frac{y}{3}\right)\frac{1}{3}=\frac{1}{3}\times 3$

$f'\left(\frac{y}{3}\right)=3$

put $y = 3x$

$f'(x) = 3$

integrate both sides

$f(x) = 3x + c$

from eq. (1)

$f(0) = 0$

$\Rightarrow f(x) = 3x$

Now

$g(x) = 3 + e^x \cdot 3x$

$g'(x) = 3[e^x + x \cdot e^x]$

$= 3e^x(x + 1)$

$g'(x) = 0$, at $x = -1$

$(g(x))_{min} = 3 + e^{-1}(-3)$

$=3\left[1-\frac{1}{e}\right]=\frac{3(e-1)}{e}$

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

20. The value of the integral $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}}\left(\frac{4-\operatorname{cosec}^2 x}{\cos^4 x}\right)dx$ is :

(1) $\frac{11}{\sqrt{3}}$ (2) $\frac{16}{\sqrt{3}}$

(3) $\frac{32}{3\sqrt{3}}$ (4) $\frac{64}{3\sqrt{3}}$

Ans. (3)

Sol. $=\int_{\pi/6}^{\pi/3}\frac{4}{\cos^4 x}dx - \int_{\pi/6}^{\pi/3}\frac{\operatorname{cosec}^2 x}{\cos^4 x}dx$

$=\int_{\pi/6}^{\pi/3}\frac{4}{\cos^4 x}dx - \left[-\frac{\cot x}{\cos^4 x} - \int_{\pi/6}^{\pi/3}(-\cot x)\frac{-4}{\cos^5 x}(-\sin x)\,dx\right]$

$=\frac{\cot x}{\cos^4 x}\Big|_{\frac{\pi}{6}}^{\frac{\pi}{3}}$

$=\frac{1/\sqrt{3}}{(1/2)^4} - \frac{\sqrt{3}}{\left(\frac{\sqrt{3}}{2}\right)^4}$

$=\frac{16}{\sqrt{3}} - \frac{\sqrt{3}.16}{9} = \frac{16}{\sqrt{3}} - \frac{16}{3\sqrt{3}} = \frac{16}{\sqrt{3}}\left(1-\frac{1}{3}\right)$

$\frac{32}{3\sqrt{3}} = \frac{32\sqrt{3}}{9}$

SECTION-B

21. Let A = {1, 2, 3, 4, 5, 6}. The number of one-one functions f : A → A such that f(1) ≥ 3, f(3) ≤ 4 and f(2) + f(3) = 5, is _____.

Ans. (72)

Sol. f(2) + f(3) = 5

C-I : f(2) = 4, f(3) = 1

so, f(1) = 3, 5, 6 → 3 choices

f(4) → 3 choices

f(5) → 2 choices

f(6) → 1 choices

so total function for C-I : 3 × 3 × 2 × 1 = 18

Similarly

C-II : f(2) = 1, f(3) = 4 → 18 function

C-III : f(2) = 3, f(3) = 2 → 18 function

C-IV : f(2) = 2, f(3) = 2 → 18 function

Answer total 72 function

22. Two players A and B play a series of games of badminton. The player, who wins 5 games first, wins the series. Assuming that no game ends in a draw, the number of ways, in which player A wins the series is _____.

Ans. (126)

Sol. Maximum number of games will be 9

A must win 5 gamese so,

Number of ways = ${}^{9}C_{5}$ = 126

Answer = 126

23. If the sum of the coefficients of x^7 and x^{14} in the expansion of $\left(\frac{1}{x^3}-x^4\right)^n$, $x \neq 0$, is zero, then the value of n is _____.

Ans. (21)

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. $T_{r+1} = {}^nC_r \left(\frac{1}{x^3}\right)^{n-r} \left(-x^4\right)^r$

$T_{r+1} = {}^nC_r\, x^{7r-3n} (-1)^r$

for coeff. of $x^7 \Rightarrow 7r_1 - 3n = 7$

$\Rightarrow r_1 = \frac{7+3n}{7}$

coeff. of $x^7 \Rightarrow {}^nC_{r_1} (-1)^{r_1}$

for coeff. of $x^{14} \Rightarrow 7r_2 - 3n = 14$

$\Rightarrow r_2 = \frac{14+3n}{7} = r_1 + 1$

coeff. of $x^{14} \Rightarrow {}^nC_{r_1+1} (-1)^{r_1+1}$

$\Rightarrow {}^nC_{r_1} (-1)^{r_1} + {}^nC_{r_1+1} (-1)^{r_1+1} = 0$

$\Rightarrow {}^nC_{r_1} = {}^nC_{r_1+1} \Rightarrow r_1 + r_1 + 1 = n$

$\Rightarrow 2r_1 + 1 = n \Rightarrow 2\left(\frac{7+3n}{7}\right) + 1 = n$

Ans. $\Rightarrow \boxed{n = 21}$

24. If $\frac{\pi}{4} + \sum_{p=1}^{11} \tan^{-1}\left(\frac{2^{p-1}}{1+2^{2p-1}}\right) = \alpha$, then $\tan\alpha$ is equal to _____.

Ans. (2048)

Sol. $\alpha = \frac{\pi}{4} + \sum_{p=1}^{11} \tan^{-1}\left(\frac{2^{p-1}}{1+2^{2p-1}}\right)$

$= \frac{\pi}{4} + \sum_{p=1}^{11} \tan^{-1}\left(\frac{2^p - 2^{p-1}}{1+2^p 2^{p-1}}\right)$

$= \frac{\pi}{4} + \sum_{p=1}^{11} \left(\tan^{-1}\left(2^p\right) - \tan^{-1}\left(2^{p-1}\right)\right)$

$= \frac{\pi}{4} + \tan^{-1}\left(2^{11}\right) - \tan^{-1}\left(2^0\right) = 2^{11}$

Ans. 2048

25. Let $y = y(x)$ be the solution of the differential equation

$x \sin\left(\frac{y}{x}\right) dy = \left(y \sin\left(\frac{y}{x}\right) - x\right) dx$, $y(1) = \frac{\pi}{2}$ and let $\alpha = \cos\left(\frac{y\left(e^{12}\right)}{e^{12}}\right)$. Then the number of integral value of p, for which the equation $x^2 + y^2 - 2px + 2py + \alpha + 2 = 0$ represents a circle of radius $r \le 6$, is _____.

Ans. (11)

Sol. $\sin\frac{y}{x}\frac{dy}{dx} = \frac{y}{x}\sin\frac{y}{x} - 1$

put $y = tx \Rightarrow \frac{dy}{dx} = t + x\frac{dt}{dx}$

$\sin t\left(t + x\frac{dt}{dx}\right) = t\sin t - 1$

$x\sin t\frac{dt}{dx} + 1 = 0$

$\sin t\, dt + \frac{dx}{x} = 0$

$\Rightarrow -\cos t + \ell nx = C$

$\Rightarrow -\cos\left(\frac{y}{x}\right) + \ell nx = C$

$y(1) = \frac{\pi}{2} \Rightarrow C = 0$

$\Rightarrow \cos\left(\frac{y}{x}\right) = \ell nx$

$\operatorname{cosec}\left(\frac{y(e^{12})}{e^{12}}\right) = 12 \Rightarrow \alpha = 12$

$x^2 + y^2 - 2px + 2py + 14 = 0$

$r = \sqrt{p^2 + p^2 - 14}$

$\Rightarrow r \le 6 \Rightarrow r^2 \le 36$

$2p^2 - 14 \le 36$

$p^2 \le 25$

$p \in [-5, 5]$

number of integral value of p = 11

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

भरोसा जताया नहीं, जीता जाता है

जब नतीजे मायने रखते हैं,
तो स्टूडेंट्स **ALLEN** KOTA को चुनते हैं

JOIN
LEADER COURSE
For Class 12th pass students

Target: JEE (Main+Adv.) 2027

Batches Starting from

15 April &
06, 27 May 2026

Click to know more



AIR 1
JEE Adv. 2025
Rajit Gupta
7 Year Classroom Course

ALLEN

ALLEN Scholarship Admission Test (ASAT)

Win up to
90% Scholarship*
On JEE, NEET &
Foundation Courses



Test Dates: 5, 12, 19 & 26 April

Register for ~~500~~ ₹99/-

*Limited period offer, subject to the scholarship rules and the T&Cs.

ALLEN ONLINE

Ace **JEE 2027** with our

Leader Online Course

- **900+** hours of LIVE classes
- **30** latest pattern tests
- **24/7** AI-powered doubt support & more



Avail up to a

90% **scholarship** with **ASAT.**

Enrol Now

ALLEN ONLINE

Prefer learning at your own pace?

Start anytime with the **Leader Self-Study PLUS Course**

- **2000+** recorded video lectures
- **32** latest-pattern tests
- Digital study material & more




Enrol Now

DON'T PREPARE HARDER
PREPARE RIGHT !

Advanced-level MCQs

Prepared By ALLEN Experts



Well Graded Exercise

Detailed Answer Key

SHOP NOW on ALLEN e-Store


ALLEN

Get The Latest
IIT-JEE Special Books
at Your Door Steps...!!

JOIN THE JOURNEY OF LEARNING

with

HANDBOOKS | ADVANCED-QB | SCORE PAPERS
PHYSICS EXPRESS | MAIN PYQ's | Adv. PYQ's



SHOP NOW

Available in HINDI & ENGLISH