JEE–MAIN EXAMINATION – APRIL 2026

(HELD ON SUNDAY 05th APRIL 2026) **TIME : 3:00 PM TO 6:00 PM**

CHEMISTRY — TEST PAPER WITH SOLUTION

SECTION-A

51. What volume of hydrogen gas at STP would be liberated by action of 50 mL of H_2SO_4 of 50% purity (density = 1.3 g mL^{-1}) on 20 g of zinc ?

Given : Molar mass of H, O, S, Zn are 1, 16, 32, 65 g mol^{-1} respectively.

(1) 5.824 L (2) 7.428 L

(3) 6.892 L (4) 8.375 L

Ans. (3)

Sol. Mass of $H_2SO_4 = \frac{50\times1.3}{98}\times\frac{50}{100} = \frac{32.5}{98}$

$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq.) + H_2(g)$

$\frac{20}{65}$ $\frac{32.5}{98}$

Moles of H_2 formed = $\frac{20}{65}$

Volume of $H_2 = \frac{20}{65} \times 22.7 = 6.9$ L.

52. Which of the following statement(s) is/are **true** ?

(A) If two orbitals have the same value of $(n + l)$, the orbital with lower value of n will have lower energy.

(B) Energies of the orbitals in the same subshell increase with increase in atomic number.

(C) The size of $2p_x$ orbital is less than the size of $3p_x$ orbital.

(D) Among 5f,6s, 4d, 5p and 5d orbitals, none of the orbitals have 2 radial nodes.

Choose the **correct** answer from the options given below :

(1) A, B and C only (2) A and C only

(3) C and D only (4) A only

Ans. (2)

Sol. (a) If two orbitals are having same value of "$n+\ell$" then the orbital having lower value of 'n' will have lower energy.

(b) Energies of the orbitals in the same subshell decreases with increase in atomic number.

(c) Size of $2p_x < 3p_x$

(d)

Orbital	Radial node (n-ℓ-1)
5f	1
6s	5
4d	1
5p	3
5d	2

53. The covalent radii of atoms A and B are r_A and r_B, respectively. The covalent bond length and total length of AB molecule are respectively :

(1) $(r_A + r_B)$, $2(r_A + r_B)$ (2) $\frac{1}{2}(r_A + r_B)$, $(r_A + r_B)$

(3) $(r_A + r_B)$, $(r_A + r_B)$ (4) $2(r_A + r_B)$, $\frac{1}{2}(r_A + r_B)$

Ans. (1)

Sol. According to NCERT

Bond length = $r_A + r_B$

Total length of molecule = $2(r_A + r_B)$

54. Consider the following data for the reaction

$X_2(g) + Y_2(g) \rightleftharpoons 2XY(g)$

at 600 K. The $\Delta_rG^\ominus$ (in kJ mol^{-1}) for the reaction is:

Compound	$\Delta_fH^\ominus_{600K}$ (kJ mol^{-1})	$S^\ominus_{600K}$ (J $mol^{-1}K^{-1}$)
XY(g)	42	200
$X_2(g)$	8	140
$Y_2(g)$	80	250

(1) –21000 (2) –10

(3) –1000 (4) –9.012

Ans. (2)

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. $\Delta_rG = \Delta_rH - T.\Delta_rS$

$\Delta_rH = (2 \times 42 - 80 - 8) = -4$ kJ/mol

$\Delta_rS = (400 - 250 - 140) = +10$ J/K.mol

$\Delta_rG = -4000 - 600 \times (10)$

$= -10{,}000$ J/mol $= -10$ kJ/mol

55. The **correct** order of molar heat capacities measured at 298 K and 1 bar is :

(1) Copper (s) > Bromine (l) > Helium (g)

(2) Bromine (l) > Copper (s) > Helium (g)

(3) Helium (g) > Bromine (l) > Copper (s)

(4) Helium (g) > Bromine (l) = Copper (s)

Ans. (2)

Sol. Generally molar heat capacity of liquids is greater than solids because liquids have greater degrees of freedom & thus can store more energy with lesser temperature rise.

Also for monoatomic gases molar heat capacity is low as only transalational degrees of freedom are present.

$\therefore$ order is :

$Br_2(\ell) > Cu(s) > He(g)$

56. The reaction $A(g) \rightleftharpoons B(g) + C(g)$ was initiated with the amount 'a' of A(g). At equilibrium it is found that the amount of A(g) remaining is (a –x) at a total pressure of p.

The equilibrium constant K_p of the reaction can be calculated from the expression :

(1) $\frac{x^2}{a^2+x^2} \times p$ (2) $\frac{x^2}{a^2-x^2} \times p$

(3) $\frac{a+x^2}{x^2} \times p$ (4) $\frac{a-x^2}{x^2} \times p$

Ans. (2)

Sol.

	$A(g)$	$\rightleftharpoons B(g)$	$+ C(g)$
t = 0	a	0	0
t_{eq}	a–x	x	x

$$K_p = \frac{(x)(x)}{(a-x)} \times \left(\frac{P}{a+x}\right)$$

$$K_P = \frac{x^2P}{a^2-x^2}$$

57. One half cell in a voltaic cell is constructed by dipping silver rod in $AgNO_3$ solution of unknown concentration, other half cell is Zn rod dipped in 1 molar solution of $ZnSO_4$.

A voltage of 1.60 V is measured at 298 K for this cell. What is the concentration of Ag^+ ions used in terms of log x($x = [Ag^+]$) ?

$E^{\Theta}_{Zn^{2+}/Zn} = -0.76V$, $E^{\Theta}_{Ag^+/Ag} = +0.80V$,

$\frac{2.303RT}{F} = 0.059$ V

(1) $\frac{2}{3.9}$ (2) $\frac{4}{5.9}$

(3) $\frac{2.9}{2}$ (4) $\frac{5.9}{4}$

Ans. (2)

Sol. Cell Reaction :

$Zn(s) + 2Ag^{\oplus}(aq) \longrightarrow 2Ag(s) + Zn^{+2}(aq)$

$E_{cell} = E^{o}_{cell} - \frac{0.059}{n} \log(Q)$

$1.6 = 1.56 - \frac{0.059}{2} \log \frac{1}{(Ag^+)^2}$

$0.04 = +0.059 \log[Ag^+]$

$\text{Log}[Ag^+] = \frac{4}{5.9}$

58. Given below are two statements :

Statement-I : The number of pairs among $[Al_2O_3, Cr_2O_3]$, $[Cl_2O_7, Mn_2O_7]$, $[Na_2O, V_2O_3]$ and $[CO, N_2O]$ that contain oxides of same nature (acidic, basic, neutral or amphoteric) is 4.

Statement-II : Among Na_2O, Al_2O_3, CO and Cl_2O_7, the most basic and acidic oxides are Na_2O and Cl_2O_7, respectively.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both **Statement-I** and **Statement-II** are true.

(2) Both **Statement-I** and **Statement-II** are false.

(3) **Statement-I** is true but **Statement-II** is false.

(4) **Statement-I** is false but **Statement-II** is true.

Ans. (1)

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. **Statement I :**

Al_2O_3, Cr_2O_3 : Amphoteric

CO, N_2O : Neutral

Na_2O, V_2O_3 : Basic

Cl_2O_7, Mn_2O_7 : Acidic

Statement II :

Most basic oxide : Na_2O

Most acidic oxide : Cl_2O_7

59. Given below are two statements :

Statement-I : Aluminium upon reaction with NaOH forms $[Al(OH)_6]^{3-}$ ion.

Statement-II : The geometry of ICl_4^-, ClO_3^- and IBr_2^- is square planar, pyramidal and linear respectively.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both **Statement-I** and **Statement-II** are true.

(2) Both **Statement-I** and **Statement-II** are false.

(3) **Statement-I** is true but **Statement-II** is false.

(4) **Statement-I** is false but **Statement-II** is true.

Ans. **(4)**

Sol. **Statement I :**

$Al + NaOH\ (excess) \rightarrow Na[Al(OH)_4] + H_2(g)$

Statement II :

$ClO_3^{\ominus}$ is sp^3 and pyramidal

$ICl_4^{\ominus}$ is sp^3d^2 and square planar

$IBr_2^{\ominus}$ is sp^3d and linear.

60. Given below are two statements :

Statement-I : Presence of large number of unpaired electrons in transition metal atoms results in higher enthalpies of their atomisation.

Statement-II : $d_{xy} = d_{xz} = d_{yz} < d_{x^2-y^2} = d_{z^2}$ and $d_{x^2-y^2} = d_{z^2} < d_{xy} = d_{xz} = d_{yz}$ are the d-orbital splittings in $[Fe(H_2O)_6]^{3+}$ and $[Ni(Cl)_4]^{2-}$ complex ions respectively.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both **Statement-I** and **Statement-II** are correct.

(2) Both **Statement-I** and **Statement-II** are incorrect.

(3) **Statement-I** is correct but **Statement-II** is incorrect.

(4) **Statement-I** is incorrect but **Statement-II** is correct.

Ans. **(1)**

Sol. **Statemet I :**

More are the number of unpaired electrons , stronger is metal-metal boding thus higher enthalpy of atomisation.

Statement II :

$[Fe(H_2O)_6]^{3+}$ is octahedral complex.

Thus, $\left[t_{2g}\left(d_{xy} = d_{yz} = d_{zx}\right) < e_g\left(d_{x^2-y^2} = d_{z^2}\right)\right]$

$[NiCl_4]^{2-}$ is tetrahedral complex.

Thus, $\left[e\left(d_{x^2-y^2} = d_{z^2}\right) < t_2\left(d_{xy} = d_{yz} = d_{zx}\right)\right]$

61. Identify the **correct** statements from the following:

(A) $[Fe(C_2O_4)_3]^{3-}$ is the most stable complex among $[Fe(OH)_6]^{3-}$, $[Fe(C_2O_4)_3]^{3-}$ and $[Fe(SCN)_6]^{3-}$

(B) The stability of $[Cu(NH_3)_4]^{2+}$ is greater than that of $[Cu(en)_2]^{2+}$

(C) The hybridization of Fe in $K_4[Fe(CN)_6]$ is d^2sp^3

(D) $[Fe(NO_2)_3Cl_3]^{3-}$ exhibits linkage isomerism

(E) NO_2^- and SCN^- ligands are NOT ambidentate ligands

Choose the **correct** answer from the options given below :

(1) A, B, C, D and E (2) B, C and D only

(3) A, C and D only (4) A, C and E only

Ans. **(3)**

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. (A) $[Fe(C_2O_4)_3]^{3-}$ has oxalate ion as ligand which leads to chelation thus maximum stability among given complexes.

(B) Stability : $[Cu(NH_3)_4]^{2+} < [Cu(en)_2]^{2+}$; due to higher CFSE and chelation.

(C) $K_4[Fe(CN)_6]$

$\Rightarrow Fe^{2+} = [Ar]3d^6$

In presence of SFL the electronic configuration of Fe^{+2} $[t_{2g}^6, e_g^0]$, thus d^2sp^3 hybridized.

(D) $[Fe(NO_2)_3Cl_3]^{3-}$ has NO_2^- as ambidentate ligand thus can show linakge isomerism.

(E) NO_2^- and SCN^- both are ambidentate ligands.

62. Match **List-I** with **List-II.**

	List-I (Purification technique)		**List-II (Used to separate)**
A.	Simple distillation	I.	Steam volatile compound
B.	Fractional distillation	II.	Two liquids with large difference in boiling points
C.	Steam distillation	III.	Liquid decomposing at its boiling point
D.	Distillation under reduced pressure	IV.	Two liquids with close boiling points

Choose the **correct** answer from the options given below :

(1) A-II, B-III, C-I, D-IV (2) A-II, B-IV, C-I, D-III

(3) A-II, B-IV, C-III, D-I (4) A-IV, B-III, C-II, D-I

Ans. **(3)**

Sol. It is theory based question.

63. IUPAC name of the some alkenes are given below :

Find out the correct stability order.

A. 2-Methylbut-2-ene

B. *cis*-But-2-ene

C. 2,3-Dimethylbut-2-ene

D. Prop-1-ene

Choose the **correct** answer from the options given below :

(1) C > A > B > D


(2) C > A > D > B

(3) B > D > A > C

(4) A > B > C > D

Ans. **(1)**

Sol.



Stability of alkene ∝ Number of α hydrogen.

64. Identify the correct IUPAC name of hydrocarbon (x) containing three primary carbon atoms and with molar mass 72 g mol^{-1}.

(1) 1,1-Dimethylcyclopropane


(2) 2,2-Dimethylpropane

(3) 2-Methylbutane


(4) n-pentane

Ans. **(3)**


Sol.



Mol. mass = 72



65. Complete the following reaction sequence and give the name of major product 'P'.



(1) 2-Chloropropanoic acid

(2) 3-Chloropropanoic acid

(3) 1-Chloropropane

(4) 2-Chloropropane

Ans. (1)

Sol. $CH_3–CH–C\equiv N \xrightarrow[\text{(hydrolysis of cyanide}]{OH^-/H_2O} CH_3–CH_2–COO^- \xrightarrow{H_3O^+} CH_3–CH_2–COOH \xrightarrow[\text{(HVZ Rxn)}]{\text{(i) Red P + } Cl_2,\ \text{(ii) } H_2O} CH_3–\underset{Cl}{CH}–COOH$

66. Given below are two statements :

Statement-I : The condensation reaction between $CH_3–CH=O$ and $H_2N–\underset{H}{N}–\underset{O}{C}–NH_2$ under optimum pH will produce $CH_3–CH=N–\overset{\overset{O}{\|}}{C}(H)–N–NH_2$

Statement-II : The molecule, $Ph–CH(O–H)(O–CH_3)$ will generate Ph–CH=O in the presence of dilute acid.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both **Statement-I** and **Statement-II** are true.


(2) Both **Statement-I** and **Statement-II** are false.

(3) **Statement-I** is true but **Statement-II** is false.

(4) **Statement-I** is false but **Statement-II** is true.


Ans. (4)

Sol. Statement-I



Hence statement-I is incorrect.

Statement-II



Statement-II is correct.

67. Given below are two statements :

Statement-I : Heating benzamide with bromine in an ethanolic solution of sodium hydroxide will give benzylamine.

Statement-II : Nitration of aniline with HNO_3/H_2SO_4 at 288 K produces *m*-nitroaniline in higher amount than *o*-nitroaniline (pH adjusted).

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both **Statement-I** and **Statement-II** are true.

(2) Both **Statement-I** and **Statement-II** are false.


(3) **Statement-I** is true but **Statement-II** is false.

(4) **Statement-I** is false but **Statement-II** is true.

Ans. (4)


ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. Statement-I



Hence Statement-I is incorrect

Statement-II



Statement-II is correct

68. Identify the **incorrect** statement about tertiary structure of proteins.

(1) They can be fibrous or globular in structure.

(2) The main forces that stabilize the structure are hydrogen bonding, disulphide links, van der Waals and electrostatic forces of attraction.


(3) The structure remains intact when exposed to pH changes.


(4) A linear polypeptide chain will convert to a secondary structure and then further folding of the secondary structure will convert to tertiary structure.

Ans. **(3)**

Sol. On changing pH tertiary strutcure of protein disrupted and amino acids intact only by peptide linkage.

69. Given below are two statements :

Statement-I :  and

 are two anomers of

D-(+)-glucose.

Statement-II : The open chain forms of D-glucose and D-fructose contain three similar chiral carbons at C_3, C_4 and C_5.

In the light of the above statements, choose the **correct** answer from the options given below :

(1) Both **Statement-I** and **Statement-II** are true.

(2) Both **Statement-I** and **Statement-II** are false.

(3) **Statement-I** is true but **Statement-II** is false.

(4) **Statement-I** is false but **Statement-II** is true.

Ans. **(1)**

Sol. α-D-Glucose and β-D-Glucose are anomers.

At C-3, C-4 and C-5 are identical configuration in glucose and fructose.

70. A paper dipped in a dil. H_2SO_4 solution of 'X' upon treatment with SO_2 gas turns into green. The compound 'X' is :

(1) KI-starch (2) $KMnO_4$

(3) $Pb(CH_3COO)_2$ (4) $K_2Cr_2O_7$

Ans. **(4)**

Sol. SO_2 gas acts as reducing agent.

$$K_2Cr_2O_7 + SO_2 \xrightarrow{H_2SO_4} \underset{\text{Green solution}}{Cr_2(SO_4)_3} + SO_4^{2-}$$

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

SECTION-B

71. The total number of unpaired electrons present in the d^3, d^4 (low spin) d^5 (high spin), d^6 (high spin) and d^7 (low spin) octahedral complex systems is ____.

Ans. **(15)**

Sol. $d^3 \rightarrow \left[t_{2g}^3, e_g^0\right]$, 3 unpaired electrons

d^4 (low spin) $\rightarrow \left[t_{2g}^4, e_g^0\right]$, 2 unpaired electrons

d^5 (high spin) $\rightarrow \left[t_{2g}^3, e_g^2\right]$, 5 unpaired electrons

d^6 (high spin) $\rightarrow \left[t_{2g}^4, e_g^2\right]$, 4 unpaired electrons

d^7 (low spin) $\rightarrow \left[t_{2g}^6, e_g^1\right]$, 1 unpaired electrons

72. RMgI when treated with ice cold water liberated a gas which occupied 1.4 dm^3/g at STP. The gas produced is further reacted with iodine in presence of HIO_3 to give compound (X). Compound (X) in presence of Na and dry ether produced compound (Y). Molar mass of compound (Y) is ___ $g\ mol^{-1}$. (Nearest integer)

Ans. **(30)**

Sol. $RMgI \xrightarrow[H_2O]{\text{Ice cold}} \underset{\text{Gas}}{R{-}H} + Mg\begin{matrix} I \\ OH \end{matrix}$

Vol. = 1.4 dm^3/gm at STP

Hence according to this mol. Mass of gas is = 16

Hence liberated gas is = CH_4 (Methane)

$$CH_4 + I_2 \xrightarrow{HIO_3} \underset{X}{CH_3{-}I} + HI$$

$$\underset{X}{CH_3 + I} \xrightarrow[\text{(Wurtz reaction)}]{\text{Na/Dry ether}} \underset{Y}{CH_3 - CH_3}$$

Molar mass = 30

73. 20 g hemoglobin in a 1 L aqueous solution (A) at 300 K is separated from pure water by semi permeable membrane. At equilibrium the height of solution in a tube dipped in a solution (A) is found to be 80.0 mm higher than the tube dipped in water.

The molar mass of hemoglobin is _____ $kg\ mol^{-1}$. (Nearest integer)

(Given : g = 10 ms^{-2}, R = 8.3 kPa $dm^3K^{-1}mol^{-1}$, density of solution = 1000 $kg\ m^{-3}$)

Ans. **(62)**

Sol. Osmotic pressure, $\pi = \rho gh$

$= 1000 \times 10 \times 80 \times 10^{-3}$

= 800 Pa .....(1)

Let molar mass of haemoglobin = M g/mol

conc. of haemoglobin = $\frac{20/M}{10^{-3}}\left(\frac{mol}{m^3}\right)$

$\pi = CRT$ (S.I. units)

$\Rightarrow \pi = \left[\frac{20}{10^{-3}M}\right] \times 8.3 \times 300$

= 800 [From (1)]

Solving we get :

$M = \frac{20 \times 8.3 \times 300}{0.8}$ g/mol

$= 62.25\ kg\ mol^{-1}$

Ans = 62

74. At 298 K, the molar conductivity of x% (w/w) MX solution (aqueous) is 123.5 S $cm^2\ mol^{-1}$. The conductance of same solution is 1.9×10^{-3} S. The value of x is _____ $\times 10^{-2}$.

(Given : cell constant = 1.3 cm^{-1}; molar mass of MX is 75 g mol^{-1}, density of aqueous solution of MX at 298 K is 1.0 g mL^{-1})

Ans. **(15)**

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

Sol. Let molarity of MX solution = y M

$$\Rightarrow \Lambda_m = c(\ell/A) \times \frac{1000}{y}$$

$$\Rightarrow 123.5 = 1.9 \times 10^{-3} \times 1.3 \times \frac{1000}{y}$$

$$\Rightarrow y = 0.02 \text{ M}$$

Let molar mass of MX = M g mol^{-1}

0.2 mol MX is dissolve in 1L solution


(0.02 × 75) gm MX is dissolved in 1000 g solution

$$\% \text{ w/w} = x\% = \frac{0.02 \times 75}{1000} \times 100$$

= 0.15%

Ans = 15

75. For a reaction A → P at T K, the half life ($t_{1/2}$) is plotted as a function of initial concentration $[A]_0$ of A as given below :



The value of x in the given figure is ____ s (Nearest integer)

Ans. (90)

Sol. From the plot, $t_{1/2} \propto [A]_0$

$$\propto [A_0]^{1-N}$$

1 – N = 1

N = 0 (order)

$$t_{1/2} = \frac{A_0}{2K}$$

$$240 = \frac{4 \times 10^{-3}}{2K} \quad \text{....(1)}$$

$$x = \frac{1.5 \times 10^{-3}}{2K} \quad \text{....(2)}$$

Divide (1) / (2)

$$\frac{240}{x} = \frac{4}{1.5}$$

x = 90

ALLEN Scholarship Admission Test (ASAT)
Win up to 90% Scholarship* On JEE, NEET & Foundation Courses
Test Dates: 5, 12, 19 & 26 April
Register for ~~500~~ ₹99/-
*Limited period offer, subject to the scholarship rules and the T&Cs.

भरोसा जताया नहीं, जीता जाता है

जब नतीजे मायने रखते हैं,
तो स्टूडेंट्स **ALLEN** KOTA को चुनते हैं

JOIN
LEADER COURSE
For Class 12th pass students

Target: JEE (Main+Adv.) 2027

Batches Starting from

15 April &
06, 27 May 2026

Click to know more



AIR 1
JEE Adv. 2025
Rajit Gupta
7 Year Classroom Course

ALLEN

ALLEN Scholarship Admission Test (ASAT)

Win up to
90% Scholarship*
On JEE, NEET &
Foundation Courses



Test Dates: 5, 12, 19 & 26 April

Register for ~~500~~ ₹99/-

*Limited period offer, subject to the scholarship rules and the T&Cs.

ALLEN ONLINE

Ace **JEE 2027** with our

Leader Online Course

- **900+** hours of LIVE classes
- **30** latest pattern tests
- **24/7** AI-powered doubt support & more



Avail up to a

90% **scholarship** with **ASAT.**

Enrol Now

ALLEN ONLINE

Prefer learning at your own pace?

Start anytime with the **Leader Self-Study PLUS Course**

- **2000+** recorded video lectures
- **32** latest-pattern tests
- Digital study material & more




Enrol Now

DON'T PREPARE HARDER
PREPARE RIGHT !

Advanced-level MCQs

Prepared By ALLEN Experts



Well Graded Exercise

Detailed Answer Key

SHOP NOW on ALLEN e-Store


ALLEN

Get The Latest
IIT-JEE Special Books
at Your Door Steps...!!

JOIN THE JOURNEY OF LEARNING

with

HANDBOOKS | ADVANCED-QB | SCORE PAPERS
PHYSICS EXPRESS | MAIN PYQ's | Adv. PYQ's



SHOP NOW

Available in HINDI & ENGLISH