

Estimation of elastic properties of KOQ-HTI sample

Author: Martin Rühlmann, Abakumov Ivan

Publication date: 24th October 2019

E-mail: abakumov_ivan@mail.ru

Introduction

Some bla bla bla here))

Add MLIB library

```
clear; close all; clc;
mlibfolder = '/home/ivan/Desktop/MLIB';
path(path, mlibfolder);
add_mlib_path;
```

1. Upload the data

```
Sample = MLD('Sample_KOQ-HTI_data.mat');

% Data = load('/remote/data/ivan/Ultrasonic_data/KOQ-HTI/Processed/KOQ-HTI_PR.txt');
% Sample.Theta = Data(:,1);      % beta [DEG] Grad
% Sample.Vp33 = Data(:,2);      % P-velocity in vertikaler (axial) "slow"
% Sample.Vp11 = Data(:,3);      % P-velocity horizontal (radial) "fast"
% Sample.Vs31 = Data(:,4);      % S-velocity vertical (axial) "slow"
% Sample.Vs21 = Data(:,5);      % S-velocity horizontal (radial) "fast"
% Sample.Vqp = Data(:,6);
% Sample.rho = 2.205;

ind1 = 1:8;          % seria of measurements
ind2 = 9:16;         % another at 90 degree
```


Plot velocities

```
figure(123)          % P velocities
fig = figure('Position', [1 1 700 550]);
plot (Sample.Theta, Sample.Vp33, 'b+', 'MarkerSize', 7, 'LineWidth', 2)
hold on
plot (Sample.Theta, Sample.Vp11, 'kx', 'MarkerSize', 7, 'LineWidth', 2)
plot (Sample.Theta(ind1)-0.5, Sample.Vqp(ind1), 'rd', 'MarkerSize', 7, 'LineWidth', 2)
err = 0.029*ones(size(Sample.Vqp(ind1)));
errorbar(Sample.Theta(ind1)-0.5, Sample.Vqp(ind1), err, 'r', 'LineStyle', 'none', 'LineWidth', 2)
plot (Sample.Theta(ind2)+0.5, Sample.Vqp(ind2), 'cd', 'MarkerSize', 7, 'LineWidth', 2)
err = 0.029*ones(size(Sample.Vqp(ind2)));
errorbar(Sample.Theta(ind2)+0.5, Sample.Vqp(ind2), err, 'c', 'LineStyle', 'none', 'LineWidth', 2)
xlabel('Theta [\circ]', 'LineWidth', 2)
```

```

ylabel('Velocity [km/s]', 'LineWidth', 2)
legend('V_{p 33}', 'V_{p 11}', 'V_{qp PR1}', 'err 2.9%', 'V_{qp PR2}', 'err 2.9%', 'Location')
title('KOQ-HTI')
xticks(-45:45:270)
xticklabels(-45:45:270)
axis([-45 245 3.375 3.675]);
grid on

```



2. Conventional analyses of Thomsen's parameters

```

C11 = Sample.rho*Sample.Vp11.^2;
C33 = Sample.rho*Sample.Vp33.^2;
C44 = Sample.rho*Sample.Vs31.^2;
C66 = Sample.rho*Sample.Vs21.^2;
Cqp = Sample.rho*Sample.Vqp.^2;
C12 = C11 - 2*C66;

ST2 = (sin(Sample.Theta/180*pi)).^2;
CT2 = (cos(Sample.Theta/180*pi)).^2;

```

```

A = (C11 + C44).*ST2 + (C33 + C44).*CT2;
B = (C11 - C44).*ST2 - (C33 - C44).*CT2;

C13 = sqrt(((2*Cqp-A).^2 - B.^2)./(4*ST2.*CT2)) - C44;

gamma = (C66 - C44)./(2*C44);
epsilon = (C11 - C33)./(2*C33);
delta = ((C13+C44).^2 - (C33-C44).^2)./(2*C33.*(C33-C44));
eta = (epsilon-delta)./(1+2*delta);

ind = ((Sample.Theta == 45) + (Sample.Theta == 135) + (Sample.Theta == 225)) == 1;
delta45.mean = mean(delta(ind));
delta45.std = std(delta(ind));

figure(21);
fig = figure('Position', [1 1 550 550]);

subplot(2,2,1)
plot(Sample.Theta,epsilon,'kd');
xlabel('Theta [\circ]','LineWidth', 2)
axis([-45 245 0 0.2]);
legend('\epsilon')
grid on
title('\epsilon')


subplot(2,2,2)
plot(Sample.Theta,gamma,'kd');
xlabel('Theta [\circ]','LineWidth', 2)
axis([-45 245 0 0.2]);
legend('\gamma')
grid on
title('\gamma')

subplot(2,2,3)
l1 = plot(Sample.Theta(ind1),delta(ind1),'rd');
hold on
l2 = plot(Sample.Theta(ind2),delta(ind2),'cd');
l1 = plot(Sample.Theta,delta45.mean*ones(size(Sample.Theta)),'r--');
legend(l1, '\delta(45\circ)')
xlabel('Theta [\circ]','LineWidth', 2)
axis([-45 245 -0.05 0.2]);
grid on
title('\delta')

subplot(2,2,4)
plot(Sample.Theta(ind1),eta(ind1),'rd');
hold on
plot(Sample.Theta(ind2),eta(ind2),'cd');
xlabel('Theta [\circ]','LineWidth', 2)
axis([-45 245 -0.2 0.2]);
grid on

```

```
title('\eta')
```



3. Least squares analysis of Thomsen's parameters


Visualize the data

```
figure(122)
fig = figure('Position', [1 1 1000 400]);
subplot(1,2,1)
plot(Sample.Theta, Sample.Vp33, '+b');
hold on
plot(Sample.Theta, Sample.Vp11, 'xk');
plot(Sample.Theta(ind1), Sample.Vqp(ind1), 'dr');
plot(Sample.Theta(ind2), Sample.Vqp(ind2), 'dc');
xlabel('Theta [\circ]')
ylabel('Velocity [km/s]')
legend('V_{p 33}', 'V_{p 11}', 'V_{qp} PR1', 'V_{qp} PR2','Location','best')
title('P-wave velocity')
```

```

subplot(1,2,2)
plot(Sample.Theta, Sample.Vs21, 'ko');
hold on
plot(Sample.Theta, Sample.Vs31, 'bo');
xlabel('Theta [\circ]')
ylabel('Velocity [km/s]')
legend('V_{s 21}', 'V_{s 31}', 'Location', 'best')
title('S-wave velocity')

```



Assign C values

```

Sample.C11 = C11;
Sample.C33 = C33;
Sample.C44 = C44;
Sample.C66 = C66;
Sample.Vqp = Sample.Vqp;

```

Find optimum delta

```

testdelta = -0.3:0.0001:0.5;
dtheta = -10:0.1:10;
nSample = Sample;
JJ = zeros(length(dtheta),length(testdelta));
for i=1:length(dtheta)
    nSample.Theta = Sample.Theta + dtheta(i);
    J = costFunction_delta(nSample,testdelta);
    JJ(i,:) = J;
end

[~, minind] = min(JJ(:));
[inda, ind] = ind2sub(size(JJ),minind);


thetapl = dtheta(ind);

```

```
delta = testdelta(indd);
```

Plot final results

```
figure(343)
fig = figure('Position', [1 1 500 400]);
plot(testdelta, JJ(indd,:), 'k-');
hold on
plot(testdelta(indd), JJ(indd,indd), 'r.','MarkerSize', 7);
xlabel('\delta')
ylabel('Cost function')
title('Cost function')
```




```
nSample.Theta = -22.5:1:225;
Vqp = get_Vqp_VTI(nSample,delta);

figure(1222)
fig = figure('Position', [1 1 700 550]);
plot (Sample.Theta, Sample.Vp33,'b+', 'MarkerSize', 7, 'LineWidth', 2)
hold on
plot (Sample.Theta, Sample.Vp11,'kx', 'MarkerSize', 7, 'LineWidth', 2)
plot (Sample.Theta(ind1)-0.5, Sample.Vqp(ind1),'rd', 'MarkerSize', 7, 'LineWidth', 2)
err = 0.029*ones(size(Sample.Vqp(ind1)));
errorbar(Sample.Theta(ind1)-0.5,Sample.Vqp(ind1),err,'r','LineStyle','none', 'LineWidth', 2)
plot (Sample.Theta(ind2)+0.5, Sample.Vqp(ind2),'cd', 'MarkerSize', 7, 'LineWidth', 2)
err = 0.029*ones(size(Sample.Vqp(ind2)));
errorbar(Sample.Theta(ind2)+0.5,Sample.Vqp(ind2),err,'c','LineStyle','none', 'LineWidth', 2)
```

```

plot(nSample.Theta-thetapl, Vqp, 'g--', 'LineWidth', 2)
xlabel('Theta [\circ]', 'LineWidth', 2)
ylabel('Velocity [km/s]', 'LineWidth', 2)
legend('V_{p 33}', 'V_{p 11}', 'V_{qp PR1}', 'err 2.9%', 'V_{qp PR2}', 'err 2.9%', 'V_{qp}^m')
title('KOQ-HTI')
xticks(-45:45:270)
xticklabels(-45:45:270)
axis([-45 245 3.375 3.675]);
grid on

```



3. Least squares analysis of Thomsen's parameters (weak anisotropy approximation)

```

Alpha = linspace(3.3,3.5,101);
Delta = linspace(-0.05,0.2,51);
Epsilon = linspace(0.02,0.12,21);
dTheta = linspace(-10,10,51);

```

```
tic
J = costFunction_delta_weak(Sample,Alpha,Delta,Epsilon,dTheta);
toc
```

Elapsed time is 6.387254 seconds.

```
[~, ind] = min(J(:));
[inda,indd,inde,indt] = ind2sub(size(J),ind);


result.alpha = Alpha(inda);
result.delta = Delta(indd);
result.epsilon = Epsilon(inde);
result.dtheta = dTheta(indt);
```

Plot the result:

```
figure(27)
fig = figure('Position', [1 1 1200 800]);

nSample.Theta = -22.5:1:225;
Vqp = get_Vqp_VTI_weak(nSample,result.alpha,result.delta,result.epsilon,result.dtheta)

figure(144)
fig = figure('Position', [1 1 700 550]);
plot (Sample.Theta, Sample.Vp33,'b+', 'MarkerSize', 7, 'LineWidth', 2)
hold on
plot (Sample.Theta, Sample.Vp11,'kx', 'MarkerSize', 7, 'LineWidth', 2)
plot (Sample.Theta(ind1)-0.5, Sample.Vqp(ind1),'rd', 'MarkerSize', 7, 'LineWidth', 2)
err = 0.029*ones(size(Sample.Vqp(ind1)));
errorbar(Sample.Theta(ind1)-0.5,Sample.Vqp(ind1),err,'r','LineStyle','none', 'LineWidth', 2)
plot (Sample.Theta(ind2)+0.5, Sample.Vqp(ind2),'cd', 'MarkerSize', 7, 'LineWidth', 2)
err = 0.029*ones(size(Sample.Vqp(ind2)));
errorbar(Sample.Theta(ind2)+0.5,Sample.Vqp(ind2),err,'c','LineStyle','none', 'LineWidth', 2)
plot(nSample.Theta+result.dtheta, Vqp, 'g--', 'LineWidth', 2)
xlabel('Theta [\circ]', 'LineWidth', 2)
ylabel('Velocity [km/s]', 'LineWidth', 2)
legend('V_{p 33}', 'V_{p 11}', 'V_{qp PR1}', 'err 2.9%', 'V_{qp PR2}', 'err 2.9%', 'V_{qp}^{\wedge}')
title('KOQ-HTI')
xticks(-45:45:270)
xticklabels(-45:45:270)
axis([-45 245 3.375 3.675]);
grid on
```

4. Find errors of estimated parameters

```
% angle error std: 3 grad ==> error in velocity 25 m/s
% error in velocity due to measurements ==> 15 m/s
% total error: 30 m
%
%                               37 mmeasurement 73
% variance for velocity error 30 m = 0.010;    0.0060
% variance for velocity error 100 m = 0.030    0.020

Theta = Sample.Theta;
iSample = Sample;

clear test;

for i=1:1:500

    iSample.Theta = Theta + 1*randn(size(Theta));
    Vqptrue = get_Vqp_VTI_weak(iSample,result.alpha,result.delta,result.epsilon,result
```

```

VqpE = Vqptrue.*(1+0.015*randn(size(Vqptrue)));
iSample.Vqp = VqpE;
iSample.Theta = Theta;

J = costFunction_delta_weak(iSample,Alpha,Delta,Epsilon,dTheta);

[~, ind] = min(J(:));
[inda,indd,inde,indt] = ind2sub(size(J),ind);

test.alpha(i) = Alpha(inda);
test.delta(i) = Delta(indd);
test.epsilon(i) = Epsilon(inde);
test.dtheta(i) = dTheta(indt);
disp(i)

```

end

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500

Plot results:

```
figure(33)
subplot(2,2,1);
histfit(test.alpha',21)
xlabel('V_{P0}, [m/s]')
dist = fitdist(test.alpha(1:100)', 'Normal');
disp(['ALPHA: Relative error: ' num2str(abs(result.alpha-dist.mu)/result.alpha*100) '%'])
```

ALPHA: Relative error: 0.12763%

```
alpha_err = 2*dist.sigma;
```

```

title(['\mu=' num2str(dist.mu,4) ' \sigma=' num2str(dist.sigma,2)])

subplot(2,2,2);
histfit(test.delta',11)
xlabel('\delta')
dist = fitdist(test.delta(1:100)', 'Normal');
disp(['DELTA: Relative error: ' num2str(abs(result.delta-dist.mu)/result.delta*100) '%

```

DELTA: Relative error: 12%

```

delta_err = 2*dist.sigma;
title(['\mu=' num2str(dist.mu,2) ' \sigma=' num2str(dist.sigma,2)])

subplot(2,2,3);
histfit(test.epsilon',11)
xlabel('\epsilon')
dist = fitdist(test.epsilon(1:100)', 'Normal');
disp(['EPSILON: Relative error: ' num2str(abs(result.epsilon-dist.mu)/result.epsilon*100) '%

```

EPSILON: Relative error: 1.8462%

```

epsilon_err = 2*dist.sigma;
title(['\mu=' num2str(dist.mu,2) ' \sigma=' num2str(dist.sigma,2)])
%
subplot(2,2,4);
histfit(test.dtheta',11)
xlabel('\Delta\theta')
dist = fitdist(test.dtheta(1:100)', 'Normal');
disp(['DTHETA: Absolute error: ' num2str(result.dtheta-dist.mu)])

```

DTHETA: Absolute error: -0.288

```

dtheta_err = 2*dist.sigma;
title(['\mu=' num2str(dist.mu,2) ' \sigma=' num2str(dist.sigma,2)])

```

