

Review of Probability Theory and Linear Algebra

Mário A. T. Figueiredo

Instituto Superior Técnico & Instituto de Telecomunicações

Lisboa, **Portugal**

LxMLS: Lisbon Machine Learning School

July 19, 2012

Outline

- Probability Theory
- Linear Algebra

What is probability?

- Classical definition: $\mathbb{P}(A) = \frac{N_A}{N}$

...with N mutually exclusive equally likely outcomes,
 N_A of which result in the occurrence of A .

Laplace, 1814

Example: $\mathbb{P}(\text{randomly drawn card is } \clubsuit) = 13/52$.

Example: $\mathbb{P}(\text{getting 1 in throw a die}) = 1/6$.

What is probability?

- Classical definition: $\mathbb{P}(A) = \frac{N_A}{N}$

...with N mutually exclusive equally likely outcomes,
 N_A of which result in the occurrence of A .

Laplace, 1814

Example: $\mathbb{P}(\text{randomly drawn card is } \clubsuit) = 13/52$.

Example: $\mathbb{P}(\text{getting 1 in throw a die}) = 1/6$.

- Frequentist definition: $\mathbb{P}(A) = \lim_{N \rightarrow \infty} \frac{N_A}{N}$

...relative frequency of occurrence of A in infinite number of trials.

What is probability?

- Classical definition: $\mathbb{P}(A) = \frac{N_A}{N}$

...with N mutually exclusive equally likely outcomes,
 N_A of which result in the occurrence of A .

Laplace, 1814

Example: $\mathbb{P}(\text{randomly drawn card is } \clubsuit) = 13/52$.

Example: $\mathbb{P}(\text{getting 1 in throw a die}) = 1/6$.

- Frequentist definition: $\mathbb{P}(A) = \lim_{N \rightarrow \infty} \frac{N_A}{N}$

...relative frequency of occurrence of A in infinite number of trials.

- Subjective probability: $\mathbb{P}(A)$ is a degree of belief.

...gives meaning to $\mathbb{P}(\text{"tomorrow will rain"})$.

Key concepts: Sample space and events

- **Sample space** \mathcal{X} = set of possible outcomes of a random experiment.

Examples:

- ▶ Tossing two coins: $\mathcal{X} = \{HH, TH, HT, TT\}$
- ▶ Roulette: $\mathcal{X} = \{1, 2, \dots, 36\}$
- ▶ Draw a card from a shuffled deck: $\mathcal{X} = \{A\clubsuit, 2\clubsuit, \dots, Q\diamondsuit, K\diamondsuit\}$.

Key concepts: Sample space and events

- **Sample space** \mathcal{X} = set of possible outcomes of a random experiment.

Examples:

- ▶ Tossing two coins: $\mathcal{X} = \{HH, TH, HT, TT\}$
- ▶ Roulette: $\mathcal{X} = \{1, 2, \dots, 36\}$
- ▶ Draw a card from a shuffled deck: $\mathcal{X} = \{A\clubsuit, 2\clubsuit, \dots, Q\spadesuit, K\spadesuit\}$.

- An **event** is a subset of \mathcal{X}

Examples:

- ▶ “exactly one H in 2-coin toss”: $A = \{TH, HT\} \subset \{HH, TH, HT, TT\}$.
- ▶ “odd number in the roulette”: $B = \{1, 3, \dots, 35\} \subset \{1, 2, \dots, 36\}$.
- ▶ “drawn a \heartsuit card”: $C = \{A\heartsuit, 2\heartsuit, \dots, K\heartsuit\} \subset \{A\clubsuit, \dots, K\spadesuit\}$

Kolmogorov's Axioms for Probability

- Probability is a function that maps events A into the interval $[0, 1]$.

Kolmogorov's axioms for probability (1933):

Kolmogorov's Axioms for Probability

- Probability is a function that maps events A into the interval $[0, 1]$.

Kolmogorov's axioms for probability (1933):

- ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A) \geq 0$

Kolmogorov's Axioms for Probability

- Probability is a function that maps events A into the interval $[0, 1]$.

Kolmogorov's axioms for probability (1933):

- ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A) \geq 0$
- ▶ $\mathbb{P}(\mathcal{X}) = 1$

Kolmogorov's Axioms for Probability

- Probability is a function that maps events A into the interval $[0, 1]$.

Kolmogorov's axioms for probability (1933):

- ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A) \geq 0$
- ▶ $\mathbb{P}(\mathcal{X}) = 1$
- ▶ If $A_1, A_2 \dots \subseteq \mathcal{X}$ are disjoint events, then $\mathbb{P}\left(\bigcup_i A_i\right) = \sum_i \mathbb{P}(A_i)$


Kolmogorov's Axioms for Probability

- Probability is a function that maps events A into the interval $[0, 1]$.

Kolmogorov's axioms for probability (1933):

- ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A) \geq 0$
 - ▶ $\mathbb{P}(\mathcal{X}) = 1$
 - ▶ If $A_1, A_2 \dots \subseteq \mathcal{X}$ are disjoint events, then $\mathbb{P}\left(\bigcup_i A_i\right) = \sum_i \mathbb{P}(A_i)$
- From these axioms, many results can be derived. **Examples:**

- ▶ $\mathbb{P}(\emptyset) = 0$
- ▶ $C \subset D \Rightarrow \mathbb{P}(C) \leq \mathbb{P}(D)$
- ▶ $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$




Conditional Probability and Independence

- If $\mathbb{P}(B) > 0$, $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$ (conditional prob. of A given B)

Conditional Probability and Independence

- If $\mathbb{P}(B) > 0$, $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$ (conditional prob. of A given B)
- ...satisfies all Kolmogorov's axioms:
 - ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A|B) \geq 0$
 - ▶ $\mathbb{P}(\mathcal{X}|B) = 1$
 - ▶ If $A_1, A_2 \dots \subseteq \mathcal{X}$ are disjoint, then
$$\mathbb{P}\left(\bigcup_i A_i \middle| B\right) = \sum_i \mathbb{P}(A_i|B)$$




Conditional Probability and Independence

- If $\mathbb{P}(B) > 0$, $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$ (conditional prob. of A given B)

- ...satisfies all Kolmogorov's axioms:

- ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A|B) \geq 0$
- ▶ $\mathbb{P}(\mathcal{X}|B) = 1$
- ▶ If $A_1, A_2 \dots \subseteq \mathcal{X}$ are disjoint, then
$$\mathbb{P}\left(\bigcup_i A_i \middle| B\right) = \sum_i \mathbb{P}(A_i|B)$$

- Events A, B are independent ($A \perp B$) $\Leftrightarrow \mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.




Conditional Probability and Independence

- If $\mathbb{P}(B) > 0$, $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$ (conditional prob. of A given B)

- ...satisfies all Kolmogorov's axioms:

- ▶ For any $A \subseteq \mathcal{X}$, $\mathbb{P}(A|B) \geq 0$
- ▶ $\mathbb{P}(\mathcal{X}|B) = 1$
- ▶ If $A_1, A_2 \dots \subseteq \mathcal{X}$ are disjoint, then
$$\mathbb{P}\left(\bigcup_i A_i \middle| B\right) = \sum_i \mathbb{P}(A_i|B)$$




- Events A, B are independent ($A \perp B$) $\Leftrightarrow \mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.
- Relationship with conditional probabilities:

$$A \perp B \Leftrightarrow \mathbb{P}(A|B) = \mathbb{P}(A)$$

Bayes Theorem

- Law of total probability: if A_1, \dots, A_n are a partition of \mathcal{X}

$$\mathbb{P}(B) = \sum_i \mathbb{P}(B|A_i)\mathbb{P}(A_i)$$




- Bayes' theorem: if A_1, \dots, A_n are a partition of \mathcal{X}

$$\mathbb{P}(A_i|B) = \frac{\mathbb{P}(B \cap A_i)}{\mathbb{P}(B)} = \frac{\mathbb{P}(B|A_i) \mathbb{P}(A_i)}{\sum_i \mathbb{P}(B|A_i)\mathbb{P}(A_i)}$$


Random Variables

- A (real) **random variable** (RV) is a function: $X : \mathcal{X} \rightarrow \mathbb{R}$



Random Variables


- A (real) **random variable** (RV) is a function: $X : \mathcal{X} \rightarrow \mathbb{R}$



- ▶ **Discrete RV**: range of X is countable (e.g., \mathbb{N} or $\{0, 1\}$)

Random Variables


- A (real) **random variable** (RV) is a function: $X : \mathcal{X} \rightarrow \mathbb{R}$



- ▶ **Discrete RV**: range of X is countable (e.g., \mathbb{N} or $\{0, 1\}$)
- ▶ **Continuous RV**: range of X is uncountable (e.g., \mathbb{R} or $[0, 1]$)

Random Variables


- A (real) **random variable** (RV) is a function: $X : \mathcal{X} \rightarrow \mathbb{R}$



- ▶ **Discrete RV**: range of X is countable (e.g., \mathbb{N} or $\{0, 1\}$)
- ▶ **Continuous RV**: range of X is uncountable (e.g., \mathbb{R} or $[0, 1]$)
- ▶ **Example**: number of head in tossing two coins,
 $\mathcal{X} = \{HH, HT, TH, TT\}$,
 $X(HH) = 2, X(HT) = X(TH) = 1, X(TT) = 0$.
Range of $X = \{0, 1, 2\}$.

Random Variables


- A (real) **random variable** (RV) is a function: $X : \mathcal{X} \rightarrow \mathbb{R}$



- ▶ **Discrete RV**: range of X is countable (e.g., \mathbb{N} or $\{0, 1\}$)
- ▶ **Continuous RV**: range of X is uncountable (e.g., \mathbb{R} or $[0, 1]$)
- ▶ **Example**: number of head in tossing two coins,
 $\mathcal{X} = \{HH, HT, TH, TT\}$,
 $X(HH) = 2, X(HT) = X(TH) = 1, X(TT) = 0$.
Range of $X = \{0, 1, 2\}$.
- ▶ **Example**: distance traveled by a tossed coin; range of $X = \mathbb{R}_+$.


Random Variables: Distribution Function

- **Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$





Random Variables: Distribution Function

- **Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$




- **Example:** number of heads in tossing 2 coins; $\text{range}(X) = \{0, 1, 2\}$.





Random Variables: Distribution Function

- Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$



- Example:** number of heads in tossing 2 coins; $\text{range}(X) = \{0, 1, 2\}$.



- Probability mass function** (discrete RV): $f_X(x) = \mathbb{P}(X = x)$,

$$F_X(x) = \sum_{x_i \leq x} f_X(x_i).$$

Important Discrete Random Variables

- **Uniform:** $X \in \{x_1, \dots, x_K\}$, pmf $f_X(x_i) = 1/K$.

Important Discrete Random Variables

- **Uniform:** $X \in \{x_1, \dots, x_K\}$, pmf $f_X(x_i) = 1/K$.
- **Bernoulli RV:** $X \in \{0, 1\}$, pmf $f_X(x) = \begin{cases} p & \Leftarrow x = 1 \\ 1 - p & \Leftarrow x = 0 \end{cases}$

Can be written compactly as $f_X(x) = p^x(1-p)^{1-x}$.

Important Discrete Random Variables

- **Uniform:** $X \in \{x_1, \dots, x_K\}$, pmf $f_X(x_i) = 1/K$.
- **Bernoulli RV:** $X \in \{0, 1\}$, pmf $f_X(x) = \begin{cases} p & \Leftarrow x = 1 \\ 1 - p & \Leftarrow x = 0 \end{cases}$

Can be written compactly as $f_X(x) = p^x(1-p)^{1-x}$.

- **Binomial RV:** $X \in \{0, 1, \dots, n\}$ (sum on n Bernoulli RVs)

$$f_X(x) = \text{Binomial}(x; n, p) = \binom{n}{x} p^x (1-p)^{(n-x)}$$

Important Discrete Random Variables

- **Uniform:** $X \in \{x_1, \dots, x_K\}$, pmf $f_X(x_i) = 1/K$.
- **Bernoulli RV:** $X \in \{0, 1\}$, pmf $f_X(x) = \begin{cases} p & \Leftarrow x = 1 \\ 1 - p & \Leftarrow x = 0 \end{cases}$


Can be written compactly as $f_X(x) = p^x(1-p)^{1-x}$.

- **Binomial RV:** $X \in \{0, 1, \dots, n\}$ (sum on n Bernoulli RVs)

$$f_X(x) = \text{Binomial}(x; n, p) = \binom{n}{x} p^x (1-p)^{(n-x)}$$


Binomial coefficients:

$$\binom{n}{x} = \frac{n!}{(n-x)! x!}$$




Random Variables: Distribution Function

- **Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$





Random Variables: Distribution Function

- **Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$




- **Example:** continuous RV with uniform distribution on $[a, b]$.





Random Variables: Distribution Function

- **Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$




- **Example:** continuous RV with uniform distribution on $[a, b]$.





- **Probability density function (pdf, continuous RV):** $f_X(x)$

Random Variables: Distribution Function

- Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$



- Example:** continuous RV with uniform distribution on $[a, b]$.




- Probability density function (pdf, continuous RV):** $f_X(x)$



$$F_X(x) = \int_{-\infty}^x f_X(u) du,$$

Random Variables: Distribution Function

- Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$



- Example:** continuous RV with uniform distribution on $[a, b]$.




- Probability density function (pdf, continuous RV):** $f_X(x)$



$$F_X(x) = \int_{-\infty}^x f_X(u) du, \quad \mathbb{P}(X \in [c, d]) = \int_c^d f_X(x) dx,$$

Random Variables: Distribution Function

- Distribution function:** $F_X(x) = \mathbb{P}(\{\omega \in \mathcal{X} : X(\omega) \leq x\})$



- Example:** continuous RV with uniform distribution on $[a, b]$.



- Probability density function (pdf, continuous RV):** $f_X(x)$

$$F_X(x) = \int_{-\infty}^x f_X(u) du, \quad \mathbb{P}(X \in [c, d]) = \int_c^d f_X(x) dx, \quad \mathbb{P}(X=x) = 0$$


Important Continuous Random Variables

- **Uniform:** $f_X(x) = \text{Uniform}(x; a, b) = \begin{cases} \frac{1}{b-a} & \Leftarrow x \in [a, b] \\ 0 & \Leftarrow x \notin [a, b] \end{cases}$
(see previous slide).

Important Continuous Random Variables

- **Uniform:** $f_X(x) = \text{Uniform}(x; a, b) = \begin{cases} \frac{1}{b-a} & \Leftarrow x \in [a, b] \\ 0 & \Leftarrow x \notin [a, b] \end{cases}$
(see previous slide).



- **Gaussian:** $f_X(x) = \mathcal{N}(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$



Important Continuous Random Variables

- **Uniform:** $f_X(x) = \text{Uniform}(x; a, b) = \begin{cases} \frac{1}{b-a} & \Leftarrow x \in [a, b] \\ 0 & \Leftarrow x \notin [a, b] \end{cases}$
(see previous slide).

- **Gaussian:** $f_X(x) = \mathcal{N}(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$



- **Exponential:** $f_X(x) = \text{Exp}(x; \lambda) = \begin{cases} \lambda e^{-\lambda x} & \Leftarrow x \geq 0 \\ 0 & \Leftarrow x < 0 \end{cases}$

Expectation of Random Variables

- **Expectation:** $\mathbb{E}(X) = \begin{cases} \sum_i x_i f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} x f_X(x) dx & X \text{ continuous} \end{cases}$

Expectation of Random Variables

- **Expectation:** $\mathbb{E}(X) = \begin{cases} \sum_i x_i f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} x f_X(x) dx & X \text{ continuous} \end{cases}$

- **Example:** Bernoulli, $f_X(x) = p^x (1 - p)^{1-x}$, for $x \in \{0, 1\}$.

$$\mathbb{E}(X) = 0(1 - p) + 1p = p.$$

Expectation of Random Variables

- **Expectation:** $\mathbb{E}(X) = \begin{cases} \sum_i x_i f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} x f_X(x) dx & X \text{ continuous} \end{cases}$

- **Example:** Bernoulli, $f_X(x) = p^x (1-p)^{1-x}$, for $x \in \{0, 1\}$.

$$\mathbb{E}(X) = 0(1-p) + 1p = p.$$

- **Example:** Binomial, $f_X(x) = \binom{n}{x} p^x (1-p)^{n-x}$, for $x \in \{0, \dots, n\}$.

$$\mathbb{E}(X) = np.$$

Expectation of Random Variables

- **Expectation:** $\mathbb{E}(X) = \begin{cases} \sum_i x_i f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} x f_X(x) dx & X \text{ continuous} \end{cases}$

- **Example:** Bernoulli, $f_X(x) = p^x (1-p)^{1-x}$, for $x \in \{0, 1\}$.

$$\mathbb{E}(X) = 0(1-p) + 1p = p.$$

- **Example:** Binomial, $f_X(x) = \binom{n}{x} p^x (1-p)^{n-x}$, for $x \in \{0, \dots, n\}$.

$$\mathbb{E}(X) = np.$$

- **Example:** Gaussian, $f_X(x) = \mathcal{N}(x; \mu, \sigma^2)$; $\mathbb{E}(X) = \mu$.

Expectation of Random Variables

- **Expectation:** $\mathbb{E}(X) = \begin{cases} \sum_i x_i f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} x f_X(x) dx & X \text{ continuous} \end{cases}$

- **Example:** Bernoulli, $f_X(x) = p^x (1-p)^{1-x}$, for $x \in \{0, 1\}$.

$$\mathbb{E}(X) = 0(1-p) + 1p = p.$$

- **Example:** Binomial, $f_X(x) = \binom{n}{x} p^x (1-p)^{n-x}$, for $x \in \{0, \dots, n\}$.

$$\mathbb{E}(X) = np.$$

- **Example:** Gaussian, $f_X(x) = \mathcal{N}(x; \mu, \sigma^2)$; $\mathbb{E}(X) = \mu$.

- **Linearity of expectation:** $\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$.

Expectation of Functions of Random Variables

- $\mathbb{E}(g(X)) = \begin{cases} \sum_i g(x_i) f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx & X \text{ continuous} \end{cases}$

Expectation of Functions of Random Variables

- $\mathbb{E}(g(X)) = \begin{cases} \sum_i g(x_i) f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx & X \text{ continuous} \end{cases}$
- **Example:** variance, $\text{var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right)$

Expectation of Functions of Random Variables

- $\mathbb{E}(g(X)) = \begin{cases} \sum_i g(x_i) f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx & X \text{ continuous} \end{cases}$
- **Example:** variance, $\text{var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right) = \mathbb{E}(X^2) - \mathbb{E}(X)^2$
- **Example:** Bernoulli variance, $\mathbb{E}(X^2) = \mathbb{E}(X) = p$

Expectation of Functions of Random Variables

- $\mathbb{E}(g(X)) = \begin{cases} \sum_i g(x_i) f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx & X \text{ continuous} \end{cases}$
- **Example:** variance, $\text{var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right) = \mathbb{E}(X^2) - \mathbb{E}(X)^2$
- **Example:** Bernoulli variance, $\mathbb{E}(X^2) = \mathbb{E}(X) = p$, thus $\text{var}(X) = p(1 - p)$.

Expectation of Functions of Random Variables

- $\mathbb{E}(g(X)) = \begin{cases} \sum_i g(x_i) f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx & X \text{ continuous} \end{cases}$
- **Example:** variance, $\text{var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right) = \mathbb{E}(X^2) - \mathbb{E}(X)^2$
- **Example:** Bernoulli variance, $\mathbb{E}(X^2) = \mathbb{E}(X) = p$, thus $\text{var}(X) = p(1 - p)$.
- **Example:** Gaussian variance, $\mathbb{E}((X - \mu)^2) = \sigma^2$.

Expectation of Functions of Random Variables

- $\mathbb{E}(g(X)) = \begin{cases} \sum_i g(x_i) f_X(x_i) & X \text{ discrete on } \{x_1, \dots, x_K\} \\ \int_{-\infty}^{\infty} g(x) f_X(x) dx & X \text{ continuous} \end{cases}$

- **Example:** variance, $\text{var}(X) = \mathbb{E}\left((X - \mathbb{E}(X))^2\right) = \mathbb{E}(X^2) - \mathbb{E}(X)^2$

- **Example:** Bernoulli variance, $\mathbb{E}(X^2) = \mathbb{E}(X) = p$, thus $\text{var}(X) = p(1 - p)$.

- **Example:** Gaussian variance, $\mathbb{E}((X - \mu)^2) = \sigma^2$.

- Probability as expectation of indicator, $\mathbf{1}_A(x) = \begin{cases} 1 & \Leftarrow x \in A \\ 0 & \Leftarrow x \notin A \end{cases}$

$$\mathbb{P}(X \in A) = \int_A f_X(x) dx = \int \mathbf{1}_A(x) f_X(x) dx = \mathbb{E}(\mathbf{1}_A(X))$$

Two (or More) Random Variables

- **Joint pmf** of two discrete RVs: $f_{X,Y}(x,y) = \mathbb{P}(X = x \wedge Y = y)$.

Extends trivially to more than two RVs.

Two (or More) Random Variables

- **Joint pmf** of two discrete RVs: $f_{X,Y}(x,y) = \mathbb{P}(X = x \wedge Y = y)$.

Extends trivially to more than two RVs.

- **Joint pdf** of two continuous RVs: $f_{X,Y}(x,y)$, such that

$$\mathbb{P}(X \in A) = \int \int_A f_{X,Y}(x,y) dx dy, \quad A \subset \mathbb{R}^2$$

Extends trivially to more than two RVs.

Two (or More) Random Variables

- **Joint pmf** of two discrete RVs: $f_{X,Y}(x,y) = \mathbb{P}(X = x \wedge Y = y)$.

Extends trivially to more than two RVs.

- **Joint pdf** of two continuous RVs: $f_{X,Y}(x,y)$, such that

$$\mathbb{P}(X \in A) = \int \int_A f_{X,Y}(x,y) dx dy, \quad A \subset \mathbb{R}^2$$

Extends trivially to more than two RVs.

- **Marginalization**: $f_Y(y) = \begin{cases} \sum_x f_{X,Y}(x,y), & \text{if } X \text{ is discrete} \\ \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx, & \text{if } X \text{ continuous} \end{cases}$

Two (or More) Random Variables

- **Joint pmf** of two discrete RVs: $f_{X,Y}(x,y) = \mathbb{P}(X = x \wedge Y = y)$.

Extends trivially to more than two RVs.

- **Joint pdf** of two continuous RVs: $f_{X,Y}(x,y)$, such that

$$\mathbb{P}(X \in A) = \int \int_A f_{X,Y}(x,y) dx dy, \quad A \subset \mathbb{R}^2$$

Extends trivially to more than two RVs.

- **Marginalization**: $f_Y(y) = \begin{cases} \sum_x f_{X,Y}(x,y), & \text{if } X \text{ is discrete} \\ \int_{-\infty}^{\infty} f_{X,Y}(x,y) dx, & \text{if } X \text{ continuous} \end{cases}$

- **Independence**: $X \perp\!\!\!\perp Y \Leftrightarrow f_{X,Y}(x,y) = f_X(x) f_Y(y)$.

Conditionals and Bayes' Theorem

- **Conditional pmf** (discrete RVs):

$$f_{X|Y}(x|y) = \mathbb{P}(X = x | Y = y) = \frac{\mathbb{P}(X = x \wedge Y = y)}{\mathbb{P}(Y = y)} = \frac{f_{X,Y}(x, y)}{f_Y(y)}.$$

Conditionals and Bayes' Theorem

- **Conditional pmf** (discrete RVs):

$$f_{X|Y}(x|y) = \mathbb{P}(X = x | Y = y) = \frac{\mathbb{P}(X = x \wedge Y = y)}{\mathbb{P}(Y = y)} = \frac{f_{X,Y}(x, y)}{f_Y(y)}.$$

- **Conditional pdf** (continuous RVs): $f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}$

...the meaning is technically delicate.

Conditionals and Bayes' Theorem

- **Conditional pmf** (discrete RVs):

$$f_{X|Y}(x|y) = \mathbb{P}(X = x | Y = y) = \frac{\mathbb{P}(X = x \wedge Y = y)}{\mathbb{P}(Y = y)} = \frac{f_{X,Y}(x, y)}{f_Y(y)}.$$

- **Conditional pdf** (continuous RVs): $f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}$

...the meaning is technically delicate.

- **Bayes' theorem**: $f_{X|Y}(x|y) = \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)}$ (pdf or pmf).

Conditionals and Bayes' Theorem

- **Conditional pmf** (discrete RVs):

$$f_{X|Y}(x|y) = \mathbb{P}(X = x | Y = y) = \frac{\mathbb{P}(X = x \wedge Y = y)}{\mathbb{P}(Y = y)} = \frac{f_{X,Y}(x, y)}{f_Y(y)}.$$

- **Conditional pdf** (continuous RVs): $f_{X|Y}(x|y) = \frac{f_{X,Y}(x, y)}{f_Y(y)}$

...the meaning is technically delicate.

- **Bayes' theorem**: $f_{X|Y}(x|y) = \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)}$ (pdf or pmf).

- Also valid in the mixed case (e.g., X continuous, Y discrete).

Joint, Marginal, and Conditional Probabilities: An Example

- A pair of binary variables $X, Y \in \{0, 1\}$, with **joint** pmf:

$f_{X,Y}(x, y)$	$Y = 0$	$Y = 1$
$X = 0$	1/5	2/5
$X = 1$	1/10	3/10

Joint, Marginal, and Conditional Probabilities: An Example

- A pair of binary variables $X, Y \in \{0, 1\}$, with **joint** pmf:

$f_{X,Y}(x, y)$	$Y = 0$	$Y = 1$
$X = 0$	1/5	2/5
$X = 1$	1/10	3/10

- Marginals:** $f_X(0) = \frac{1}{5} + \frac{2}{5} = \frac{3}{5}$, $f_X(1) = \frac{1}{10} + \frac{3}{10} = \frac{4}{10}$,
 $f_Y(0) = \frac{1}{5} + \frac{1}{10} = \frac{3}{10}$, $f_Y(1) = \frac{2}{5} + \frac{3}{10} = \frac{7}{10}$.

Joint, Marginal, and Conditional Probabilities: An Example

- A pair of binary variables $X, Y \in \{0, 1\}$, with **joint** pmf:

$f_{X,Y}(x, y)$	$Y = 0$	$Y = 1$
$X = 0$	1/5	2/5
$X = 1$	1/10	3/10

- Marginals:** $f_X(0) = \frac{1}{5} + \frac{2}{5} = \frac{3}{5}$, $f_X(1) = \frac{1}{10} + \frac{3}{10} = \frac{4}{10}$,
 $f_Y(0) = \frac{1}{5} + \frac{1}{10} = \frac{3}{10}$, $f_Y(1) = \frac{2}{5} + \frac{3}{10} = \frac{7}{10}$.

- Conditional** probabilities:

$f_{X Y}(x y)$	$Y = 0$	$Y = 1$
$X = 0$	2/3	4/7
$X = 1$	1/3	3/7

$f_{Y X}(y x)$	$Y = 0$	$Y = 1$
$X = 0$	1/3	2/3
$X = 1$	1/4	3/4

An Important Multivariate RV: Multinomial

- **Multinomial:** $X = (X_1, \dots, X_K)$, $X_i \in \{0, \dots, n\}$, such that $\sum_i X_i = n$,

$$f_X(x_1, \dots, x_K) = \begin{cases} \binom{n}{x_1 \ x_2 \ \dots \ x_K} p_1^{x_1} p_2^{x_2} \cdots p_K^{x_K} & \Leftrightarrow \sum_i x_i = n \\ 0 & \Leftrightarrow \sum_i x_i \neq n \end{cases}$$

$$\binom{n}{x_1 \ x_2 \ \dots \ x_K} = \frac{n!}{x_1! x_2! \cdots x_K!}$$

Parameters: $p_1, \dots, p_K \geq 0$, such that $\sum_i p_i = 1$.

An Important Multivariate RV: Multinomial

- **Multinomial:** $X = (X_1, \dots, X_K)$, $X_i \in \{0, \dots, n\}$, such that $\sum_i X_i = n$,

$$f_X(x_1, \dots, x_K) = \begin{cases} \binom{n}{x_1 \ x_2 \ \dots \ x_K} p_1^{x_1} p_2^{x_2} \cdots p_K^{x_K} & \Leftrightarrow \sum_i x_i = n \\ 0 & \Leftrightarrow \sum_i x_i \neq n \end{cases}$$

$$\binom{n}{x_1 \ x_2 \ \dots \ x_K} = \frac{n!}{x_1! x_2! \cdots x_K!}$$

Parameters: $p_1, \dots, p_K \geq 0$, such that $\sum_i p_i = 1$.

- Generalizes the binomial from binary to K -classes.

An Important Multivariate RV: Multinomial

- **Multinomial:** $X = (X_1, \dots, X_K)$, $X_i \in \{0, \dots, n\}$, such that $\sum_i X_i = n$,

$$f_X(x_1, \dots, x_K) = \begin{cases} \binom{n}{x_1 \ x_2 \ \dots \ x_K} p_1^{x_1} p_2^{x_2} \cdots p_K^{x_K} & \Leftarrow \sum_i x_i = n \\ 0 & \Leftarrow \sum_i x_i \neq n \end{cases}$$

$$\binom{n}{x_1 \ x_2 \ \dots \ x_K} = \frac{n!}{x_1! x_2! \cdots x_K!}$$

Parameters: $p_1, \dots, p_K \geq 0$, such that $\sum_i p_i = 1$.

- Generalizes the binomial from binary to K -classes.
- **Example:** tossing n independent fair dice, $p_1 = \dots = p_6 = 1/6$.
 x_i = number of outcomes with i dots. Of course, $\sum_i x_i = n$.

An Important Multivariate RV: Gaussian

- **Multivariate Gaussian:** $X \in \mathbb{R}^n$,

$$f_X(x) = \mathcal{N}(x; \mu, C) = \frac{1}{\sqrt{\det(2\pi C)}} \exp\left(-\frac{1}{2}(x - \mu)^T C^{-1}(x - \mu)\right)$$

An Important Multivariate RV: Gaussian

- **Multivariate Gaussian:** $X \in \mathbb{R}^n$,

$$f_X(x) = \mathcal{N}(x; \mu, C) = \frac{1}{\sqrt{\det(2\pi C)}} \exp\left(-\frac{1}{2}(x - \mu)^T C^{-1}(x - \mu)\right)$$


- Parameters: mean $\mu \in \mathbb{R}^n$ and covariance matrix $C \in \mathbb{R}^{n \times n}$.

An Important Multivariate RV: Gaussian

- **Multivariate Gaussian:** $X \in \mathbb{R}^n$,

$$f_X(x) = \mathcal{N}(x; \mu, C) = \frac{1}{\sqrt{\det(2\pi C)}} \exp\left(-\frac{1}{2}(x - \mu)^T C^{-1}(x - \mu)\right)$$

- Parameters: mean $\mu \in \mathbb{R}^n$ and covariance matrix $C \in \mathbb{R}^{n \times n}$.



Statistical Inference 101 (A Lot More Tomorrow)

- Two RVs, with joint pdf or pmf $f_{X,Y}(x,y) = f_{Y|X}(y|x) f_X(x)$.
 $f_{Y|X}(y|x)$ is often called **likelihood function**; $f_X(x)$ is called **prior**.

Statistical Inference 101 (A Lot More Tomorrow)

- Two RVs, with joint pdf or pmf $f_{X,Y}(x,y) = f_{Y|X}(y|x) f_X(x)$.
 $f_{Y|X}(y|x)$ is often called **likelihood function**; $f_X(x)$ is called **prior**.
- One RV is observed, say $Y = y$; goal is to infer the other RV X .

Statistical Inference 101 (A Lot More Tomorrow)

- Two RVs, with joint pdf or pmf $f_{X,Y}(x,y) = f_{Y|X}(y|x) f_X(x)$.
 $f_{Y|X}(y|x)$ is often called **likelihood function**; $f_X(x)$ is called **prior**.
- One RV is observed, say $Y = y$; goal is to infer the other RV X .
- **Maximum likelihood** (ML) criterion: $\hat{x}_{\text{ML}} = \arg \max_x f_{Y|X}(y|x)$

Statistical Inference 101 (A Lot More Tomorrow)

- Two RVs, with joint pdf or pmf $f_{X,Y}(x,y) = f_{Y|X}(y|x) f_X(x)$.
 $f_{Y|X}(y|x)$ is often called **likelihood function**; $f_X(x)$ is called **prior**.
- One RV is observed, say $Y = y$; goal is to infer the other RV X .
- **Maximum likelihood** (ML) criterion: $\hat{x}_{\text{ML}} = \arg \max_x f_{Y|X}(y|x)$
- **Maximum *a posteriori*** (MAP) criterion:

$$\hat{x}_{\text{MAP}} = \arg \max_x f_{X|Y}(x|y) = \arg \max_x \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)} = \arg \max_x f_{Y|X}(y|x) f_X(x)$$

Statistical Inference 101 (A Lot More Tomorrow)

- Two RVs, with joint pdf or pmf $f_{X,Y}(x,y) = f_{Y|X}(y|x) f_X(x)$.
 $f_{Y|X}(y|x)$ is often called **likelihood function**; $f_X(x)$ is called **prior**.
- One RV is observed, say $Y = y$; goal is to infer the other RV X .
- **Maximum likelihood** (ML) criterion: $\hat{x}_{\text{ML}} = \arg \max_x f_{Y|X}(y|x)$
- **Maximum a posteriori** (MAP) criterion:

$$\hat{x}_{\text{MAP}} = \arg \max_x f_{X|Y}(x|y) = \arg \max_x \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)} = \arg \max_x f_{Y|X}(y|x) f_X(x)$$

- **Posterior mean** (PM) criterion (for continuous RVs):

$$\hat{x}_{\text{PM}} = \mathbb{E}(X|Y = y) = \int x f_{X|Y}(x|y) dx$$

Statistical Inference: Example

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs:
 $Y = (Y_1, \dots, Y_n)$, with $Y_i \in \{0, 1\}$.
Common pmf $f_{Y_i|X}(y|x) = x^y(1-x)^{1-y}$, where $x \in [0, 1]$.

Statistical Inference: Example

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs: $Y = (Y_1, \dots, Y_n)$, with $Y_i \in \{0, 1\}$.
Common pmf $f_{Y_i|X}(y|x) = x^y(1-x)^{1-y}$, where $x \in [0, 1]$.

- Likelihood function: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i}(1-x)^{1-y_i}$

Log-likelihood function:

$$\log f_{Y|X}((y_1, \dots, y_n)|x) = n \log(1-x) + \log \frac{x}{1-x} \sum_{i=1}^n y_i$$

Statistical Inference: Example

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs: $Y = (Y_1, \dots, Y_n)$, with $Y_i \in \{0, 1\}$.
Common pmf $f_{Y_i|X}(y|x) = x^y(1-x)^{1-y}$, where $x \in [0, 1]$.

- Likelihood function: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i}(1-x)^{1-y_i}$

Log-likelihood function:

$$\log f_{Y|X}((y_1, \dots, y_n)|x) = n \log(1-x) + \log \frac{x}{1-x} \sum_{i=1}^n y_i$$

- Maximum likelihood: $\hat{x}_{\text{ML}} = \arg \max_x f_{Y|X}(y|x) = \frac{1}{n} \sum_{i=1}^n y_i$

Statistical Inference: Example

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs: $Y = (Y_1, \dots, Y_n)$, with $Y_i \in \{0, 1\}$.
Common pmf $f_{Y_i|X}(y|x) = x^y(1-x)^{1-y}$, where $x \in [0, 1]$.

- Likelihood function: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i}(1-x)^{1-y_i}$

Log-likelihood function:

$$\log f_{Y|X}((y_1, \dots, y_n)|x) = n \log(1-x) + \log \frac{x}{1-x} \sum_{i=1}^n y_i$$

- Maximum likelihood: $\hat{x}_{\text{ML}} = \arg \max_x f_{Y|X}(y|x) = \frac{1}{n} \sum_{i=1}^n y_i$
- Example: $n = 10$, observed $y = (1, 1, 1, 0, 1, 0, 0, 1, 1, 1)$, $\hat{x}_{\text{ML}} = 7/10$.

Statistical Inference: Example (Continuation)

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs.


Statistical Inference: Example (Continuation)

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs.

- Likelihood: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i} (1-x)^{1-y_i} = x^{\sum_i y_i} (1-x)^{n-\sum_i y_i}$

Statistical Inference: Example (Continuation)

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs.
- Likelihood: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i} (1-x)^{1-y_i} = x^{\sum_i y_i} (1-x)^{n-\sum_i y_i}$
- Beta (conjugate) prior: $f_X(x) \propto x^{\alpha-1} (1-x)^{\beta-1}$, $\alpha, \beta > 0$




Statistical Inference: Example (Continuation)

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs.

- Likelihood: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i} (1-x)^{1-y_i} = x^{\sum_i y_i} (1-x)^{n-\sum_i y_i}$

- Beta (conjugate) prior: $f_X(x) \propto x^{\alpha-1} (1-x)^{\beta-1}$, $\alpha, \beta > 0$

► MAP estimate: $\hat{x}_{\text{MAP}} = \frac{\alpha + \sum_i y_i - 1}{\alpha + \beta + n - 2}$



Statistical Inference: Example (Continuation)

- Observed n i.i.d. (independent identically distributed) Bernoulli RVs.


- Likelihood: $f_{Y|X}((y_1, \dots, y_n)|x) = \prod_{i=1}^n x^{y_i} (1-x)^{1-y_i} = x^{\sum_i y_i} (1-x)^{n-\sum_i y_i}$

- Beta (conjugate) prior: $f_X(x) \propto x^{\alpha-1} (1-x)^{\beta-1}$, $\alpha, \beta > 0$

► MAP estimate: $\hat{x}_{\text{MAP}} = \frac{\alpha + \sum_i y_i - 1}{\alpha + \beta + n - 2}$

► Example: $\alpha = 2$, $\beta = 5$, $n = 10$,
 $y = (1, 1, 1, 0, 1, 0, 0, 1, 1, 1)$,

$$\hat{x}_{\text{MAP}} = \frac{8}{15} \quad (\text{recall } \hat{x}_{\text{ML}} = 7/10)$$



Agenda

- ~~Probability Theory~~ ✓
- Linear Algebra

Linear Algebra

- Linear algebra provides (among many other things) a compact way of representing, studying, and solving linear systems of equations

Linear Algebra

- Linear algebra provides (among many other things) a compact way of representing, studying, and solving linear systems of equations
- **Example:** the system

$$4x_1 - 5x_2 = -13$$

$$-2x_1 + 3x_2 = 9$$

can be written compactly as $Ax = b$, where

$$A = \begin{bmatrix} 4 & -5 \\ -2 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} -13 \\ 9 \end{bmatrix}$$

Linear Algebra

- Linear algebra provides (among many other things) a compact way of representing, studying, and solving linear systems of equations
- **Example:** the system

$$\begin{aligned}4x_1 - 5x_2 &= -13 \\ -2x_1 + 3x_2 &= 9\end{aligned}$$

can be written compactly as $Ax = b$, where

$$A = \begin{bmatrix} 4 & -5 \\ -2 & 3 \end{bmatrix}, \quad b = \begin{bmatrix} -13 \\ 9 \end{bmatrix}$$

- It can be solved as $x = A^{-1}b$ (if A^{-1} exists).

Notation: Matrices and Vectors

- $A \in \mathbb{R}^{m \times n}$ is a **matrix** with m rows and n columns.

$$A = \begin{bmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & \ddots & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{bmatrix}.$$

Notation: Matrices and Vectors

- $A \in \mathbb{R}^{m \times n}$ is a **matrix** with m rows and n columns.

$$A = \begin{bmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & \ddots & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{bmatrix}.$$

- $x \in \mathbb{R}^n$ is a **vector** with n components,

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

Notation: Matrices and Vectors

- $A \in \mathbb{R}^{m \times n}$ is a **matrix** with m rows and n columns.

$$A = \begin{bmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & \ddots & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{bmatrix}.$$

- $x \in \mathbb{R}^n$ is a **vector** with n components,

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

- A **(column) vector** is a matrix with n rows and 1 column.

Notation: Matrices and Vectors

- $A \in \mathbb{R}^{m \times n}$ is a **matrix** with m rows and n columns.

$$A = \begin{bmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & \ddots & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{bmatrix}.$$

- $x \in \mathbb{R}^n$ is a **vector** with n components,

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

- A **(column) vector** is a matrix with n rows and 1 column.
- A matrix with 1 row and n columns is called a **row vector**.

Matrix Transpose and Products

- Given matrix $A \in \mathbb{R}^{m \times n}$, its **transpose** A^T is such that $(A^T)_{ij} = A_{j,i}$.

Matrix Transpose and Products

- Given matrix $A \in \mathbb{R}^{m \times n}$, its **transpose** A^T is such that $(A^T)_{i,j} = A_{j,i}$.
- A matrix A is **symmetric** if $A^T = A$.

Matrix Transpose and Products

- Given matrix $A \in \mathbb{R}^{m \times n}$, its **transpose** A^T is such that $(A^T)_{i,j} = A_{j,i}$.
- A matrix A is **symmetric** if $A^T = A$.
- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

Matrix Transpose and Products

- Given matrix $A \in \mathbb{R}^{m \times n}$, its **transpose** A^T is such that $(A^T)_{i,j} = A_{j,i}$.
- A matrix A is **symmetric** if $A^T = A$.
- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

- **Inner product** between vectors $x, y \in \mathbb{R}^n$:

$$\langle x, y \rangle = x^T y = y^T x = \sum_{i=1}^n x_i y_i \in \mathbb{R}.$$

Matrix Transpose and Products

- Given matrix $A \in \mathbb{R}^{m \times n}$, its **transpose** A^T is such that $(A^T)_{i,j} = A_{j,i}$.
- A matrix A is **symmetric** if $A^T = A$.
- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

- Inner product** between vectors $x, y \in \mathbb{R}^n$:

$$\langle x, y \rangle = x^T y = y^T x = \sum_{i=1}^n x_i y_i \in \mathbb{R}.$$

- Outer product** between vectors $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$: $xy^T \in \mathbb{R}^{n \times m}$, where $(xy^T)_{i,j} = x_i y_j$.

Properties of Matrix Products and Transposes

- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

Properties of Matrix Products and Transposes

- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

- Matrix product is **associative**: $(AB)C = A(BC)$.

Properties of Matrix Products and Transposes

- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

- Matrix product is **associative**: $(AB)C = A(BC)$.
- In general, matrix product is not **commutative**: $AB \neq BA$.

Properties of Matrix Products and Transposes

- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

- Matrix product is **associative**: $(AB)C = A(BC)$.
- In general, matrix product is not **commutative**: $AB \neq BA$.
- Transpose of product: $(AB)^T = B^T A^T$.

Properties of Matrix Products and Transposes

- Given matrices $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, their **product** is

$$C = AB \in \mathbb{R}^{m \times p} \quad \text{where} \quad C_{i,j} = \sum_{k=1}^n A_{i,k} B_{k,j}$$

- Matrix product is **associative**: $(AB)C = A(BC)$.
- In general, matrix product is not **commutative**: $AB \neq BA$.
- Transpose of product: $(AB)^T = B^T A^T$.
- Transpose of sum: $(A + B)^T = A^T + B^T$.

Norms

- The **norm** of a vector is (informally) its “length”. Euclidean norm:

$$\|x\|_2 = \sqrt{\langle x, x \rangle} = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}.$$

Norms

- The **norm** of a vector is (informally) its “length”. Euclidean norm:

$$\|x\|_2 = \sqrt{\langle x, x \rangle} = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}.$$

- More generally, the ℓ_p norm of a vector $x \in \mathbb{R}^n$, where $p \geq 1$,

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}.$$

Norms

- The **norm** of a vector is (informally) its “length”. Euclidean norm:

$$\|x\|_2 = \sqrt{\langle x, x \rangle} = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}.$$

- More generally, the ℓ_p norm of a vector $x \in \mathbb{R}^n$, where $p \geq 1$,

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}.$$

- Notable case: the ℓ_1 norm, $\|x\|_1 = \sum_i |x_i|$.

Norms

- The **norm** of a vector is (informally) its “length”. Euclidean norm:

$$\|x\|_2 = \sqrt{\langle x, x \rangle} = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}.$$

- More generally, the ℓ_p norm of a vector $x \in \mathbb{R}^n$, where $p \geq 1$,

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}.$$

- Notable case: the ℓ_1 norm, $\|x\|_1 = \sum_i |x_i|$.
- Notable case: the ℓ_∞ norm, $\|x\|_\infty = \max\{|x_1|, \dots, |x_n|\}$.

Norms

- The **norm** of a vector is (informally) its “length”. Euclidean norm:

$$\|x\|_2 = \sqrt{\langle x, x \rangle} = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}.$$

- More generally, the ℓ_p norm of a vector $x \in \mathbb{R}^n$, where $p \geq 1$,

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}.$$

- Notable case: the ℓ_1 norm, $\|x\|_1 = \sum_i |x_i|$.
- Notable case: the ℓ_∞ norm, $\|x\|_\infty = \max\{|x_1|, \dots, |x_n|\}$.
- Notable case: the ℓ_0 “norm” (not): $\|x\|_0 = |\{i : x_i \neq 0\}|$.

Special Matrices

- The **identity matrix** $I \in \mathbb{R}^{n \times n}$ is a square matrix such that

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \quad I = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Special Matrices

- The **identity matrix** $I \in \mathbb{R}^{n \times n}$ is a square matrix such that

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \quad I = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

- Neutral element of matrix product: $AI = IA = A$.

Special Matrices

- The **identity matrix** $I \in \mathbb{R}^{n \times n}$ is a square matrix such that

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \quad I = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

- Neutral element of matrix product: $AI = IA = A$.
- Diagonal matrix: $A \in \mathbb{R}^{n \times n}$ is diagonal if $(i \neq j) \Rightarrow A_{ij} = 0$.

Special Matrices

- The **identity matrix** $I \in \mathbb{R}^{n \times n}$ is a square matrix such that

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \quad I = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

- Neutral element of matrix product: $AI = IA = A$.
- Diagonal matrix: $A \in \mathbb{R}^{n \times n}$ is diagonal if $(i \neq j) \Rightarrow A_{i,j} = 0$.
- Upper triangular matrix: $(j < i) \Rightarrow A_{i,j} = 0$.

Special Matrices

- The **identity matrix** $I \in \mathbb{R}^{n \times n}$ is a square matrix such that

$$I_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases} \quad I = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

- Neutral element of matrix product: $AI = IA = A$.
- Diagonal matrix: $A \in \mathbb{R}^{n \times n}$ is diagonal if $(i \neq j) \Rightarrow A_{i,j} = 0$.
- Upper triangular matrix: $(j < i) \Rightarrow A_{i,j} = 0$.
- Lower triangular matrix: $(j > i) \Rightarrow A_{i,j} = 0$.

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

- Eigenvalues of diagonal matrix are the elements in the diagonal.

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

- Eigenvalues of diagonal matrix are the elements in the diagonal.
- Matrix **trace**:

$$\text{trace}(A) = \sum_i A_{i,i} = \sum_i \lambda_i$$

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

- Eigenvalues of diagonal matrix are the elements in the diagonal.
- Matrix **trace**:

$$\text{trace}(A) = \sum_i A_{i,i} = \sum_i \lambda_i$$

- Matrix **determinant**:

$$|A| = \det(A) = \prod_i \lambda_i$$

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

- Eigenvalues of diagonal matrix are the elements in the diagonal.
- Matrix **trace**:

$$\text{trace}(A) = \sum_i A_{i,i} = \sum_i \lambda_i$$

- Matrix **determinant**:

$$|A| = \det(A) = \prod_i \lambda_i$$

- Properties: $|AB| = |A||B|$,

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

- Eigenvalues of diagonal matrix are the elements in the diagonal.
- Matrix **trace**:

$$\text{trace}(A) = \sum_i A_{i,i} = \sum_i \lambda_i$$

- Matrix **determinant**:

$$|A| = \det(A) = \prod_i \lambda_i$$

- Properties: $|AB| = |A||B|$, $|A^T| = |A|$,

Eigenvalues, eigenvectors, determinant, trace

- A vector $x \in \mathbb{R}^n$ is an **eigenvector** of matrix $A \in \mathbb{R}^{n \times n}$ if

$$Ax = \lambda x,$$

where $\lambda \in \mathbb{R}$ is the corresponding **eigenvalue**.

- Eigenvalues of diagonal matrix are the elements in the diagonal.
- Matrix **trace**:

$$\text{trace}(A) = \sum_i A_{i,i} = \sum_i \lambda_i$$

- Matrix **determinant**:

$$|A| = \det(A) = \prod_i \lambda_i$$

- Properties: $|AB| = |A||B|$, $|A^T| = |A|$, $|\alpha A| = \alpha^n |A|$

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.
- Determinant of inverse: $\det(A^{-1}) = \frac{1}{\det(A)}$.

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.
- Determinant of inverse: $\det(A^{-1}) = \frac{1}{\det(A)}$.
- Solving system $Ax = b$, if A is invertible: $x = A^{-1}b$.

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.
- Determinant of inverse: $\det(A^{-1}) = \frac{1}{\det(A)}$.
- Solving system $Ax = b$, if A is invertible: $x = A^{-1}b$.
- Properties: $(A^{-1})^{-1} = A$,

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.
- Determinant of inverse: $\det(A^{-1}) = \frac{1}{\det(A)}$.
- Solving system $Ax = b$, if A is invertible: $x = A^{-1}b$.
- Properties: $(A^{-1})^{-1} = A$, $(A^{-1})^T = (A^T)^{-1}$,

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.
- Determinant of inverse: $\det(A^{-1}) = \frac{1}{\det(A)}$.
- Solving system $Ax = b$, if A is invertible: $x = A^{-1}b$.
- Properties: $(A^{-1})^{-1} = A$, $(A^{-1})^T = (A^T)^{-1}$, $(AB)^{-1} = B^{-1}A^{-1}$

Matrix Inverse

- Matrix $A \in \mathbb{R}^{n \times n}$ is **invertible** if there is $B \in \mathbb{R}^{n \times n}$ s.t. $AB = BA = I$.
- ...matrix B such that $AB = BA = I$ denoted $B = A^{-1}$ (**inverse** of A).
- Matrix $A \in \mathbb{R}^{n \times n}$ is invertible $\Leftrightarrow \det(A) \neq 0$.
- Determinant of inverse: $\det(A^{-1}) = \frac{1}{\det(A)}$.
- Solving system $Ax = b$, if A is invertible: $x = A^{-1}b$.
- Properties: $(A^{-1})^{-1} = A$, $(A^{-1})^T = (A^T)^{-1}$, $(AB)^{-1} = B^{-1}A^{-1}$
- There are many algorithms to compute A^{-1} ; general case, computational cost $O(n^3)$.

Quadratic Forms and Positive (Semi-)Definite Matrices

- Given matrix $A \in \mathbb{R}^{n \times n}$ and vector $x \in \mathbb{R}^n$,

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{i,j} x_i x_j \in \mathbb{R}$$

is called a **quadratic form**.

Quadratic Forms and Positive (Semi-)Definite Matrices

- Given matrix $A \in \mathbb{R}^{n \times n}$ and vector $x \in \mathbb{R}^n$,

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{i,j} x_i x_j \in \mathbb{R}$$

is called a **quadratic form**.

- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive semi-definite** (PSD) if, for any $x \in \mathbb{R}^n$, $x^T A x \geq 0$.

Quadratic Forms and Positive (Semi-)Definite Matrices

- Given matrix $A \in \mathbb{R}^{n \times n}$ and vector $x \in \mathbb{R}^n$,

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{i,j} x_i x_j \in \mathbb{R}$$

is called a **quadratic form**.

- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive semi-definite** (PSD) if, for any $x \in \mathbb{R}^n$, $x^T A x \geq 0$.
- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive definite** (PD) if, for any $x \in \mathbb{R}^n$, $(x \neq 0) \Rightarrow x^T A x > 0$.

Quadratic Forms and Positive (Semi-)Definite Matrices

- Given matrix $A \in \mathbb{R}^{n \times n}$ and vector $x \in \mathbb{R}^n$,

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{i,j} x_i x_j \in \mathbb{R}$$

is called a **quadratic form**.

- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive semi-definite** (PSD) if, for any $x \in \mathbb{R}^n$, $x^T A x \geq 0$.
- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive definite** (PD) if, for any $x \in \mathbb{R}^n$, $(x \neq 0) \Rightarrow x^T A x > 0$.
- Matrix $A \in \mathbb{R}^{n \times n}$ is PSD \Leftrightarrow all $\lambda_i(A) \geq 0$.

Quadratic Forms and Positive (Semi-)Definite Matrices

- Given matrix $A \in \mathbb{R}^{n \times n}$ and vector $x \in \mathbb{R}^n$,

$$x^T A x = \sum_{i=1}^n \sum_{j=1}^n A_{i,j} x_i x_j \in \mathbb{R}$$

is called a **quadratic form**.

- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive semi-definite** (PSD) if, for any $x \in \mathbb{R}^n$, $x^T A x \geq 0$.
- A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is **positive definite** (PD) if, for any $x \in \mathbb{R}^n$, $(x \neq 0) \Rightarrow x^T A x > 0$.
- Matrix $A \in \mathbb{R}^{n \times n}$ is PSD \Leftrightarrow all $\lambda_i(A) \geq 0$.
- Matrix $A \in \mathbb{R}^{n \times n}$ is PD \Leftrightarrow all $\lambda_i(A) > 0$.

Agenda

- ~~Probability Theory~~ ✓

- ~~Linear Algebra~~ ✓

Enjoy LxMLS!