1. The probability of a leap year selected at random contain 53 Sunday is:				
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365				
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at				
random. The probability of drawing a black ball is :				
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5				
3. The probability that it will rain tomorrow is 0.85. What is the				
probability that it will not rain tomorrow				
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these				
4. What is the probability that a number selected from the numbers				
(1, 2, 3,,15) is a multiple of 4?				
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3				
5. What are the total outcomes when we throw three coins?				
(a) 4 (b) 5 (c) 8 (d) 7				
6. The probability that a prime number selected at random from the				
numbers (1,2,3,35) is :				
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these				
7. The sum of the probability of an event and non event is :				
(a) 2 (b) 1_{-} (c) 0 (d) none of these.				
8. The following probabilities are given; choose the correct answer				
for that which is not possible.				
(a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.				
9. If three coins are tossed simultaneously, than the probability of				
getting at least two heads, is:				
(a) 1/4 (b) 3/8 (c) ½ (d) 1/8				
10. A letter is chosen at random from the letters of the word				
♦ ASSASSINATION ♦. The probability that the letter chosen has:				
(a) 6/13 (b) 7/13 (c) 1 (d) none of these.				
11. A dice is thrown. Find the probability of getting an even number.				
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2				
12. Two coins are thrown at the same time. Find the probability of				
getting both heads.				
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0				
13. Two dice are thrown simultaneously. The probability of getting a				

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9_	(D) 4/9		
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.					
(A) 3/4	(B) 27/50	(C) 1/4	(D) 29/10	00	
_	a blue ball is c	louble that of a	blue balls .If the pred ball, then the	-	
taken out at non-defecti	random fromve bulb is:		tive bulbs. One b the probability th	at it is	
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100					
18. What is (A) 1/7	-	ty of getting 53 (C) 2/7	Mondays in a lea (D) 7/366	p year?	
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) $1/26$ (B) $3/26$ (C) $7/52$ (D) $1/13$					
equally likel 1,2,312	ly to come to then the prob	rest pointing to	ng an arrow whic one of the numbe Il point to an odd (D) 5/12	er	
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4					

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:					
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225		
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$					
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10_{-} (B) 6_{-} (C) 8_{-} (D) 7_{-}					
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) 7/50 (B) 4/25 (C) 1/25 (D) 2/25					
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?					
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$		
	ored by batsman in The standard devia	_	nes are 50, 70, 82,		
	b) 25.49		d) 25.69		
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.					
	b) 13, 18		d) 13, 16		
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is					
		c) ¹ / ₄	d) $^{1}/_{6}$		
	iate between 0 an b) 7		E(X²) is <mark>) 9</mark>		
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?					

32.Out of the following values, which one is not possible in probability?					
a) $P(x) = 1$ c) $P(x) = 0.5$	b) ∑ x P(d) P(x)	(x) = 3 = - 0.5			
33.If E(X) =	2 and E(z) = 4, t b) 6	c) 0	•	sufficient data	
34.The cova	ariance of two in	dependent	random variab	le is	
a) 1	b) 0	c) - 1	d) Un	defined	
35.If Σ P(x) a) 0	b) 1			sufficient data	
• • •	0.5 and x = 4, th b) 0.5	, ,	d) 2		
37.In a discrete probability distribution, the sum of all probabilities is always?					
•	b) Infinite	c) 1_	d) Undefined		
38.If the probability of hitting the target is 0.4, find mean and					
variance. a) 0.4, 0.24	b) 0.6, 0.	24	c) 0.4, 0.16	d) 0.6, 0.16	
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6					
40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution?					


c) 5

d) 7

a) 3

b) 4

a) Mean is 0 and variance is 1 $_{-}$ b) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0							
42.\ a) E(/ariance (X)	of a rand b) E(X	om varia (2)	nble X is c) E(X2	given by <u>2</u>) – (E(X))	 <mark>2</mark>	d) (E(X))2
43.Mean of a random variable X is given by a) E(X)							
44.N a) 0	lean of a	a constan b) a	t 'a' is	c) a/2	_ ·	d) 1	
45.Variance of a constant 'a' is . a) 0							
46.Find the mean and variance of X?							
	Х	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2,	4/3	b) 3	3, 4/3		c) 2, 2/3		d) 3, 2/3
47. Find the expectation of a random variable X?							



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}