

رياضيات 2

9:11

الثلاثاء 22/6/2021

أ.د. فتحى هشام



Faculty of Computers & Information, Assiut University

1st Level

Final Exam

Duration: 2 hours

1

* الإسم الرباعي (بالعربي فقط)

نرمين محب خير عوض الله

2

* رقم الجلوس

162020677

3

* المستوى

- ☒ الاول
- ☐ الثاني
- ☐ الثالث
- ☐ رابعة 2013
- ☐ رابعة 2014
- ☐ رابعة 2015
- ☐ رابعة 2016
- ☐ رابعة 2017

4

* البرنامج

- ☒ عام
- ☐ بايو
- ☐ هندسة

5

* رقم المعمل

- ☐ ج.
- ☐ د.
- ☐

- ☐ ا ب
- ☐ ا د
- ☐ ا هـ
- ☐ ا٣
- ☐ ا٢ ب
- ☐ ا٢ ج
- ☐ ا٢ د
- ☐ ا٢ هـ
- ☐ ا٣
- ☐ ا٣ ب
- ☒ ا٣ ج
- ☐ ا٣ د
- ☐ ا٣ هـ
- ☐ ا٤
- ☐ ا٤ ب

6

* رقم الكمبيوتر

19

7

* الكود (قد تمت مراجعة بيانات الطالب ورقم الجلوس)

8

السؤال

(2 Points)

The partial fraction decomposition of $\frac{2}{x^2-1}$ is

(a) $\frac{1}{x-1} + \frac{1}{x+1}$

(b) $\frac{1}{x-1} - \frac{1}{x+1}$

(c) $\frac{1}{x+1} - \frac{1}{x-1}$

☐ a

☒ b

☐ c

9

السؤال

(2 Points)

Which statement about the set S is false where $S = \{(1,1,3), (2,3,7), (2,1,1)\}$

a. The set S is linearly independent.

b. The set S contain an element which is solution of the equation $5x_1 + 3x_2 + 7x_3 = 1$.

c. The set S contain two elements, which are multiple of each other.

d. The set S is linearly dependent.

☒ a

☐ b

☐ c

☐ d

10

السؤال
(2 Points)

The augmented matrix of the system of equations

$$2x + 3y - 2z = 0$$

$$6x + 6y + 4z = 6$$

$$-4x - 9y + 6z = 2$$

in reduced row echelon form is

(a) $\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1/3 \\ 0 & 0 & 1 & 2 \end{array} \right]$

(b) $\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1/3 \\ 0 & 0 & 1 & 1/2 \end{array} \right]$

(c) $\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -1/2 \end{array} \right]$

(d) none of these.

☐ a

☒ b

☐ c

☐ d

11

السؤال
(2 Points)

The augmented matrix of the system of equations

$$x + y + z = a$$

$$x + 2y + 2z = 0$$

$$2x + 3y + 3z = a$$

in reduced row echelon form is

(a) $\left[\begin{array}{ccc|c} 1 & 0 & 0 & -2a \\ 0 & 1 & 1 & -a \\ 0 & 0 & 0 & 0 \end{array} \right]$

(b) $\left[\begin{array}{ccc|c} 1 & 0 & 0 & -2a \\ 0 & 1 & 1 & -a \\ 0 & 0 & 1 & 0 \end{array} \right]$

(c) $\left[\begin{array}{ccc|c} 1 & 0 & 0 & -2a \\ 0 & 1 & 0 & -a \\ 0 & 0 & 0 & 0 \end{array} \right],$

(d) none of these.

☐ a

☐ b

☐ c

☒ d

12

السؤال
(2 Points)

Is the function $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x^2 + y^2 \\ x y \end{bmatrix}$ is

- (a) a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
- (b) not a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
- (c) not applicable.

☐ a

☒ b

☐ c

13

السؤال
(2 Points)

Is the set $U = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : 2x + 3y = 0 \right\}$

- (a) a subspace of \mathbb{R}^2
- (b) not a subspace of \mathbb{R}^2
- (c) not applicable.

☐ a

☒ b

☐ c

14

السؤال
(2 Points)

Let V be a vector space, and let S be a subset of V . What does it mean we say that S is linearly independent?

- a. All the elements of S are distinct from each other.
- b. S has nullity zero.
- c. The only way to write 0 as a linear combination of elements of S is the trivial combination (where one takes zero multiples of each element of S).
- d. S is closed under both addition and scalar multiplication.

☐ a

☐ b

☒ c

☐ d

15

السؤال
(2 Points)

If the order of matrix A is $m \times p$ and the order of B is $p \times n$. Then the order of matrix AB is

- (a) $m \times n$ (b) $n \times m$ (c) $n \times p$ (d) $m \times p$

☒ a

☐ b

☐ c

☐ d

السؤال
(2 Points)

State the type of partial fraction $\frac{6x + 5}{(2x - 1)^2}$

a. linear factor.

b. repeated factor.

c. quadratic factor.

d. improper fraction.

☐ a

☒ b

☐ c

☐ d

السؤال
(2 Points)

The system of equations

$$x - y - z = 4$$

$$2x - 2y - 2z = 8$$

$$5x - 5y - 5z = 20$$

has:

a) Unique Solution

b) No solution

c) Infinitely many Solutions

d) Finite solutions

☐ a

☐ b

☒ c

☐ d

السؤال
(2 Points)

The reduced form of the Matrix in Gauss Elimination method is also

- a) Column Echelon Form
- b) Row-Column Echelon Form
- c) Column-Row Echelon Form
- d) Row Echelon Form

☐ a

☐ b

☐ c

☒ d

السؤال
(2 Points)

Is the function $\langle u, v \rangle = 5u_1v_1 + 4u_2v_2$

(a) inner product on R^2

(b) not inner product on R^2

(c) not applicable .

☒ a

☐ b

☐ c

20

السؤال
(2 Points)

If $A = \begin{bmatrix} -1 & -1 & 2 \\ 2 & 1 & -2 \\ 1 & 1 & -1 \end{bmatrix}$, then $A^{-1} =$

(a) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & -1 & 2 \\ 1 & 1 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 1 & 0 \\ 1 & -1 & 2 \\ 1 & 0 & 1 \end{bmatrix}$

(c) $\begin{bmatrix} 1 & 1 & 0 \\ 0 & -1 & 2 \\ 1 & 0 & 1 \end{bmatrix}$

(d) none of these.

☐ a

☐ b

☒ c

☐ d

21

السؤال
(2 Points)

What is true regarding Determinant of a Matrix?

- a. The concept of determinant is applicable to square matrices only.
- b. To find determinant, subtract diagonal elements together.
- c. determinant is a vector value that can be computed from the elements of a square matrix.
- d. Both a and c

- ☒ a
- ☐ b
- ☐ c
- ☐ d

22

السؤال
(2 Points)

The element a_{ij} of any matrix A is present in

- (a) i^{th} row and j^{th} column
- (b) i^{th} column and j^{th} row
- (c) $(i + j)^{th}$ row and column
- (d) $(i - j)^{th}$ row and column

- ☒ a
- ☐ b
- ☐ c
- ☐ d

السؤال
(2 Points)

If determinant of a matrix is equal to zero, then it is sa

- (a) square matrix
- (b) singular matrix
- (c) non-singular matrix
- (d) identical matrix

- ☐ a
- ☒ b
- ☐ c
- ☐ d

السؤال
(2 Points)

Solve the equations using Gauss Jordan method.

$$2x - 3y + z = -1$$

$$x + 4y + 5z = 25$$

$$3x - 4y + z = 2$$

- (a) $x = 1, y = 3, z = 4$
- (b) $x = 1, y = 3, z = 5$
- (c) $x = 1, y = 3, z = 7$
- (d) $x = 1, y = 3, z = 2$

- ☐ a
- ☒ b
- ☐ c

☐ d

25

السؤال
(2 Points)

The aim of elimination steps in Gauss elimination method is to 1
the coefficient matrix to

- a) diagonal
- b) identity
- c) lower triangular
- d) upper triangular

☐ a

☐ b

☐ c

☒ d

26

السؤال
(2 Points)

Let $A; B; C$ be square invertible matrices satisfying $AB = B^2C$.
that $\det B = 3$ and $\det C = 2$. Find a formula for A and calculate
determinant of A .

- a. $A = BC, \det A = 6$.
- b. $A = B^3C, \det A = 11$.
- c. $A = B^2CB^{-1}, \det A = 6$.
- d. $A = B^2CB^{-1}, \det A = 5$.

- ☐ a
- ☐ b
- ☒ c
- ☐ d

27

السؤال
(2 Points)

Solve the equations using Gauss Jordan method.

$$x + 2y + 6z = 22$$

$$3x + 4y + z = 26$$

$$6x - y - z = 19$$

- (a) $x = 2, y = 3, z = 4$
- (b) $x = 4, y = 2, z = 3$
- (c) $x = 4, y = 3, z = 2$
- (d) $x = 3, y = 4, z = 2$

- ☐ a
- ☐ b
- ☒ c
- ☐ d

السؤال
(2 Points)

What is 'a', if $B = \begin{bmatrix} 1 & 4 \\ 2 & a \end{bmatrix}$ is a singular matrix ?

- (a) 5 (b) 6 (c) 7

- ☐ a
☐ b
☐ c
☒ d

السؤال
(2 Points)

If $\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} A = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}$, then order of matrix A

- (a) 2 x 2 (b) 2 x 3 (c) 3 x 2 (d)

- ☐ a
☒ b
☐ c
☐ d

السؤال
(2 Points)

The augmented matrix of the system of equations

$$x_1 + 2x_2 + 3x_3 = 0$$

$$2x_1 - x_2 + x_3 = 0$$

$$x_1 - 5x_2 - 4x_3 = 0$$

in the row-echelon form is

(a) $\left[\begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]$

(b) $\left[\begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$

(c) $\left[\begin{array}{ccc|c} 1 & 2 & 3 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 3 \end{array} \right]$

(d) none of these.

☐ a

☒ b

☐ c

☐ d

السؤال
(2 Points)

Which of the following equations is a variable separable

a. $(x + x^2 y)dy = (2x + xy^2)dx$

b. $(x + y)dx - 2ydy = 0$

c. $2ydx = (x^2 + 1)dy$

d. $y^2 dx + (2x - 3y)dy = 0$

☐ a

☐ b

☒ c

☐ d

32

السؤال
(2 Points)

The correct form of the partial fraction decomposition for $\frac{1}{x^3 - 5x^2 - 6x}$

(a) $\frac{A}{x} + \frac{B}{x^2 - 5x - 6}$

(b) $\frac{A}{x} + \frac{Bx + C}{x^2 - 5x - 6}$

(c) $\frac{A}{x} + \frac{B}{x - 6} + \frac{C}{x}$

☐ a

☐ b

☒ c

33

السؤال
(2 Points)

Is the set $S = \{1, x + 1, x^2 - x\}$ a base for

(a) R^2

(b) P_2

(c) P_3

☐ a

☒ b

☐ c

34

السؤال
(2 Points)

Is the function $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \cos x \\ \sin y \end{bmatrix}$ is

- (a) a linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$
- (b) a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
- (c) a linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$
- (d) none of these.

- ☐ a
- ☐ b
- ☐ c
- ☒ d

35

السؤال
(2 Points)

The solution (x, y, z) of the system

$$\begin{aligned} 2x + y + z &= -6 \\ x + 5y + 2z &= 3 \\ -3x + 2y + 3z &= -3 \end{aligned}$$

is

- (a) $(2, -3, 5)$.
- (b) $(-2, 3, 5)$.
- (c) $(-2, 3, -5)$.
- (d) none of these.

- ☐ a
- ☐ b
- ☒ c
- ☐ d

السؤال
(2 Points)

In Gaussian elimination method, original equations are transformed

- a) Column operations
- b) Row Operations
- c) Mathematical Operations
- d) Subset Operation

- ☐ a
- ☒ b
- ☐ c
- ☐ d

السؤال
(2 Points)

Solve the differential equation $dy - xdx = 0$ when $y(1)$

- a. $3x^2 + 2y - 3 = 0$
- b. $2y^2 + x^2 - 1 = 0$
- c. $x^2 - 2y - 1 = 0$
- d. $2x^2 + 2y - 2 = 0$

- ☐ a

☐ b

☒ c

☐ d

38

السؤال
(2 Points)

With $B = \begin{bmatrix} -1 & 2 \\ 2 & 1 \\ -1 & 2 \end{bmatrix}$, $C = \begin{bmatrix} 0 & -2 \\ 3 & 1 \\ 2 & -3 \end{bmatrix}$, $D = \begin{bmatrix} -1 & 4 & 2 \\ 2 & 0 & 1 \end{bmatrix}$, $E = \begin{bmatrix} 2 & 3 & 1 \\ 2 & 0 & 0 \end{bmatrix}$

We have $(2B - C)(3D - E) =$

(a) $\begin{bmatrix} 34 & -8 & 8 \\ -1 & 9 & 8 \\ 48 & -36 & 1 \end{bmatrix}$

(b) $\begin{bmatrix} 34 & -18 & 8 \\ -1 & 9 & 8 \\ 48 & -36 & 1 \end{bmatrix}$

(c) $\begin{bmatrix} 34 & -18 & 8 \\ -1 & 9 & 8 \\ 48 & -6 & 1 \end{bmatrix}$

(d) none of these.

☐ a

☒ b

☐ c

☐ d

39

السؤال
(2 Points)

The system of equations

$$x_1 + 2x_2 + x_3 = a$$

$$2x_1 + x_2 - x_3 = b$$

$$-4x_1 + x_2 + 5x_3 = c$$

is consistent if

(a) $-2a + 3b + c = 0$

(b) $-2a + 3b + c \neq 0$

(c) $-2a + 3b + 4c = 0$

(d) none of these.

☒ a

☐ b

☐ c

☐ d

40

السؤال
(2 Points)

Let V be a vector space, and let W be a subset of V . What does it mean we say that W is closed under scalar multiplication?

- a. Whenever x is in W and c is a scalar, then cx is in V .
- b. Whenever x is in V and c is a scalar, then cx is in W .
- c. If cx is in W and x is in W , then c is a scalar.
- d. Whenever x is in W and c is a scalar, then cx is in W .

☐ a

☐ b

☐ c

☒ d

41

السؤال
(2 Points)

The correct form of the partial fraction decomposition for $\frac{1}{(x-4)^2(x^2+x+9)}$

is

(a) $\frac{Ax+B}{(x-4)^2} + \frac{Cx+D}{x^2+x+9}$

(b) $\frac{A}{x-4} + \frac{B}{(x-4)^2} + \frac{Cx+D}{x^2+x+9}$

(c) $\frac{A}{(x-4)^2} + \frac{B}{x^2+x+9}$

☐ a

☒ b

☐ c

42

السؤال

(2 Points)

A finite set $S = \{v_1, v_2, v_3\}$ of elements of a linear space V is called for V if:

(a) S is linearly dependent and S spans V

(b) S is linearly independent and S spans V

(c) S is linearly independent and S not spans V

☐ a

☒ b

☐ c

43

السؤال
(2 Points)

State the type of partial fraction $\frac{125 + 4x - 9x^2}{(x-1)(x+3)(x+4)}$

a. <u>linear</u> factor.	b. <u>repeated</u> factor.
c. <u>quadratic</u> factor.	d. <u>improper</u> fraction.

☒ a

☐ b

☐ c

☐ d

44

السؤال
(2 Points)

The equation $y^2 = cx$ is general solution of

a. $\frac{dy}{dx} = \frac{2y}{x}$

b. $\frac{dy}{dx} =$

c. $\frac{dy}{dx} = \frac{y}{2x}$

d. $\frac{dy}{dx} =$

☐ a

☐ b

☒ c

☐ d

45

السؤال
(2 Points)

Is the set $U = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} : a + b = 1 \right\}$

(a) a subspace of R^2

(b) not a subspace of R^2

(c) not applicable .

☐ a

☒ b

☐ c

السؤال
(2 Points)

Is the function $\langle u, v \rangle = 2u_1v_1 + 3u_2v_2 + 4u_3v_3$

- (a) inner product on R^3
- (b) not inner product on R^3
- (c) not applicable .

- ☒ a
- ☐ b
- ☐ c

السؤال
(2 Points)

The system of equations

$$2x_1 - x_2 + x_3 = a$$

$$x_1 + x_2 - x_3 = b$$

$$7x_1 - 2x_2 + 2x_3 = c$$

is consistent if

- (a) $a - 3b + c = 0$
- (b) $a - 3b + c \neq 0$
- (c) $-3a - b + c = 0$,
- (d) none of these.

- ☐ a
- ☐ b
- ☐ c

☒ c

☐ d

48

السؤال
(2 Points)

Solve this system of equations and comment on the nature of the solution using Gauss Elimination method.

$$x + y + z = 0$$

$$-x - y + 3z = 3$$

$$-x - y - z = 2$$

a) Unique Solution

b) No solution

c) Infinitely many Solutions

d) Finite solution

☐ a

☒ b

☐ c

☐ d

49

السؤال
(2 Points)

The base for the subspace $U = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a+b+c=0 \right\}$ of R^3 is:

(a) $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$

(b) $\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$

(c) $\begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$

(d) all of these

☐ a

☐ b

☐ c

☒ d

50

السؤال
(2 Points)

The order of the differential equation $2x^2 \frac{d^2 y}{dx^2} - 3 \frac{dy}{dx} + y =$

a. 2 b. 1 c. 0 d. not

☒ a

☐ b

☐ c

☐ d

51

السؤال
(2 Points)

Let V be the set of all 2-vectors whose components as follows:

$$V = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : x + y = 0 \right\}, \text{ the set } V \text{ is nonempty since :}$$

(a) $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is an element of V

(b) $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$ is an element of V

(c) $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is an element of V

☐ a

☒ b

☐ c

52

السؤال
(2 Points)

Let V be a vector space, and let W be a subset of V . What does it mean we say that W is closed under addition?

a. Whenever x and y are in V , then $x + y$ is in W .

b. Whenever x and y are in W , then $x + y$ is in W .

c. Whenever x and y are in W , then $x + y$ is in V .

d. Every vector in W is the sum of two vectors in W .

☐ a

☒ b

☐ c

☐ d

53

السؤال
(2 Points)

The augmented matrix of the system of equations

$$2x_1 + 4x_2 - 4x_3 = 3$$

$$x_1 + 8x_2 + 2x_3 = 7$$

$$2x_1 + x_2 + x_3 = 2$$

in the row-echelon form is

(a) $\left[\begin{array}{ccc|c} 1 & 5 & 2 & 7 \\ 0 & 1 & \frac{2}{3} & \frac{11}{12} \\ 0 & 0 & 1 & \frac{1}{4} \end{array} \right]$

(b) $\left[\begin{array}{ccc|c} 1 & 8 & 5 & 7 \\ 0 & 1 & \frac{2}{3} & \frac{11}{12} \\ 0 & 0 & 1 & \frac{1}{4} \end{array} \right]$

(c) $\left[\begin{array}{ccc|c} 1 & 8 & 2 & 5 \\ 0 & 1 & \frac{2}{3} & \frac{11}{12} \\ 0 & 0 & 1 & \frac{1}{4} \end{array} \right]$

(d) none of these.

☐ a

☐ b

☐ c

☒ d

54

السؤال
(2 Points)

Solve the given system of equation by Gauss Elimination method.

$$3x + 4y - z = -6$$

$$-2y + 10z = -8$$

$$4y - 2z = -2$$

(a) $(-2, -1, -1)$

(b) $(-1, -2, -1)$

(c) $(-1, -1, -2)$

(d) $(-1, -1, -1)$

☐ a

☐ b

☐ c

☒ d

السؤال
(2 Points)

If A and B are matrices, then which from the following is

(a) $A + B \neq B + A$

(b) $(A^t)^t \neq A$

(c) $AB \neq BA$

(d) all are true

☐ a

☐ b

☒ c

☐ d

السؤال
(2 Points)

Solve the linear differential equation $\frac{dy}{dx} + \frac{y}{x} = x$

a. $xy^2 = \frac{x^3}{4} + C$

b. $xy = \frac{x^4}{4} + C$

c. $x^2y = \frac{x^4}{4} + C$

d. $y = \frac{x^3}{4} + C$

- ☐ a
- ☒ b
- ☐ c
- ☐ d

57

السؤال
(2 Points)

Is the function $T\left(\begin{bmatrix} a \\ b \\ c \end{bmatrix}\right) = \begin{bmatrix} 3a + c \\ b - 2c \end{bmatrix}$ is

- (a) a linear transformation $T: R^3 \rightarrow R^2$
- (b) a linear transformation $T: R^2 \rightarrow R^3$
- (c) not applicable.

- ☒ a
- ☐ b
- ☐ c

This content is created by the owner of the form. The data you submit will be sent to the form owner. Microsoft is not responsible for the privacy or security practices of its customers, including those of this form owner. Never give out your password.

Powered by Microsoft Forms | [Privacy and cookies](#) | [Terms of use](#)