أساسيات رياضيات 9:11 الخميس 1/4/2021 أ.د. طارق محمد عبد اللطيف

Faculty of Computers & Information, Assiut University 1st Level Final Exam Duration: 2 hours

Hi Ziyad, when you submit this form, the owner will be able to see your name and email address.

- * Required
- * الإسم الرباعي (بالعربي فقط) .1

زیاد علی محمد علی

* رقم الحلوس .2

162020252

* المستوي .3



الثاني
الثالث 🔾
الرابع (
4. البرنامج *
عام 💿
بايو
هندسة
* رقم المعمل
٧ ٧
* رقم الكمبيوتر .6
19
7. كود المراقب * كود المراقب
DE14
* (إسم الملف بالمسار (تملي بمعرفة المراقب.8

	Enter your answer
9.	The coefficient of x^4 of the expansion $(x+3/x)^10$ equals 1080 * (1 Point)
	○ True
	False
10.	The summation of the series -1/6+1/6+1/2+··· equals * (2 Points)
	② 2√
	○ 3√
	another answer
1.	For the equation $z^4=2+2i$, the roots take the form * (2 Points)
	$\bigcirc (z_j=2\sqrt{2} \operatorname{cis}(\pi/16$
	$(z_j = (2\sqrt{2})^4 \operatorname{cis}((\pi+j)/16)$
	$(z_j = (2\sqrt{2})^{(1/4)} \operatorname{cis}((\pi/4 + \pi j)/4)$
	another answer
2.	(The polar form of the following complex number $\sqrt{3}$ - i is 2(cos $\pi/6$ -isin $\pi/6$ * (1 Point)
	True

· aisc		False
--------	--	-------

True

42.5 (422) \ (422) \ (422)
13. For the expansion (4x+32)^(1/5) , n equals 5 * (1 Point)
○ True
False
14. The fifth term of the series 100+100(2)+100(2)^2+ is 1500 * (1 Point)
○ True
False
15. The value of any determent may be negative * (1 Point)
○ True
False
16. The number of terms of the expansion (x+3/x)^10 is 10 * (1 Point)
○ True
False
17. The 10^th term of the series 100+100(2)+100(2)^2+ is 100 (3)^10 * (1 Point)



18. For the series 300(1/3)^2+300(1/3)^3+300(1/3)^4++300(1/3)^7, the first term equals * (2 Points)
O 300
O 100
100/3
another answer
19. The summation of infinite terms of the series 300(1/3)^2+300(1/3)^3+300(1/3)^4+ equals * (2 Points)
50
<u> </u>
O 450
another answer
20. Using De Moivre's theorem, cos3θ is * (2 Points)
Cos^3 θ-sin^3 θ
○ cos^3 θ-3cosθsin^2 θ
3sinθ cos^2 θ-sin^3 θ
another answer

21. For $z=(1+i)^10$, then $arg(z)=\pi$ (1 Point)
○ True
False
22. The summation of series -1/6+1/6+1/2+··· equals approximately -0.25 * (1 Point)
○ True
False
23. The summation of the series 100+100(2)+100(2)^2+ equals -50 * (1 Point)
○ True
False
24. The coefficient of x^3 in the expansion (x-2/x)^7 equals * (2 Points)
-280
O 84
another answer
25. If z=((2+i)/(i-1))^3, then z=13/4-9/4 i * (1 Point)
○ True



26. If $A=\{x\in N: x^2-4x-5=0\}$, then A has one element only * (1 Point)
True
○ False
27. By truth table, if A=0 and B=1, then A/B is * (2 Points)
<u> </u>
0
<u> </u>
another answer
28. This series 100+100(2)+100(2)^2+ is convergence * (1 Point)
True
○ False
29. For the expansion (4x+32)^(1/5), x corresponding to x/8 (1 Point)
True
○ False

30. This series -1/6+1/6+1/2+··· is arithmetic * (1 Point)
True
○ False
31. For the complex number z=(2+2i)/i, arg(z) equals * (2 Points)
\bigcirc 7 π /4
\bigcirc 3 π /4
\odot $\pi/4$
 another answer
32. For the system: $ax+3y+5z=8+b,-3z-3y+4x=-14$, $y+ax+z=b$, if $x=-1$, $y=0$ and $z=2$ by Cramer's rule, the value of b equals * (2 Points)
O 1
O 2
○ 3
another answer
33. If z=(2+2i)/i , then z is * (2 Points)
2i-2
○ 2i+2
○ 2i+2-

another answer
34. If we calculate 35^(1/5) by binomial theorem, x equals 2/3 * (1 Point)
○ True
False
35. {N={1, 2, 3,} ∧ Z={a/b: a,b∈N,b≠0 * (1 Point)
○ True
False
36. If A={1, 2} and U={1, 2, 3, 5}, Then ((A)^c)^c equivalent * (2 Points)
A
○ ø
another answer
37. The coefficient of 1/x^5 of the expansion (x-2/x)^7 is * (2 Points)
280

another answer

38	The summation of the series 300(1/3)^2+300(1/3)^3+300(1/3)^4++300(1/3)^7 is * (2 Points)
	49.93
	<u> </u>
	<u>444.8</u>
	 another answer
39.	. [The convergence interval, for the expansion $(4x+32)^{(1/5)}$, is $x \in [-8,8 * (1 Point)]$
	○ True
	False
40	. By De Moivre's theorem, sin4θ equivalent * (2 Points)
	4cos ³ θ-4sin ³ θ
	(4cosθsinθ(cos^2 θ-sin^2 θ)
	4sin^2 θ cos^2 θ-4sin^4 θ
	 another answer
41.	For the system: $ax+3y+5z=8+b$, $-3z-3y+4x=-14$, $y+ax+z=b$, if $x=-1$, $y=0$ and $z=2$ by Cramer's rule, the value of a equals * (2 Points)
	<u> </u>
	O 2

	يف	رق محمد عبد اللط	1/4/2021 أ.د. طار	، 9:11 الخميس
<u> </u>				
another a	answer			
42. The series (1 Point)	1+1/4+(1.3)/(4.8)-	⊦ is a	binomial	series *

- 43. The necessary condition for existing a solution for a linear system is $\Delta \neq 0$ * (1 Point)
 - True

True

Flase

False

Submit

This content is created by the owner of the form. The data you submit will be sent to the form owner. Microsoft is not responsible for the privacy or security practices of its customers, including those of this form owner. Never give out your password.

Powered by Microsoft Forms | Privacy and cookies | Terms of use