

# MCPIP1 expression is positively correlated with an immunosuppressive microenvironment and is associated with aggressive clinicopathological features in glioma patients.

Kenza Miyara<sup>1</sup>, Hamza Benthami<sup>1</sup>, Hayat Miftah<sup>1</sup>, Saadia Ait Ssi<sup>1</sup>, Abdelhakim Lakhdar<sup>2,3</sup>, Abdallah Badou<sup>1,4,5</sup>

<sup>1</sup> Laboratory of Immunogenetics and Human Pathology, Faculty of Medicine and Pharmacy, Hassan II University, Casablanca, Morocco

<sup>2</sup> Neurosurgery Department, University Hospital Center (CHU) Ibn Rochd, Casablanca, Morocco


<sup>3</sup> Research Laboratory on Neurological, Neurosensory Diseases and Disability, Faculty of Medicine and Pharmacy, Hassan II University, Casablanca, Morocco

<sup>4</sup> Mohammed VI Center for Research and Innovation, Rabat, Morocco

<sup>5</sup> Mohammed VI University of Sciences and Health, Casablanca, Morocco

## INTRODUCTION

Glioma is the most common primary brain tumor. Recently, studies have shown that failure of PD1/PDL-1 immunotherapy may be linked to upregulation of various inhibitory immune checkpoints in glioma patients. This highlights the importance of deciphering novel biomarkers for additive or synergistic impact on glioblastoma patients to guide and improve immune-mediated therapy concepts. MCPIP1 known also as Regnase 1 an inflammation-related endoribonuclease, expressed in macrophages, T and B cells could be a good target for immunotherapy.



## AIM OF THE STUDY

Explore the expression, the clinical and immunological significance of the MCPIP1, an emerging immune checkpoint, in human gliomas by exploiting transcriptomic bases.

## MATERIALS & METHODS



International dataset transcriptome of glioma patients

667 Glioma cases




GraphPad Prism 6.0  
Software




R Software

Statistical Analysis


## RESULTS



**Figure 1:** MCPIP1 expression is significantly associated with aggressive clinicopathological parameters of glioma



**Figure 2:** MCPIP1 expression is associated with protumoral cell infiltration and immunosuppressive microenvironment



**Figure 3:** High expression of MCPIP1 is associated with an inhibitory immune microenvironment

## CONCLUSION

MCPIP1 is significantly associated with aggressive clinicopathological parameters of glioma. Its overexpression is positively associated with an inhibitory immune microenvironment as it is also associated with protumoral cell infiltration. Overall, our preliminary results suggest that MCPIP1 could act as a potential immunosuppression-mediated molecule in the glioma microenvironment in several ways.