

## MAG3-99mTC DYNAMIC RENAL SCINTIGRAPHY: A VALUABLE TOOL IN DETECTING INTERMITTENT PYELO-URETERAL JUNCTION SYNDROME - INSIGHTS FROM TWO CASE REPORTS

**K.Bouikhsaine, F.El Fartass, H.Alaoui, C.El Ouatiq, M.A. Bsis, A. Matrane**  
Nuclear Medicine Service, Mohammed VI University Hospital, Marrakech  
e-mail :Kenzabouikhaine@gmail.com

Keywords : Intermittent Pyelo-Ureteral Junction Syndrome, MAG3-99mTC, HOMSY sign.

### INTRODUCTION:

Intermittent pyelo-ureteral junction obstruction in pediatric patients presents as a distinctive clinical condition. The primary causes often involve either external compression, typically from the inferior polar pedicle crossing the pyelo-ureteral junction anteriorly, or an intrinsic abnormality within a segment of the proximal ureter. Dynamic renal scintigraphy serves as a diagnostic tool for investigating urine drainage obstructions. It aids in distinguishing between organic and functional obstructive syndromes while also providing an assessment of the individual functional capacity of each kidney.

**OBJECTIVE:** The contribution of dynamic renal scintigraphy with MAG3 in the diagnosis of intermittent pyeloureteral junction syndrome.

### MATERIAL AND METHOD :


In this report, we highlight two distinct cases :

1. A 13-year-old child who has a medical history marked by recurrent urinary tract infections and presents with moderate right hydronephrosis. Intriguingly, a CT urogram reveals no apparent obstruction.
2. Our second case features a 2-year-old child with a recurrent history of urinary tract infections, showing left hydronephrosis. Remarkably, a CT urogram fails to identify any obstructions in this instance as well.


Dynamic imaging with a gamma camera was performed post intravenous injection of 70.67 MBq (1.91 mCi) for the first patient and 23 MBq (0.62 mCi) for the second, using 99mTc-labeled MAG3. Two phases of imaging, along with 1 mg/kg furosemide at the 20th minute, were conducted. Evaluation included early and late post-voiding static images.

### BIBLIOGRAPHY:

- 1-Yves L. Homsy, Prashant H. Mehta, Denis Huot, Slevin Danais, Intermittent Hydronephrosis: A Diagnostic Challenge, The Journal of Urology, Volume 140, Issue 5, Part 2, 1988, Pages 1222-1226, ISSN 0022-5347, [https://doi.org/10.1016/S0022-5347\(17\)42007](https://doi.org/10.1016/S0022-5347(17)42007).
- 2- Signes d'une possible obstruction pyelo-urétérale à haut débit chez les enfants et les adolescents Sophie Turpin, Diego Barrieras, Marie-Josée Clermont, Julie Franc-Guimond, Anne-Marie Houle, Veronique Phan, Orchidee Djahang, Raymond Lambert CHU Sainte-Justine, Montréal, QC, Canada. Can Urol Assoc J 2017;11(10Suppl7):S364-80. <http://dx.doi.org/10.5489/cuaj.4968>



**Figure 1:** The isotopic renogram of the first patient shows a double delayed HOMSY peak, suggesting intermittent obstruction of the pyeloureteral junction (volume-dependent obstruction).



**Figure 2:** The isotopic renogram for the second patient displays a dual delayed HOMSY peak, indicating intermittent obstruction at the pyeloureteral junction.

### CONCLUSION :

The presence of a double-peaked delayed curve or the HOMSY sign observed in dynamic renal scintigraphy with MAG3 is a rare characteristic of intermittent hydronephrosis. Without treatment, this condition typically progresses to complete obstruction within a few months to a few years.

### RESULTS :

The scintigraphic examination revealed the following:

#### **Case n° 1 :**

The right kidney displayed an increased size, heterogeneous uptake, marked by central and medial photopenic areas with a double-peaked HOMSY delayed image, suggestive of intermittent pyeloureteral junction obstruction (volume-dependent obstruction). The relative renal function (RRF) was assessed at 41.7%.

The isotopic nephrogram of the left kidney appeared normal with a relative renal function evaluated at 58.3%.

#### **Case n° 2 :**

The left kidney exhibited an enlarged size and showed heterogeneous uptake. Notably, it displayed central and medial photopenic areas, which were further confirmed by a double-peaked HOMSY delayed image. These findings collectively suggest intermittent pyeloureteral junction obstruction. The relative renal function (RRF) for the left kidney was assessed at 38%.

In contrast, the isotopic nephrogram of the right kidney appeared to be within normal parameters, with a relative renal function assessed at 62%.

### DISCUSSION:

Isotopic renography was first described in the 1970s to assess upper urinary tract obstruction, and it was categorized into three types based on the renogram appearance: I= Normal Pattern, II= Dilated Obstructive Pattern, IIIa= Dilated-Nonobstructive Pattern, IIIb= Equivocal Patterns. In 1988, a new type, IV, was introduced as the "HOMSY sign," characterized by a characteristic double-peaked delayed curve indicative of intermittent hydronephrosis during non-painful intervals. This sign is rare but plays a crucial role in early diagnosis of intermittent obstruction and, consequently, in therapeutic intervention before progressing to complete and overt obstruction.