

**Rx:229789.1**  
**ISM, LISA**

1200 MAIN ST PO BOX 1500 APT UPSTAIRSPATERSON,NJ 07424

**TAKE 1 TABLET(S) 3 TIMES A DAY THEN GO  
TO THE DOCTOR ON THURSDAY THEN GO  
HOME AND TO BED FOR THE NIGHT**

3.33 Refills Before 31Dec08  
Qty: 30 **LIPITOR 10 MG TABLET**

Generic:ATORVASTATIN CALCIUM 10MG TAB  
Dr. NURSE ISM Use By:

Pfizer Us Pharm

LAP

S/c APM

RX: 229789.1 LAP Disp: 01/26/08  
30 LIPITOR 10 MG TABLET

3.33 Refills Percent AWP Paid:  
71-155-23 ISMNURSE

Total Paid:  
TPPD:\$50.00Cpy: 0.00

Orig: 01/01/08  
Pfizer Us

**Rx: 229789**  
**ISM, LISA**

1200 MAIN ST PO BOX 1500 APT UPSTAIRS  
PATERSON,NJ 07424

Qty:30 LIPITOR 10 MG TABLET

NDC# 71-155-23 (973) 256-5050  
NURSEISM

Auth:

3.33 Refills Before 31Dec08

Dispensed: 01/26/08

**APM**

**Copay: 40.00**

X<sup>-</sup>

N<sub>a</sub>

9<sub>±</sub>

x

**RX: 229789.1**

APM

ISMLISA

1200 MAIN ST PO BOX 1500 APT UPSTAIRSPATERSON,NJ 07424

**TAKE 1 TABLET(S) 3 TIMES A DAY THEN  
GO TO THE DOCTOR ON THURSDAY THEN  
GO HOME AND TO BED FOR THE NIGHT**

3.33 Refills Before 31Dec08  
Qty: 30 LIPITOR 10MG TAB

Dr. NURSE ISM  
1500 CARDINAL DRIVE PO BOX APT

Pfizer Us Pharm

XX0000250

(973) 256-5050

Rx:229789Date: 01/26/08

APM REF# TPPD: \$50.00 Cpy: 0.00  
**ISM, LISA**

1200 MAIN ST PO BOX 1500 APT

HIPPA: Y

**Rx: 229789**  
**ISM, LISA**

1200 MAIN ST PO BOX 1500 APT UPSTAIRS  
PATERSON,NJ 07424

Qty: 30 LIPITOR 10 MG TABLET

NDC# 71-155-23 (973) 256-5050  
NURSE ISM

Auth:

3.33 Refills Before 31Dec08

Dispensed: 01/26/08

**APM**

**Copay: 40.00**

ISM Test Store

1500 Cardinal Drive  
SMITHTOWN,WV 07424  
Phone: 973-256-7633

**LIPITOR 10 MG TABLET**

**IMPORTANT NOTE: The following information is intended to supplement, not substitute for, the expertise and judgment of your physician, pharmacist or other healthcare professional.**

**It should not be construed to indicate that use of the drug is safe, appropriate, or effective for you.**

**Consult your healthcare professional before using this drug.**

**ATORVASTATIN ORAL**

**USES: Atorvastatin is an enzyme blocker (HMG-CoA reductase inhibitor), also known as a "statin".**

**It is used along with a proper diet to help lower cholesterol and fats (triglycerides) in the blood.**

**In general, this drug is prescribed after non-drug treatment options have not been fully successful at lowering cholesterol (e.g., diet change, increase in exercise, weight loss if overweight).**

**Reducing cholesterol and triglycerides help prevent strokes and heart attacks.**

**Atorvastatin is used in adults and children (10 years of age and older).**

**Young girls must have had their first menstrual period before starting this medication.**

**HOW TO USE: Take this medication by mouth usually once daily with or without food, or as directed by your doctor.**

**This drug is best taken in the evening.**

**Dosage is based on your medical condition, response to therapy, and use of certain interacting medicines.**

**Many of the drugs listed in the Drug Interactions section may increase the chances of muscle injury when used with atorvastatin.**

**Consult your doctor or pharmacist for more details.**

**Avoid eating grapefruit or drinking grapefruit juice while being treated with this medication unless your doctor instructs you otherwise.**

**Grapefruit juice can increase the amount of certain medications in your bloodstream.**

**Consult your doctor or pharmacist for more details.**

**If you also take certain other drugs to lower your cholesterol (bile acid-binding resins such as cholestyramine or colestipol), take atorvastatin at least 2 hours after these medications.**

**Use this medication regularly in order to get the most benefit from it.**

**Remember to use it at the same time each day.**

**It may take up to 4 weeks before the full benefit of this drug takes effect.**

**It is important to continue taking this medication even if you feel well.**

**Most people with high cholesterol or triglycerides do not feel sick.**

**SIDE EFFECTS: Headache, diarrhea, stomach/abdominal pain, or joint pain may occur.**

**If any of these effects persist or worsen, notify your doctor or pharmacist promptly.**

**Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects.**

**Many people using this medication do not have serious side effects.**

**This drug may infrequently cause muscle pain, tenderness, or weakness (which can rarely lead to a very serious condition called rhabdomyolysis).**

**Seek immediate medical attention if you develop these muscle problems (especially with fever or unusual tiredness).**

**Tell your doctor immediately if any of these highly unlikely but very serious side effects occur: yellowing eyes and skin, dark urine, severe fatigue, severe stomach/abdominal pain, persistent nausea, change in the amount of urine.**

**A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs.**

**Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing.**

**Rx: 229789**  
**ISM LISA**

973 256 7633