NAME	IMAGE	ACTIVE INGREDIENT & MECHANISM OF ACTION	USE
Abilify 10-15-5		**#Aripiprazol** **Mechanism** #partial agonist at D2 #Potent antagonist at serotonin	-Antipsychotics: -Schizophrenia -Bipolar disorder
Abilia -15-20 "ABILIFY"		#Aripiprazol **Mechanism** #partial agonist at D2 #Potent antagonist at serotonin	-Antipsychotics: -Schizophrenia -Bipolar disorder
Acitretin 10-25		#Acitretin **Mechanism** target specific receptors (RXR and RAR)and normalize the growth cycle of skin cell	Psoriasis
Actos 15-30		#Pioglitazone **Mechanism** -Insulin sensitizer oral anti diabetic -Enhances the transcription of several insulin receptor gen	Anti-diabetic
Acyclovir 400-800		#Acyclovir Acyclovir triphosphate competitively inhibits viral DNA polymerase by acting as an analog to deoxyguanosine triphosphate (dGTP). Incorporation of acyclovir triphosphate into DNA results in chain termination since the absence of a 3' hydroxyl group prevents the attachment of additional nucleosides	acyclovir is used to decrease pain and speed the healing of sores or blisters in people who have varicella (chickenpox), herpes z oster (shingles; a rash that can occur in people who have had chickenpox in the past), and first-time or repeat outbreaks of genital herpes
Adancor 10-20		#Nicorandil **Mechanism** -Potassium channel opener -nitric oxide donor	Anti-ischemic

Adwitine 30-60 "CYMBALTA"		#Duloxetine **Mechanism** -serotonin –norepinephrine reuptake inhibitor -thiophene derivative antidepressant	-Antidepressant -Diabetic peripheral neuropathic pain -Generalized anxiety disorder
Agovald 25		#Agomelatine **Mechanism** -Potent agonist at melatonin receptor -antagonist at serotonin receptors	Antidepressant: Major depressive disorder
anarcol	 8	#Naltrexone **Mechanism** -Opioid receptor antagonist –Competitive antagonist In alcohol Dependence and opioid dependence	#Alcoholism #Opioid dependence
Anastrazol Sandoz "ARIMIDEX"		#Anastrazole **Mechanism** -Aromatase inhibitor -Inhibits the synthesis of estrogen	-Breast cancer -Metastasis in both pre and post-menopausal women
Anastrazol synthon "ARIMIDEX"		#Anastrazole **Mechanism** -Aromatase inhibitor -inhibits the synthesis of estrogen	-Brest cancer -metastasis in both pre and post-menopausal women
Andovimpamide 50-100-200 "LACOVIMP"		#Lacosamide **Mechanism** Enhances slow inactivation of voltage-gated sodium channel resulting in Stabilization of repetitive neuronal firing	Anti-epileptic
Andorivaban 2.5-10-15-20 "XARELTO"		#Rivaroxaban **Mechanism** - Anti platelet -Direct factor XA inhibitor -Blocks the formation of blood clots	-Reduce the risk of stroke and serious blood clots in certain patients with atrial fibrillation - ttt and prevent DVT which can lead to blood clots in the lung – pulmonary embolism-

Angiobradine 5-7.5 "PROCORALAN"		#Ivabradine - Anti ischemic -Hart rate lowering agent that works through selective and specific inhibition of cardiac pacemaker if current	Chronic stable angina in coronary artery disease patient with normal sinus rhythm
Anselacox 60-90 "ARCOXIA"		#Etoricixib **Mechanism** -Selective cox inhibitor	-Rheumatoid arthritis -acute pain -gout -chronic low back pain -osteoarthritis -ankylosing spondylitis
Anvizram 200		#Fvipiravir **Mechanism** -purine nucleoside analogue which act as a competitive inhibitor of RNA-dependent RNA polymerase	Antiviral -influenza A B -avion influenza -ebola virus -lassa virus -COVID-19
Apexidon 0.5-1-2-3-4 Syrup "RISPERDAL"		#Risperidone **Mechanism** -block dopamine and serotonin receptors -has antagonistic action on alpha adrenergic and histaminic receptor	Antipsychotic
Apixatrack 2.5-5 "ELIQUIS"		#Apixaban **Mechanism** - Anti co agulant -direct factor xa inhibitor	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial
Apetoid 20 "AVARA"		#leflunomide **Mechanism** -Immunomodulatory drug -pyrimidine synthesis inhibitor -Anti-inflammatory	-Rheumatoid arthritis -psoriatic arthritis

Artixiban 2.5-5 "ELIQUIS"		#Apixaban **Mechanism** - Anti co agulant -direct factor xa inhibitor **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial
Aromasin 25		#Exemestane **Mechanism** - Anti neoplastic	Advanced breast cancer –adjusvant ttt – after primary or main therapy In postmenopausal women whose disease has progressed after tamoxifen therapy
Aricept 5-10		#Donepezil **Mechanism** -Acetyl cholinesterase inhibitor	Alzheimer disease
Arimidex 1		#Anastrazole **Mechanism** -Aromatase inhibitor -Inhibits the synthesis of estrogen	-Breast cancer -metastasis in both pre and post-menopausal women
Aripiprex 10-30-vial "ABILIFY"		#Aripiprazol **Mechanism** #partial agonist at D2 #Potent antagonist at serotonin	Antipsychotics: -Schizophrenia -Bipolar disorder
Arcoxia 60-90		#Etoricixib **Mechanism** -Selective cox inhibitor	-Rheumatoid arthritis -Acute pain -Gout -Chronic low back pain -Osteoarthritis -Ankylosing spondylitis

Atomafutix 10-25-40 "STRATTERA "		#Atomoxetine **Mechanism** Selective inhibition of presynaptic norepinephrine reuptake in the prefrontal cortex. Atomoxetine has a high affinity and selectivity for norepinephrine transporters, but little or no affinity for various neurotransmitter receptors	Ttt of ADHD Attention deficit hyperactivity disorder
Atomox apex 10-18-25-40-60 "STRATTERA "		#Atomoxetine **Mechanism** Selective inhibition of presynaptic norepinephrine reuptake in the prefrontal cortex. Atomoxetine has a high affinity and selectivity for norepinephrine transporters, but little or no affinity for various neurotransmitter receptors	Ttt of \ADHD Attention deficit hyperactivity disorder
Attensera 18 25 40 "STRATTERA "		#Atomoxetine **Mechanism** Selective inhibition of presynaptic norepinephrine reuptake in the prefrontal cortex. Atomoxetine has a high affinity and selectivity for norepinephrine transporters, but little or no affinity for various neurotransmitter receptors	Ttt of ADHD Attention deficit hyperactivity disorder
Arthfree 20 "AVARA"		#leflunomide **Mechanism** -Immunomodulatory drug -Pyrimidine synthesis inhibitor -Anti-inflammatory	-Rheumatoid arthritis -psoriatic arthritis
Atoreza 10l10 20l10 40l10 "ATOZET "		#Atorvastatin+ Ezetimibe **Mechanism** Atorvastatin -HMG-CoA reducates Inhibitor statins antilipemic agent - Ezetimibe reduce the absorption of cholesterol from the intestine	- combined hyperlipidemia -homozygous familial hyperlipidemia

Atozet 10l10 20l10 40l10		#Atorvastatin+ Ezetimibe **Mechanism** Atorvastatin -HMG-Coa reductase Inhibitor statins antilipemic agent - Ezetimibe reduce the absorption of cholesterol from the intestine	- combined hyperlipidemia -homozygous familial hyperlipidemia
Atozemb 10l10 20l10 "ATOZET "		#Atorvastatin+ Ezetimibe **Mechanism** Atorvastatin -HMG-Coa reductase Inhibitor statins antilipemic agent - Ezetimibe reduce the absorption of cholesterol from the intestine	- combined hyperlipidemia -homozygous familial hyperlipidemia
Avodart 0.5		#Dutastreride **Mechanism** Selective inhibitor of both type 1 and type2 5 alpha reductase inhibitor(an intracellular enzyme that converts testosterone to dihydrotestosterone which influences the development of prostate gland)this decreasing level of this hormone relieves the symptoms associated with BPH	-Benign prostatic hyperplasia –BPH- -Male pattern baldness
Avipravir 200		#Fvipiravir **Mechanism** -purine nucleoside analogue which act as a competitive inhibitor of RNA-dependent RNA polymerase	Antiviral -influenza A B -avion influenza -ebola virus -lassa virus -COVID-19
Avara 10-20		#leflunomide **Mechanism** -Immunomodulatory drug -Pyrimidine synthesis inhibitor -Anti-inflammatory	-Rheumatoid arthritis -psoriatic arthritis

Betmiga 50		Mirabegrone **Mechanism** Relaxes the detrusor smooth muscle during the storage phase of the urinary bladder fill-void cycle by activation of beta-3 AR which increases bladder capacity	Over active bladder (urge urinary incontinence urgency frequency)
Bladogra 50 "BETMIGA"		Mirabegrone **Mechanism** Relaxes the detrusor smooth muscle during the storage phase of the urinary bladder fill-void cycle by activation of beta-3 AR which increases bladder capacity	Over active bladder (urge urinary incontinence urgency frequency)
Brilique 60-90		#Ticagrelor **Mechanism** Platelet aggregation inhibitor Works by slowing or stopping platelet from sticking to blood vessel walls or injured tissues	-Reducing the risk of stroke heart attack -or death in certain patient who have had a heart attack or who have angina (chest pain) -it should be used along with aspirin as directed by your doctor
Bradipect 5-7.5 "PROCORALAN "		#Ivabradine **Mechanism** - Anti ischemic -Hart rate lowering agent that works through selective and specific inhibition of cardiac pacemaker if current	Chronic stable angina in coronary artery disease patient with normal sinus rhythm
Brintellix 10-20		#Vortioxetine **Mechanism** Atypical antidepressant	Ttt Major depressive disorder
Budexan inh 1-0.5		#Budesonide **Mechanism** is a potent topical anti-inflammatory agent It binds and activates glucocorticoid receptors (GR) in the effector cell (e.g., bronchial)	Budesonide is used to control and prevent symptoms (Wheezing and shortness of breath) caused by asthma. This medication belongs to a class of drugs known as corticosteroids. It works directly in the lungs to make breathing easier by reducing the irritation and swelling of the airways

Casodex 50		#Bicalutamide **Mechanism** Non-steroidal anti-estrogen anti-neoplastic -anti androgenic drug binds to receptor of androgen on prostate gland and prevent growth of androgen stimulated prostate cancer	Antineoplastic Prostate cancer
Carcemia 100 400		#Imatinib **Mechanism** Anti-cancer Tyrosine-kinase inhibitor	- chronic myelogenous leukemia -lymphoblastic leukemia -gastrointestinal stromal tumors -systemic mastocytosis -hyperosinophilic syndrome -plexiform neurofibromas -dermatofibrosarcoma protuberans
Carfalone 25-50		#Eplerenone **Mechanism** Aldosterone receptor antagonist similar to spironolactone . Eplerenone is in a class of medications called mineralocorticoid receptor It works by blocking the action of aldosterone, a natural substance in the raises blood pressure	is used alone or in combination with other medications to treat high blood pressure antagonists. body that
Cardura 1-4-dxl		#Doxazosin **Mechanism** -Selective alpha-1 antagonist -it dilates both arteries and veins -the alpha 1 blockade in non-vascular smooth muscle cause relaxation of smooth muscle of prostate	- Anti hyper tension - Benign prostatic hypertrophy
Celebrex 200-100		#Celecoxib **Mechanism** - selective Cox-2 inhibitor	1-osteoarthritis 2-rheumatoid arthritis 3-acut gout

Cellcept 250- 500		**#Mycophenolate mofetil** **Mechanism** Immunosuppressant	Prophylaxis of organ rejection in patient receiving allogeneic renal –cardiac or hepatic transplants
Champix 1 0.5		#varenicline **Mechanism** #nicotinic agonist	Smoking cessation
Cipla tropium 18 mcg "SPIRIVA"		#Tiotropium bromide **Mechanism** Anticholinergic (muscarinic antagonist) -bronchodilator	-chronic obstructive pulmonary disease(COPD) -bronchospasm -chronic bronchitis -emphysema
Cipralex10		#Escitalopram **Mechanism** SSRI-selective serotonin reuptake inhibitor	Depression
Cialis 5-20		Tadalafil	Ttt eriction dysfunction
Cipram 20		#Citalopram **Mechanism** SSRI-selective serotonin reuptake inhibitor	Depression
Cludine tech 0.5-1 "TECAVIR"		#Entecavir **Mechanism** Entecavir is a nucleoside analog, or more specifically, a deoxyguanosine analogue that belongs to a class of carbocyclic nucleosides and inhibits reverse transcription, DNA replication and transcription in the viral replication process	Ttt chronic hepatitis b virus

Concor amlo		Bisolol+ amlodipine	Ant hypertension
Constipride 1-2 "RESOLOR"		#Prucalopride **Mechanism** selective high affinity serotonin receptor agonist which targets the impaired motility associated with chronic constipation thus normalize bowel movement -it stimulates colonic mass movement	Ttt chronic constipation
Cotareg 80l25 160l12.5 160l25 320l12.5 320l25		Valsartan Hydroclorothiazide	Anti-hypertension
Coveram 5l5 5l10 10l10 10l5		Amlodipine + Perindopril	Anti-hypertension
Cortexin		Cortexin Iorn Biotin Zinc Floic acid Vitamin c Vitamin d3 Vitamin b1 Vitamin b2 Vitamin b3 Vitamin b5 Vitamin b6 Vitamin b12 Copper	

Crestor 5-10-20		#Rosuvastatin **Mechanism** HMG-COA reductase inhibitor - reduces cholesterol synthesis up to 50%	Anti-hyper lipidemic
Cymbalta 60-30		#Duloxetine **Mechanism** -serotonin –norepinephrine reuptake inhibitor -thiophene derivative antidepressant	-Antidepressant -diabetic peripheral neuropathic pain -generalized anxiety disorder
Cymbatex 20-30-60 "CYMBALTA"		#duloxetine **Mechanism** -serotonin –norepinephrine reuptake inhibitor -thiophene derivative antidepressant	-Antidepressant -diabetic peripheral neuropathic pain -generalized anxiety disorder
Dapaglif "FORXIGA"		#Dapagliflozin **Mechanism** *#Dapagliflozin* inhibits subtype 2 of the sodium-glucose transport proteins (SGLT2) which are responsible for at least 90% of the glucose reabsorption in the kidney. Blocking this transporter mechanism causes blood glucose to be eliminated through the urine	Antidaibetic
Diaflozmet 5l1000 10l1000 10l500 5l500 "XIGDUO "		#dapagliflozin +metform **Mechanism** *#Dapagliflozin* inhibits subtype 2 of the sodium-glucose transport proteins (SGLT2) which are responsible for at least 90% of the glucose reabsorption in the kidney. Blocking this transporter mechanism causes blood glucose to be eliminated through the urine *#Metformin* in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Antidaibetic

Diacurimap 10- 25		#Empagiliflozin **Mechanism** *#Empagliflozin* works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition , empagliflozin reduces renal reabsorption of glucose and increases urinary excretion of glucose	Antidaibetic
Diglifloz 10-5 "FORXIGA"		#Dapagliflozin **Mechanism** *#Dapagliflozin* inhibits subtype 2 of the sodium-glucose transport proteins (SGLT2) which are responsible for at least 90% of the glucose reabsorption in the kidney. Blocking this transporter mechanism causes blood glucose to be eliminated through the urine	Anti-diabetic
Diglifloz plus 5/1000 "XIGDUO "		#dapagliflozin+Metformin **Mechanism** *#Dapagliflozin* *#Metformin* in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Anti-diabetic
Dalfarosis e.r		#Dalfampridine **Mechanism** - Nonspecific potassium channel blocker which improves conduction in focally demyelinated axons by delaying repolarization and prolonging the duration of action potential -enhanced neuronal conduction is through to strengthen skeletal muscle fiber twitch activity -improving peripheral motor neurologic function	Ttt to improve walking in patient with multiple sclerosis(MS)

Donhimer 5 "ARICEPT"		# Donepezil **Mechanism** -Acetyl cholinesterase inhibitor	Alzheimer disease
Duloxeprin 30-60 "CYMBATEX"		#Duloxetine **Mechanism** -serotonin –norepinephrine reuptake inhibitor -thiophene derivative antidepressant	-Antidepressant -diabetic peripheral neuropathic pain -generalized anxiety disorder
Duodart 0.5\0.4		#Tamsulin+ Dutasteride **Mechanism** *#Dutasteride*: 5 alpha reductase inhibitor -testosterone is responsible for prostate hyper trophy Blocking this enzyme prevents prostate from further growing *#Tamsulin*: block alpha receptor in the prostate and bladder so relax the smooth muscles of prostate	Ttt benign prostatic hyperplasia -for preventing urinary retention
Duphaston10		#Dydrogesterone **Mechanism** Dydrogesterone is a progestogen that works by regulating the healthy growth and normal shedding of the womb lining by acting on progesterone receptors in the uterus. Rapidly absorbed in the gastrointestinal tract with a bioavailability of 28%	Threatened or recurrent miscarriage during pregnancy, dysfunctional bleeding , infertility due to luteal insufficiency, dysmenorrhea.
Ebixa		# Memantine **Mechanism** n-methyl –d-aspartate receptor antagonist (NMDA) -dopamine d2 agonist -xss glutamate plays a role in neural cell death	Antidementia agent in Alzheimer disease
Efexor xr 75-150		#venlafaxine **Mechanism** Serotonin norepinephrine reuptake inhibitor	-Depression -generalized anxiety disorder -social anxiety

Effegad 75-150 "EFEXOR"		#venlafaxine **Mechanism** Serotonin norepinephrine reuptake inhibitor	-Depression -generalized anxiety disorder -social anxiety
Eliquis5-2.5		#Apixaban **Mechanism** - Anti co- agulant -direct factor xa inhibitor **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin 	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial fibrillation
Elimbosis 2.5-5 "ELIQUIS"		#Apixaban **Mechanism** - Anti co- agulant -direct factor xa inhibitor **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial fibrillation
Empacoza 10-25 "JARDIANCE"		Empagiliflozin **Mechanism** Empagliflozin works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition, empagliflozin reduces renal reabsorption of glucose and increases urinary excretion of glucose	Anti-diabetic

Empaglimax 25-10 "JARDIANCE"		#Empagiliflozin **Mechanism** Empagliflozin works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition, empagliflozin reduces renal reabsorption of glucose and increases urinary excretion of glucose	Anti-diabetic
Empagliform 5l500 12.5l500 "SYNJARDY"		#Empagliflozin Metformin **Mechanism** *#Empagliflozin* *#Metformin* in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Anti-diabetic
Em ex		#Granisetron **Mechanism** Selective 5ht3 receptor antagonist	Nausea and vomiting associated with cytotoxic chemotherapy or radiotherapy
Emend 125mg 1 cap +80mg 2 cap		#Aprepitant **Mechanism** Neurokinin1 NK1 receptor blocker	Antiemetic

Entresto 50-100-200		#sacubitril Valsartan **Sacubitril** is a pro-drug that upon activation acts as a neprilysin inhibitor. It works by blocking the **action** of neprilysin thus preventing the breakdown of natriuretic peptides. This leads to a prolonged duration of the favorable effects of these peptides. **Neprilysin** is a key enzyme in the degradation of natriuretic peptides, and the primary rationale for **neprilysin** inhibitor therapy in cardiovascular disease was to increase endogenous natriuretic peptide levels, and thereby achieve the vasodilatation and natriuresis these peptides produce 	Heart failure Used to reduce the risk of cardiovascular death and hospitalization for heart failure in patient with chronic heart failure
Ergoconazol 200 "VFEND"		Voriconazol	Ttt of esophageal candidiasis invasive pulmonary aspergillosis -ttt serious fungal infections caused by scedosporium apiospermum and fusarium spp

Eraloner 25-50		#Eplerenone **Mechanism** Aldosterone receptor antagonist similar to spironolactone . Eplerenone is in a class of medications called mineralocorticoid receptor It works by blocking the action of aldosterone, a natural substance in the raises blood pressure	is used alone or in combination with other medications to treat high blood pressure antagonists. body that
Eslizepine 400 800		#Eslicabrazepine **Mechanism** Anti-epileptic It may work by slowing abnormal nerve impulses in the brain	Ttt partial onset seizures
Exforge 10l160 5l160 10l160l25 htc 5l160l12.5 htc		Amilodipine + Valsartan	Anti-hypertension
Exelon 1.5 3 4.5 6 9 patch 18 patch	 	Rivastagmine Central reversible non competitive anti choline esterase	Mild to moderate dementia of the Alzheimer's
Exeedolip		**#icosapent** **Mechanism** icosapent ethyl) is a type of omega-3 fatty acid only containing EPA, eicosapentaenoic acid (EPA), whereas supplemental **fish oils** contain both EPA and docosahexaenoic acid (DHA). t, **It** is thought to work by decreasing the amount of triglycerides made by the body increased ß-oxidation; inhibition of acyl-CoA:1,2-diacylglycerol acyltransferase (DGAT); decreased lipogenesis in the liver; and increased plasma lipoprotein lipase activity **It** reduces high triglycerides without raising levels of "bad" cholesterol or LDL-C	.It is **used** along with a low-fat and low-cholesterol diet to lower high triglyglyceride level (>500 mgldl) - **cardio vascular risk reduction :** -indicated as an adjunct to maximally tolerated statin therapy to reduce risk of myocardial infarction stroke coronary revasularization and un stable angina requiring hospitalization in adults with elevated TG >150 mgLdl And Established cardiovascular disease Or Diabetes mellitus and 2 or more additional risk factors for cv disease

Faverin 50-100		#Fluvoxamine **Mechanism** SSRI	Ttt OCD Obsessive compulsive disorder -management of bulimia nervosa
Femara 2,5		#Letrozole **Mechanism** Inhibit aromatase and suppress estrogen production -promotes the growth and release of eggs in women who aren't ovulating on their ovulation induction or super ovulation in women who are able to ovulate -estrogen levels are suppressed in women .FSH hormone output is increased by the brain and pituitary gland .FSH can result in the development of a mature follicle in the women's ovary to ovulate an egg	-Breast cancer in post-menopausal women -pre operative ttt in post-menopausal women with localized hormone receptor positive breast cancer -ttt infertility in women
Femaprent 2.5 "FEMARA"		#Letrozole **Mechanism** Inhibit aromatase and suppress estrogen production -promotes the growth and release of eggs in women who aren't ovulating on there ovulation induction or super ovulation in women who are able to ovulate -estrogen levels are suppressed in women .FSH hormone output is increased by the brain and pituitary gland .FSH can result in the development of a mature follicle in the women's ovary to ovulate an egg	--breast cancer in post-menopausal women -pre operative ttt in post-menopausal women with localized hormone receptor positive breast cancer -ttt infertility in women
Flopadex 8		Silodosin **Mechanism** Alpha 1 a blocker	Ttt urination difficulty in benign prostatic hyperplasia

Fosavance		#Alendronate Vd3 **Mechanism** bisphosphonates inhibit osteoclastic bone resorption via a mechanism that differs from that of other antiresorptive agents . Bisphosphonates attach to hydroxyapatite binding sites on bony surfaces, especially surfaces undergoing active resorption	Ttt post menopausal osteoporosis in women at risk of vit d insufficiency -reduce the risk of vertebral and hip fracture
Fosamax		#Alendronate **Mechanism** Bisphosphonates inhibit osteoclastic bone resorption via a mechanism that differs from that of other antiresorptive agents. Bisphosphonates attac h to hydroxyapatite binding sites on bony surfaces, especially surfaces undergoing active resorption	Ttt and prevention of osteoporosis in women and pagets disease of bone in both men and women
Forxiga 5-10		#Dapagliflozin **Mechanism** Dapagliflozin inhibits subtype 2 of the sodium-glucose transport proteins (SGLT2) which are responsible for at least 90% of the glucose reabsorption in the kidney. Blocking this transporter mechanism causes blood glucose to be eliminated through the urine	Antidaibetic
Forflozinb		#Dapagliflozin **Mechanism** Dapagliflozin inhibits subtype 2 of the sodium-glucose transport proteins (SGLT2) which are responsible for at least 90% of the glucose reabsorption in the kidney. Blocking this transporter mechanism causes blood glucose to be eliminated through the urine	Antidaibetic

Fungilimit200 "VFEND"		Voriconazol	Ttt of esophageal candidiasis invasive pulmonary aspergillosis -ttt serious fungal infections caused by scedosporium apiospermum and fusarium spp
Gardovas 5		#Ambrisentan **Mechanism** a potent type-A selective endothelin receptor antagonist. Endothelin is a peptide made by the body in the endothelium and it constricts blood vessels and elevates blood pressure. There are two classes of endothelin receptors: Endothelin A (ET-A) and Endothelin B (ET-B	Ttt pulmonary aterial hypertension
Granitryl 2		#Granisetron **Mechanism** -Selective5ht3 receptor antagonist	Nausea and vomiting associated with cytotoxic chemotherapy or radiotherapy
Galvus 50		#Vildagliptin **Mechanism** inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	Anti-diabetic

Galvus met 50l500 met 50l850 met 50l1000		Vildagliptin Metformin **Mechanism** #*Vildagliptin* *#Metformin* in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Anti-diabetic
Glimpacare		#Dapagliflozin **Mechanism** Dapagliflozin inhibits subtype 2 of the sodium-glucose transport proteins (SGLT2) which are responsible for at least 90% of the glucose reabsorption in the kidney. Blocking this transporter mechanism causes blood glucose to be eliminated through the urine	Anti-diabetic
Icosalip VASCEPA		**#icosapent** **Mechanism** icosapent ethyl) is a type of omega-3 fatty acid only containing EPA, eicosapentaenoic acid (EPA), whereas supplemental **fish oils** contain both EPA and docosahexaenoic acid (DHA). t, **It** is thought to work by decreasing the amount of triglycerides made by the body increased ß-oxidation; inhibition of acyl-CoA:1,2-diacylglycerol acyltransferase (DGAT); decreased lipogenesis in the liver; and increased plasma lipoprotein lipase activity **It** reduces high triglycerides without raising levels of “bad” cholesterol or LDL-C	.It is **used** along with a low-fat and low-cholesterol diet to lower high triglyglyceride level (>500 mgldl) - **cardio vascular risk reduction :** -indicated as an adjunct to maximally tolerated statin therapy to reduce risk of myocardial infarction stroke coronary revasularization and un stable angina requiring hospitalization in adults with elevated TG >150 mgLdl And Established cardiovascular disease Or Diabetes mellitus and 2 or more additional risk factors for cv disease

Iksaront 2.5-5 "ELIQUIS"		#Apixaban **Mechanism** - Anti co agulant -direct factor xa inhibitor - **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial fibrillation
Inhalex 18 "SPIRIVA"		#Tiotropium bromide **Mechanism** Anticholinergic (muscarinic antagonist) -bronchodilator	-chronic obstructive pulmonary disease(COPD) -bronchospasm -chronic bronchitis -emphysema
Inhiglip 12.5 25 "		#Alogliptin **Mechanism** inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	Anti-diabetic

Inegy 10l10 10l20 10l40		#Simvastatin Ezetimibe **Mechanism** *Simvastatin* is in a class of medications called HMG-CoA reductase inhibitors (statins). It works by slowing the production of cholesterol in the body to decrease the amount of cholesterol that may build up on the walls of the arteries and block blood flow to the heart, brain, and other parts of the body. *- ezetimibe* reduce the absorption of cholesterol from the intestine	Hyperlipidaemia
Inviga 3-6 Sustenna 100-75-150		#Paliperidone **Mechanism** Benzisoxazole atypical antipsycatric which is a main active metabolite of risperidone -an antagonist at dopamine-serotonin-adrenergic a1 and a2 receptors and histamine receptors	-schizophrenia -sever depression -anxiety -obsessive thought
Invokana 100 300		#Ganagliflozin **Mechanism** Inhibits the Na-glucose co-transporter 2 (SGLT-2) in the kidney to reduce glucose reabsorption, resulting in increased urinary glucose excretion, and lower plasma glucose	Antidaibetic
Invoglita 100 300 "INVOKANA "		#Ganagliflozin **Mechanism** #Inhibits the Na-glucose co-transporter 2 (SGLT-2) in the kidney to reduce glucose reabsorption, resulting in increased urinary glucose excretion, and lower plasma glucose	Antidaibetic

Januvia 100-50		#Sitagliptin **Mechanism** inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	Antidaibetic
Jardiance 25-10		#Empagiliflozin **Mechanism** Empagliflozin works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition empagliflozin reduces renal reabsorption of glucose and increases urinary excretion of glucose	Antidaibetic

Janumet 50\500 50\850 50\1000		Sitagliptin Metformin **Mechanism** *#Sitagliptin* inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.***#Metformin*** in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Antidaibetic
Karbalta 30-60 "CYMBALTA"		#Duloxetine **Mechanism** -serotonin –norepinephrine reuptake inhibitor -thiophene derivative antidepressant	-Antidepressant -diabetic peripheral neuropathic pain -generalized anxiety disorder

Keppra 250-500-1000 Oral solution		#Levetiracetam **<u>Mechanism</u>** Binds to synaptic vesicle protein SV2A which is believed to impede nerve conduction across synapses	Ttt epilepsy anti convulsant neuropathic pain anxiety disorder
Ketosteril 100		Combination of alpha ketoanalogue of amino acids (l-lysine+l-threonine+ l-tryptophan+l-histidine+l tryrosine	Prevention and therapy of damages duo to faulty or deficient protein metabolism in chronic renal in sufficiency
Kombiglyze xr 2.5\1000 5\1000		Saxagliptin + metformin **Saxagliptin** inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.[1] ***<u>Metformin</u>*** in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Antidaibetic

Lamictal 2-25-50-100		#Lamotrigine **mechanism** Prolongs the in activation state of voltage sensitive neuroneal sodium ion channel	Anti-epileptic
Lacovimp 50-100		#Lacosamide **mechanism** Enhances slow inactivation of voltage-gated sodium channel resulting in Stabilization of repetitive neuronal firing	Anti-epileptic
Lescol xl 80		Fluvastatin inhibition of HMG-CoA (hydroxymethylglutaryl-coenzyme A) reductase, the rate-limiting enzyme in the cholesterol biosynthesis pathway	
Letrozol 2.5 "FEMARA"		#Letrozole **mechanism** Inhibit aromatase and suppress estrogen production -promotes the growth and relase of eggs in women who aren't ovulating on their ovulation induction or super ovulation in women who are able to ovulate -estrogen levels are suppressed in women .FSH hormone output is increased by the brain and pituitary gland .FSH can result in the development of a mature follicle in the women's ovary to ovulate an egg	-Breast cancer in post-menopausal women -pre operative ttt in post-menopausal women with localized hormone receptor positive breast cancer -ttt infertility in women
Levepex 500-1000-750 "KEPPRA"		#Levetiracetam **mechanism** Binds to synaptic vesicle protein SV2A which is believed to impede nerve conduction across synapses	Ttt epilepsy anti convulsant neuropathic pain anxiety disorder
Lipitor 10-20-40-80		Atorvastatin	

Lustral		#Sertrakine **Mechanism** SSRI	Ttt depression
Maddovit a-z			Multivitamins
Mapi		Vitamin b1 Vitamin b2 Vitamin b3 Vitamin b6 Vitamin b12 Folic acid Fish oil EPA DHA	
Mestinon 60		#Pyridostigmin bromide **Mechanism** Para sympathomimetic and reversible cholinesterase inhibitor	Ttt myasthenia gravis and reverse the action of muscle relaxant
Mellitofix met 12.5l500 12.5l1000 "SYNJARDY"		#Empagliflozin Metformin **Mechanism** *#Empagliflozin* works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition, empagliflozin reduce s renal reabsorption of glucose and increases urinary excretion of glucose *# Metformin* in patients with type 2 diabetes mellitus Setting is inhibition of hepatic gluconeogenesis.	Anti-diabetic

Mellitofix 25 10 "JARDIANCE"		#Empagiliflozin **Mechanism** *#Empagliflozin* works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition, empagliflozin reduce s renal reabsorption of glucose and increases urinary excretion of glucose	Anti-diabetic
Micaedis plus 40\12.5 plus 80\12.5		#Telmisatran Hydroclorothiazide **Mechanism** telmisartan blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II by selectively blocking the binding of angiotensin II to the AT1 receptor in many tissues, such as vascular smooth muscle and the adrenal gland	Anti-hypertensive
Micaedis 80 40		#Telmisatran **Mechanism** #Telmisatran blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II by selectively blocking the binding of angiotensin II to the AT1 receptor in many tissues, such as vascular smooth muscle and the adrenal gland	Anti-hypertensive
Minirin melt 60-120MCG		#Desmopressin **Mechanism** Synthetic analogue of natural pituitary hormone 8-arginine vasopressin ADH antidiuretic hormone	Urinary incontinence antidiuretic

Modasomil 200		Modafinil **Mechanism** Most research on modafinil's wake-promoting mechanism has focused on monoaminergic effects showing modafinil stimulates histamine (HA), norepinephrine (NE), serotonin (5-HT), dopamine (DA), and orexin systems in the brain, but researchers have not been able to isolate a single site of action or locate major receptor	Wakefulness disorders – narcolepsy -shift work sleep disorder -excessive daytime sleepiness
Myogrel Myogrel plus		Clopidogrel + acetylsalicylic acid	
Nashilive 5 10		#Obeticholic acid **Mechanism** Suppress the bile acid production in the liver and increase the bile flow	To treat primary cholangitis-a type of liver disease that destroys the bile ducts which allows bile to stay in the liver and cause damage
Naviluca 200		Fluconazole	Antifungal
Napibradine 5-7.5 "PROCORALAN "		#Ivabradine - Anti ischemic -Hart rate lowering agent that works through selective and specific inhibition of cardiac pacemaker if current	Chronic stable angina in coronary artery disease patient with normal sinus rhythm

Nexium 40-20 Vial Sach		Esomeprazole	Ppi
Netlook 10 20 40 "ROACCUTAN "		Isotretinoin	Anti acne
Olapex5-10 "ZYPREXA"		#Olanzapine **Mechanism** The exact mechanism of action of olanzapine is not known. It may work by blocking receptors for several neurotransmitters (chemicals that nerves use to communicate with each other) in the brain. It binds to alpha-1, dopamine, histamine H-1, muscarinic, and serotonin type 2 (5-HT2) receptors	For acute and maintenance ttt of schizophrenia
Omnic 0.4 Modified release		#Tamsuline **Mechanism** Selective alpha 1 adrenergic receptor antagonist In the prostate And leads to relaxation of smooth muscle in the bladder neck and prostate	Bhp
Omnic ocas 0.4 Prolonged release		#Tamsuline **Mechanism** Selective alpha 1 adrenergic receptor antagonist In the prostate And leads to relaxation of smooth muscle in the bladder neck and prostate	Bhp
Omegapress 0.1 MG 0.2 MG		#Desmopressin Synthetic analogue of natural pituitary hormone 8-arginine vasopressin ADH antidiuretic hormone	Urinary incontinence antidiuretic

Ondalenz 8 "ZOFRAN"		Ondansteron Selective serotonin 5 HT3 receptor antagonist	Anti-emetic
Onbrez breezhaler		#Indacaterol **Mechanism** Acts locally in the lung as a bronchodilator. Indacaterol is a partial agonist at the human beta2-adrenergic receptor with Nano molar potency. In isolated human bronchus, indacaterol has a rapid onset of action and a long duration of action	Onbrez Breezhaler is a medicine that is used to keep the airways open in adults with chronic obstructive pulmonary disease (COPD). COPD is a long-term disease in which the airways and air sacs inside the lungs become damaged or blocked, leading to difficulty breathing
Onglyza		#Saxagliptin inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	

Optaminess 100 150 200			Dietary supplement the diet of hemodialysis patient with essential amino acids
Orgoroxaban 10-20 "XATRELTO"		#Rivaroxaban **Mechanism** - Anti platelet -direct factor XA inhibitor -blocks the formation of blood clots **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-reduce the risk of stroke and serious blood clots in certain patients with atrial fibrillation - ttt and prevent DVT which can lead to blood clots in the lung – pulmonary embolism-
Oseltamiver "TAMIFLU"		# Oseltamiver	
Pariet 20-10		Aripeprazol	Ppi
Pentaza Tab Sup		#Mesalazine **Mechanism** Exerts its anti-inflammatory action by inhibiting lipoxy genase enzyme this will lead to decrease the production of pro inflammatory leukotrienes and 5 hydroxy eicosa tetre enoic acid in the intestinal wall -it inhibits the production of inflammatory mediators like interleukin1 –tumor nicrosis factor	-ulcerativ colitis -crohns disease -proctitis b and proctosigmoiditis-used only suppositories=

Piascledine 300		#Avocado oil soya bean oil **Mechanism** Block the bro inflammatory chemicals Prevents the deterionation of synovial cells(which)line the joints) May help regenerate normal connective tissue	joint arthritis
Pixaspire 2.5-5 "ELIQUIS"		#Apixaban **Mechanism** - Anti co agulant -direct factor xa inhibitor **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial fibrillation
Pivastalo 1-2-4		#Pivastatin **Mechanism** Completely inhibits HMG CoA reductase, the rate-determining enzyme in hepatic cholesterol synthesis. Consequently, LDL-C receptors in the liver are increased, thereby increasing the removal of LDL-C from the blood	
Pixclot 2.5 "ELIQUIS"		#Apixaban **Mechanism** - Anti co agulant -direct factor xa inhibitor **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial fibrillation

Pletaal 50-100		#Clostazol **Mechanism** selective inhibitor of phosphodiesterase type 3 With therapeutic focus on increasing cAMP. An increase in cAMP results in an increase in the active form of protein kinase A(PKA), which is directly related with an inhibition in platelet aggregation. PKA also prevents the activation of an enzyme (myosin light-chain kinase) that is important in the contraction of smooth muscle cells, thereby exerting its vasodilatory effect.	Antiplatlet
Plavidamol 75 "PLAVIX"		#clopidogril	used to reduce the risk of myocardial infarction and stroke or established peripheral arterial disease
Plavix 75		#clopidogril	used to reduce the risk of myocardial infarction and stroke or established peripheral arterial disease

Prandaglim 25-6-25		#Alogliptin **Mechanism** inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	Anti-diabetic
Pradaxa 75 110 150		#Dabigatran **Mechanism** Reversibly binds to the active site on the thrombin molecule, preventing thrombin-mediated activation of coagulation factors. 	used to prevent stroke and harmful blood clots (such as in your legs or lunges !) if you have a certain type of irregular heartbeat atrial fibrillation) . Dabigatran is also used to treat blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) and to reduce the risk of them occurring again. This medication may also be used to prevent these blood clots from forming after hip replacement surgery.

Pregnacare			
Procoralan 5-7.5		#Ivabradine **Mechanism** - Anti ischemic -Hart rate lowering agent that works through selective and specific inhibition of cardiac pacemaker if current	Chronic stable angina in coronary artery disease patient with normal sinus rhythm
Proscar 5		#Finasteride **Mechanism** Blocks the action of an enzyme called 5-alpha-reductase. This enzyme changes testosterone to another hormone that causes the prostate to grow It will increase testosterone levels in the body, which decreases prostate size	used to shrink an enlarged prostate (benign prostatic hyperplasia or BPH) in adult men
Prostride 5 "PROSCAR"		#Finasteride **Mechanism** Blocks the action of an enzyme called 5-alpha-reductase. This enzyme changes testosterone to another hormone that causes the prostate to grow It will increase testosterone levels in the body, which decreases prostate size	used to shrink an enlarged prostate (benign prostatic hyperplasia or BPH) in adult men
Quetiazic xr 50- xr150 xr300- xr400 "SEROQUEL"		**Mechanism** #Quetiapine Second-generation antipsychotic that has affinity for D2, 5-HT2A, H1, alpha 1 and 5-HT1A receptors. Its precise mechanism of action is unknown, but according to the dopamine theory of schizophrenia, antipsychotic effects might be related to the drug's ability to reduce dopaminergic neurotransmission in the mesolimbic pathway.	used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder sudden episodes of mania or depression associated with bipolar disorder

Quetaoine 25 100 200 "SEROQUEL"		**Mechanism** #Quetiapine Second-generation antipsychotic that has affinity for D2, 5-HT2A, H1, alpha 1 and 5-HT1A receptors. Its precise mechanism of action is unknown, but according to the dopamine theory of schizophrenia, antipsychotic effects might be related to the drug's ability to reduce dopaminergic neurotransmission in the mesolimbic pathway.	used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder sudden episodes of mania or depression associated with bipolar disorder
Quitapex 25 100 200 300 400 "SEROQUEL"		**Mechanism** #Quetiapine Second-generation antipsychotic that has affinity for D2, 5-HT2A, H1, alpha 1 and 5-HT1A receptors. Its precise mechanism of action is unknown, but according to the dopamine theory of schizophrenia, antipsychotic effects might be related to the drug's ability to reduce dopaminergic neurotransmission in the mesolimbic pathway.	used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder sudden episodes of mania or depression associated with bipolar disorder
Quitcool 50-200-300 "SEROQUEL"		**Mechanism** #Quetiapine Second-generation antipsychotic that has affinity for D2, 5-HT2A, H1, alpha 1 and 5-HT1A receptors. Its precise mechanism of action is unknown, but according to the dopamine theory of schizophrenia, antipsychotic effects might be related to the drug's ability to reduce dopaminergic neurotransmission in the mesolimbic pathway.	used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder sudden episodes of mania or depression associated with bipolar disorder

Ravemantine 10 "EBIXA"		# Memantine **Mechanism** n-methyl –d-aspartate receptor antagonist (NMDA) -dopamine d2 agonist -xss glutamate plays a role in neural cell death	Antidementia agent in Alzheimer disease
Revolade 25-50		eltrombopag	Ttt primary immune thrombocytopenia (ITP), a disease in which the patient's immune system destroys platelets. Patients with ITP have low platelet counts in the blood (thrombocytopenia) and are at risk of bleeding
Resolor 1-2		#Prucalopride **Mechanism** selective height affinity serotonin receptor agonist which targets the impaired motility associated with chronic constipation thus normalize bowel movement -it stimulates colonic mass movement	Ttt chronic constipation
Recoxibright 30-60-90-120 "ARCOXIA"		#Etoricixib **Mechanism** selective co2 inhibitor	-Rheumatoid arthritis -acute pain -gout -chronic low back pain -osteoarthritis -ankylosing spondylitis
Risperdal 2 4 Solution Consta 25 Consta 37.5		#Risperidone **Mechanism** -block dopamine and serotonin receptors -has antagonistic action on alpha adrenergic and histaminic receptor	Antipychotic

Rivarospire 2.5-10-15-20 "XARELTO"		#Rivaroxaban **Mechanism** - Anti platelet -direct factor XA inhibitor -blocks the formation of blood clots **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-Reduce the risk of stroke and serious blood clots in certain patients with atrial fibrillation - ttt and prevent DVT which can lead to blood clots in the lung – pulmonary embolism-
Roaccutan 10-20		Isotretinoin	Anti-acne
Roaccutrex 10-20-40 "ROACCUTAN "		Isotretinoin	Anti-acne
Sandimmune-neoral 25 50 100		**Cyclosporine** Its major mode of action is inhibition of the production of cytokines involved in the regulation of T-cell activation. In particular, cyclosporine inhibits the transcription of interleukin 2.	used to prevent organ rejection after a kidney, heart, or liver transplant. This medicine is also used to treat severe psoriasis or severe rheumatoid arthritis
Seebri beezhaler		#glycopyronium Works by blocking the bronchoconstrictor action of acetylcholine on airway smooth muscle cells thereby dilating the airways. Of the five known muscarinic receptor subtypes (M1-5), only subtypes M1-3 have a defined physiological function in the human lung.	Seebri Breezhaler is indicated as a maintenance bronchodilator treatment to relieve symptoms in adult patients with chronic obstructive pulmonary disease (COPD). Posology The recommended dose is the inhalation of the content of one capsule once daily using the Seebri Breezhaler inhaler

Seretide diskus 100l50 250l50 500l50		salmeterol and fluticasone	asthma and chronic obstructive pulmonary disease (COPD) in people who need regular treatment
Seroquel 25-100-200- Xr 50- Xr 200- Xr 300- Xr 400		**Mechanism** #Quetiapine Second-generation antipsychotic that has affinity for D2, 5-HT2A, H1, alpha 1 and 5-HT1A receptors. Its precise mechanism of action is unknown, but according to the dopamine theory of schizophrenia, antipsychotic effects might be related to the drug's ability to reduce dopaminergic neurotransmission in the mesolimbic pathway.	used to treat certain mental/mood conditions (such as schizophrenia, bipolar disorder sudden episodes of mania or depression associated with bipolar disorder
Singulair 4 5 10		# Montelukast Montelukast is in the leukotriene receptor antagonist family of medications. It works by blocking the action of leukotriene D4 in the lungs resulting in decreased inflammation and relaxation of smooth muscle	By blocking leukotrienes, Singulair improves asthma sym ptoms, helps control asthma and improves seasonal allergy symptoms (also known as hay fever or seasonal allergic rhinitis). Your doctor has prescribed Singulair to treat asthma, preventing your asthma symptom s during the day and night
Sifrol 0.18 0.7		#Pramipexol **Mechanism** presynaptic D2 and D3 dopamine auto receptors and suppresses the synthesis and synaptic release of dopamine	to treat the symptoms of Parkinson's disease

Solian 200-400		#Amisulpride **Mechanism** amisulpride binds selectively to dopamine D2, D3 receptors in the limbic system thereby enhancing dopaminergic transmission	Antispychotics
Spiriva 18 inh Spiriva respimat 2.5		#tiotropium bromide **Mechanism** Anticholinergic (muscarinic antagonist) -bronchodilator	-chronic obstructive pulmonary disease(COPD) -bronchospasm -chronic bronchitis -emphysema
Strattera 10 18 25 40 60		#atomoxetine **Mechanism** Selective inhibition of presynaptic norepinephrine reuptake in the prefrontal cortex. Atomoxetine has a high affinity and selectivity for norepinephrine transporters, but little or no affinity for various neurotransmitter receptors	Ttt of ADHD Attention deficit hyperactivity disorder
Starkoprex 5-20			
Strakopena 2.5-5		#Apixaban **Mechanism** - Anti co agulant -direct factor xa inhibitor	-DVT prophylaxis -DVT may lead to pulmonary embolism – PE-in knee or lip replacement surgery -ttt both DVT and PE -to lower the risk of stroke and embolism in patient with non valvular atrial fibrillation
Stroka75 "PLAVIX"		#clopidogril	used to reduce the risk of myocardial infarction and stroke or established peripheral arterial disease

Symbicort 80l4.5 160l4.5 320l9		#Budesonide **Mechanism** *Budesonide* is an inhaled corticosteroid (ICS) that works by reducing and preventing respiratory tract inflammation, *formoterol* is a long-acting beta2-agonist bronchodilator (LABA) that decreases resistance in the respiratory airway and increases airflow to the lungs	Symbicort is used to control and prevent the symptoms of asthma in adults and children at least 6 years old. Symbicort is also used to help control the symptoms of chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema
Sympaprost8		#Silodosin **Mechanism** Alpha 1 a blocker	Ttt urination difficulty in benign prostatic hyperplasia
Synjardy 12.5l1000 12.5l850 5l1000 5l850		Empagliflozin Metformin **Mechanism** *#Empagliflozin* works by inhibiting the sodium-glucose co-transporter-2 (SGLT-2) found in the proximal tubules in the kidneys. Through SGLT2 inhibition, empagliflozin reduce s renal reabsorption of glucose and increases urinary excretion of glucose *Metformin* in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Anti diabetic
Tareg 40 80 160 320		#valsartan	Anti-hyper tension
Tecavir 0.5-1		Entecavir **Mechanism** Entecavir is a nucleoside analog, or more specifically, a deoxyguanosine analogue that belongs to a class of carbocyclic nucleosides and inhibits reverse transcription, DNA replication and transcription in the viral replication process	Ttt chronic hepatitis b virus

Tenaviron 300		#tenafovir disoproxil **<u>Mechanism</u>** <u>*tenofovir*</u> interferes with an enzyme called reverse transcriptase, which is used by HIV-infected cells to make new viruses. Since tenofovir inhibits, or reduces, the activity of this enzyme, this drug causes HIV-infected cells to slow down or stop producing new viruses	Tenofovir is used with other HIV medications to help control HIV infection. It helps to decrease the amount of HIV in your body so your immune system can work better. This lowers your chance of getting HIV complicati ons (such as new infections, cancer) and improves your quality of life
Thrombonorm		#Anagrelide **<u>Mechanism</u>** <u>*Anagrelide*</u> works by inhibiting the maturation of platelets from megakaryocytes. The exact mechanism of action is unclear, although it is known to be a phosphodiesterase inhibitor. It is a potent (IC_{50} = 36nM) inhibitor of phosphodiesterase-II. It inhibits PDE-3 and phospholipase A2 **Anagrelide** is in a **class** of medications called platelet-reducing agents. It works by slowing the production of platelets in the body Anagrelide is not a chemotherapy drug; however, some hospitals classify anagrelide as a form of chemotherapy because it interferes with cell development, and the prescribing regulations are similar to those governing a chemotherapy drug	**Anagrelide** is **used** to treat a certain blood disorder (thrombocythemia), which is caused by your bone marrow making too many platelets. Platelets are a blood cell that the body **uses** to form blood clots. Too many platelets can cause problems with your circulation, including unwanted blood clots and bleeding problems

Thrombolinta 90 "BRILIQUE"		#Ticagrelor **Mechanism** Platelet aggregation inhibitor Works by slowing or stopping platelet from sticking to blood vessel walls or injured tissues	-reducing the risk of stroke heart attack -or death in certain patient who have had a heart attack or who have angina (chest pain) -it should be used along with asprin as directed by your doctor
Topamax 15 25 50 100		#Topiramate blocks voltage-dependent sodium and calcium channels. It also inhibits the excitatory glutamate pathway while enhancing the inhibitory effect of GABA. Moreover, it inhibits carbonic anhydrase activity.	used to treat epilepsy. These are also called anti- epileptic medicines. It can also be taken to prevent migraine
Trileptal 600-300		#oxcarbazepine is a sodium channel blocker. It binds preferentially to voltage-gated sodium channels in their inactive conformation, which prevents repetitive and sustained firing of an **action** potential	Anticonvulsants to control certain types of seizures in adults a nd children 6 years of age and older.
Trio clar		Tinidazole Clarithromycin Omeprazole	Ttt h pylori
Tiratam 500 500xr 750 1000 Vial Oral suspension		#Levetiracetam **Mechanism** Binds to synaptic vesicle protein SV2A which is believed to impede nerve conduction across synapses	Ttt epilepsy anti convulsant neuropathic pain anxiety disorder

Trajenta 5		#linagliptin inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	treat type 2 diabetes
Tiralepsy 250-500-1000 Xr 500-750- oralsolution		#Levetiracetam **Mechanism** Binds to synaptic vesicle protein SV2A which is believed to impede nerve conduction across synapses	Ttt epilepsy anti convulsant neuropathic pain anxiety disorder
Twynsta 80l5 80l10		#Telmisatran Amlodipine	Anti-hyper tensive

ultibro		# indacaterol **Mechanism** a long-acting beta-2 agonist producing bronchodilation by stimulation of intracellular adenyl cyclase. glycopyrronium, a long-acting muscarinic antagonist/anticholinergic that provides bronchodilation by blocking the action of acetylcholine on the muscarinic receptors of airway smooth muscle cells	Indacaterol inhalation is used to control wheezing, shortness of breath, coughing, and chest tightness caused by chronic obstructive pulmonary disease (COPD; a group of diseases that affect the lungs and airways, which includes chronic bronchitis and emphysema)
Valtrex 500		# Valacyclovir **Mechanism** Is converted to acyclovir, which is converted to its triphosphate form, acyclovir triphosphate (ACV-TP). ACV-TP competitively inhibits viral DNA polymerase, incorporates into and terminates the growing viral DNA chain, and inactivates of the viral DNA polymerase	i s used to treat herpes virus infections, including herpes labial is (also known as cold sores), herpes zoster (also known as shingles), and herpes simplex (also known as genital herpes) in adults. It is also used to treat chickenpox and cold sores in children
Vaxato 10-15-20		#Rivaroxaban **Mechanism** - Anti platelet -direct factor XA inhibitor -blocks the formation of blood clots **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-reduce the risk of stroke and serious blood clots in certain patients with atrial fibrillation - ttt and prevent DVT which can lead to blood clots in the lung – pulmonary embolism

Vesicare 10-5		# Solifenacin **Mechanism** is a competitive muscarinic receptor antagonist. Muscarinic receptors play an important role in several major cholinergically mediated functions, including contractions of urinary bladder smooth muscle and stimulation of salivary secretion	is used to treat overactive bladder (a condition in which the bladder muscles contract uncontrollably and cause frequent urination, urgent need to urinate, and inability to control urination)
Vfend 200		Voriconazol	Ttt of esophageal candidiasis invasive pulmonary aspergillosis -ttt serious fungal infections caused by scedosporium apiospermum and fusarium spp
Visanne 2		#Dienogest **Mechanism** Dienogest belongs to the class of medications called *progestins*, medications that act like progesterone in the body Dienogest acts on endometriosis by reducing the endogenous production of estradiol and thereby suppressing the trophic effects of estradiol on both the eutopic and ectopic endometrium	- contraceptive -Progestins reduce the effects of estrogen on tissues such as the *endometrium* (lining of the uterus) and the breast. By reducing the growth effect of estrogen on the endometrium, **dienogest helps to reduce the pelvic pain experienced by women with endometriosis.** Endometriosis is a condition where endometrium-like tissue is found outside the uterus, which causes chronic inflammation.

Vildagluse		#Vildagliptin **<u>Mechanism</u>** inhibits the activity of DPP-4, an enzyme that inactivates and degrades incretin hormones, cytokines and other peptides. Released in response to meals, incretin hormones potentiate insulin release and decrease glucagon production, lowering serum glucose concentrations. The enzyme dipeptidyl peptidase-4 (DPP-4) begins degrading the GLP-1 and GIP incretins within minutes of their release. Consequently, GLP-1 and GIP are short-lived, disrupting glucose homeostasis.[1] , thereby prolonging the body's own action of naturally secreted incretins GLP-1 and GIP.[1] GLP-1 signals to the pancreas to decrease glucagon secretion by pancreatic alpha cells, which decreases hepatic glucose productions.[1] GLP-1 and GIP increase insulin release by signaling the pancreatic beta cells, which increase uptake of glucose by the muscle, fat, and peripheral tissues.	Anti-diabetic
Vildagluse plus 50l1000		Vildagliptin Metformin **<u>Mechanism</u>** #*<u>Vildagliptin</u>* binds covalently to the catalytic site of DPP-4, eliciting prolonged enzyme inhibition. This raises intact GLP-1 levels, both after meal ingestion and in the fasting state. Vildagliptin has been shown to stimulate insulin secretion and inhibit glucagon secretion in a glucose-dependent manner *<u>#Metformin</u>* in patients with type 2 diabetes mellitus setting is inhibition of hepatic gluconeogenesis.	Anti diabetic

Xarelto 2.5-10-20-15		#Rivaroxaban **Mechanism** - Anti platelet -direct factor XA inhibitor -blocks the formation of blood clots **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin 	-reduce the risk of stroke and serious blood clots in certain patients with atrial fibrillation - ttt and prevent DVT which can lead to blood clots in the lung – pulmonary embolism-
Xanoxiban 2.5-10-15-20		#Rivaroxaban **Mechanism** - Anti platelet -direct factor XA inhibitor -blocks the formation of blood clots **Factor Xa** is generated by both the extrinsic and intrinsic coagulation pathways and is responsible for activating prothrombin to thrombin	-reduce the risk of stroke and serious blood clots in certain patients with atrial fibrillation - ttt and prevent DVT which can lead to blood clots in the lung – pulmonary embolism-
Xeloda		# Capecitabine **Mechanism** is relatively non-cytotoxic in vitro. This drug is enzymatically converted to 5-fluorouracil (5-FU) in vivo. Mechanism of Action: Both normal and tumor cells metabolize 5-FU to 5-fluoro-2-deoxyuridine monophosphate (FdUMP) and 5-fluorouridine triphosphate (FUTP) -Xeloda belongs to a class of drugs called Antineoplastics, Antimetabolite.	is a prescription medicine **used** to treat the symptoms of cancers such as Colon Cancer, Colorectal Cancer and Breast Cancer. Xeloda may be used alone or with other medications.

Xenical 120		# orlistat **Mechanism** Acts by reversibly inhibiting the gastric and pancreatic lipases. These lipases have an important role in the digestion of dietary fat. They work by breaking down the triglycerides into absorbable free fatty acids and monoglycerides	It is used by certain overweight people, such as those who are obese or have weight-related medical problems. Taking orlistat can also help keep you from gaining back weight you have lost
Zeffix100		#Lamivudine **Mechanism** is an analogue of cytidine. It can inhibit both types (1 and 2) of HIV reverse transcriptase and also the reverse transcriptase of hepatitis B virus. It is phosphorylated to active metabolites that compete for incorporation into viral DNA	Used to treat hepatitis B infection. Lamivudine is in a class of medications called nucleoside reverse transcriptase inhibitors (NRTIs). It works by decreasing the amount of HIV and hepatitis B in the blood
Zofran 8		Ondansteron Selective serotonin 5 HT3 receptor antagonist	Anti-emetic
Zyprexa 10 velotab 10-5-7.5	 	Olazapine The xact mechanism of action of olanzapine is not known. It may work by blocking receptors for several neurotransmitters (chemicals that nerves use to communicate with each other) in the brain. It binds to alpha-1, dopamine, histamine H-1, muscarinic, and serotonin type 2 (5-HT2) receptors	For acute and maintenance ttt of schizophrenia