National University of Computer and Emerging Sciences, Lahore Campus

	and the state of t
П	TIONAL UNIVERSE
	13
	STATE OF THE PARTY WITH
	THE THE PARTY OF T
	1/3/3/87 4 111
	A CONTRACTOR OF THE PARTY OF TH

	Numerical Computing	Course Code:	CS2008	
Course Name:		Semester:	Fall 2021	
Degree Program:	BS (CS)	Total Marks:	65	
Exam Duration:	60 Minutes	Weight	15%	
Paper Date:	December 3; 2021	Page(s):		
Section:	ALL	1 490(0).		*******
Exam Type:	Mid Term-II			_

Student: Name:

Instruction/Notes:

Attempt all questions on the answer book. Programmable calculators are not allowed. Don't write anything on a question paper except your name and roll number.

Points (10)

For what values h and N composite Trapezoidal approximation of the following integral is correct to 10^{-4} .

$$\int_{1.5}^{3.5} e^x dx$$

Points (10)

Use composite Simpson rule with h = 1 and 0.5 to approximate $\int_{-1}^{1} xe^{-3x} dx$ having exact solution 4.485580235538532. Then use Romberg Integration based on Simpson rule to find $O(h^6)$ approximation. Compare your results with exact solution and make conclusion.

Note: Throughout the problem values should be taken at least 6 decimals places.

Points (5)

Find the minimum number of iterations N, needed to achieve an accuracy of 10^{-3} by using the bisection

 $f(x) = e^x - 1 - x - \frac{x^2}{2}$ in [-1, 1]

Note: Find N without solving f(x) = 0 for x.

(B)

Points (10)

Show that order of convergence of Newton - Raphson method is 2.

Points (10)

Show that Newton sequence of iterations for the function $f(x) = x^3 - 2x + 2$ with $x_0 = 0$ will oscillate between 0 and 1. Furthermore, find the real root of f(x) = 0 up to 5 decimal places using Newton's method by choosing suitable x_0 .

Q3. (A)

Points (10)

Solve the following linear system using Crout's Method.

$$\begin{bmatrix} 3 & 1 & -1 \\ -1 & 2 & 8 \\ 1 & -5 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 8 \\ -8 \\ -4 \end{bmatrix}$$

For the system given in Q3 (A), find approximate solution by performing first iteration using Gauss **(B)** Jacobi method and then second iteration by Gauss Seidal method. Use [1,1,1] as initial guess for this Points (10) problem.