

Gradient Descent

Linear Regression Single Variable


Training Set	Size in feet ² (x)	Price (\$) in 1000's (y)
	2104	460
	1416	232
	1534	315
	852	178

Hypothesis: $h_{\theta}(x) = \theta_0 + \theta_1 x$

θ_i 's: Parameters


How to choose θ_i 's ?

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$




$$\theta_0 = 1.5$$

$$\theta_1 = 0$$




$$\theta_0 = 0$$

$$\theta_1 = 0.5$$



$$\theta_0 = 1$$

$$\theta_1 = 0.5$$



Idea: Choose θ_0, θ_1 so that $h_{\theta}(x)$ is close to y for our training examples (x, y)

Simplified

Hypothesis:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Parameters:

$$\theta_0, \theta_1$$

Cost Function:

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Goal: minimize $J(\theta_0, \theta_1)$
 θ_0, θ_1

$$h_{\theta}(x) = \theta_1 x$$


$$\theta_1$$

$$J(\theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

minimize $J(\theta_1)$
 θ_1

$$h_{\theta}(x)$$


(for fixed θ_1 , this is a function of x)



$$h_{\theta}(x) = \theta_1 x$$

$$J(\theta_1)$$


(function of the parameter θ_1)



$$J(\theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

$$h_{\theta}(x)$$


(for fixed θ_1 , this is a function of x)



$$h_{\theta}(x) = \theta_1 x$$

$$J(\theta_1)$$


(function of the parameter θ_1)



$$J(\theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

$$h_{\theta}(x)$$


(for fixed θ_1 , this is a function of x)



$$h_{\theta}(x) = \theta_1 x$$

$$J(\theta_1)$$

(function of the parameter θ_1)



$$J(\theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Hypothesis: $h_{\theta}(x) = \theta_0 + \theta_1 x$

Parameters: θ_0, θ_1

Cost Function: $J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$


Goal: minimize $J(\theta_0, \theta_1)$
 θ_0, θ_1

$$h_{\theta}(x)$$


(for fixed θ_0, θ_1 , this is a function of x)

$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)



$$h_{\theta}(x) = 50 + 0.06x$$





Have some function $J(\theta_0, \theta_1)$

Want $\min_{\theta_0, \theta_1} J(\theta_0, \theta_1)$

Outline:

- Start with some θ_0, θ_1
- Keep changing θ_0, θ_1 to reduce $J(\theta_0, \theta_1)$
until we hopefully end up at a minimum





Gradient descent algorithm

repeat until convergence {
 $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$ (for $j = 0$ and $j = 1$)
}

Correct: Simultaneous update

```
temp0 :=  $\theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$   
temp1 :=  $\theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$   
 $\theta_0 :=$  temp0  
 $\theta_1 :=$  temp1
```


Incorrect:


```
temp0 :=  $\theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$   
 $\theta_0 :=$  temp0  
temp1 :=  $\theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$   
 $\theta_1 :=$  temp1
```

$$\theta_1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_1)$$

If α is too small, gradient descent can be slow.

If α is too large, gradient descent can overshoot the minimum. It may fail to converge, or even diverge.






Current value of θ_1

$$\theta_1 := \theta_1 - \alpha \frac{d}{d\theta_1} J(\theta_1)$$

Gradient descent can converge to a local minimum, even with the learning rate α fixed.

$$\theta_1 := \theta_1 - \alpha \frac{d}{d\theta_1} J(\theta_1)$$

As we approach a local minimum, gradient descent will automatically take smaller steps. So, no need to decrease α over time.



$$\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) =$$

$$j = 0 : \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) =$$

$$j = 1 : \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) =$$

Gradient descent algorithm


repeat until convergence {


$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$


$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x^{(i)}$$

}

update
 θ_0 and θ_1
simultaneously









$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)




$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)




$$h_{\theta}(x)$$

(for fixed θ_0, θ_1 , this is a function of x)



$$J(\theta_0, \theta_1)$$

(function of the parameters θ_0, θ_1)



Reference

Slides adapted from
Machine Learning By Andrew Ng