

Unit 19

NITROGEN COMPOUNDS

17 items · 22 marks

Syllabus sections in this unit


19.1 Primary amines

19.2 Nitriles and hydroxynitriles

Unit 19: Nitrogen compounds**1. M/J 26/P24/Q4(d)(ii)**

Context: part (d) (introduction)

(d) Butane is converted to pentanenitrile via a two-step synthesis, as shown in Figure 4.3.

**Figure 4.3**

(ii) Pentanenitrile reacts with dilute acid.

Draw the displayed formula of the organic product made in this reaction.

[1]

2. MAR 26/P22/Q4(a)(iv)

Context

4 Cyanoethanoic acid, $\text{CH}_2(\text{CN})\text{COOH}$, is a weak organic acid.

Context: part (a) (introduction)

(a) At room temperature, $\text{CH}_2(\text{CN})\text{COOH}$ undergoes an acid–base reaction with $\text{NaOH}(\text{aq})$.

(iv) Suggest the formula of the organic product when $\text{CH}_2(\text{CN})\text{COOH}(\text{aq})$ is heated under reflux with $\text{NaOH}(\text{aq})$.

[1]

3. O/N 25/P24/Q4(b)(ii)


Context

- 4** Compounds **P** and **Q** are structural isomers.

**Fig. 4.1**

Context: part (a)(v)

- (v)** **Q** reacts with HBr(g) to produce two structural isomers, **R** and **S**, as shown in Fig. 4.2.


**Fig. 4.2**

State and explain why isomer **S** is the major product of the reaction.

[2]


Context: part (b) (introduction)

- (b)** Fig. 4.3 shows a synthesis starting from **T**, a different isomer of **R** and **S**.

**Fig. 4.3**

- (ii)** Reaction 2 is a hydrolysis reaction.

Construct an equation for reaction 2.



[1]

4. M/J 25/P22/Q4(b)(iii)-(iv)

Context: part (b) (introduction)

- (b)** Propan-1-ol is converted to compound **T** in a three-step synthesis.

In step 1, SOCl_2 is added to propan-1-ol to produce compound **E**.

In step 2, **E** reacts with a suitable reagent to produce butanenitrile.

In step 3, butanenitrile is heated with NaOH(aq) .

- (iii)** Construct an equation to describe the reaction of butanenitrile with NaOH(aq) in step 3.

[2]


- (iv)** Name the type of reaction that occurs in step 3.

[1]


5. M/J 25/P23/Q6(c)(ii)-(iii)

Context: part (c) (introduction)

- (c) $\text{HOCH}_2\text{CH}_2\text{Cl}$ reacts in a two-step synthesis to produce $\text{HOCH}_2\text{CH}_2\text{COOH}$, as shown in Fig. 6.2.

**Fig. 6.2**

- (ii) Identify the type of reaction that occurs in step 2.
- (iii) Complete the equation to show the reaction in step 2.




[1]

[1]

6. M/J 25/P24/Q4(d)(i)

Context

- 4 The skeletal formula of **E** is shown in Fig. 4.1.

**Fig. 4.1**

Context: part (d) (introduction)

- (d) **F** is an isomer of **E**.

F reacts with HCl(aq) to produce **G** and **H**.

G is a secondary alcohol.

H is also produced when HCl(aq) is added to CH_3CN .


- (i) Construct an equation for the acid hydrolysis of CH_3CN with HCl(aq) .

[1]

7. MAR 25/P22/Q4(b)(ii)

Context

- 4 Fig. 4.1 shows compounds **J** to **M**, each of which contains four carbon atoms.

**Fig. 4.1**

Context: part (b) (introduction)

- (b) **K** reacts with HCN in the presence of a KCN catalyst, forming **N**.


- (ii) Identify the class of compound of which **N** is a member.

[1]

8. O/N 24/P21/Q4(b)

Context

- 4** Propanone, CH_3COCH_3 , is an important organic reagent. Fig. 4.1 shows some reactions of propanone and its derivatives.

**Fig. 4.1**

- (b)** Suggest the reagents and conditions for reaction 2.

[1]

9. O/N 24/P22/Q2(b)(ii)

Context

- 2** Oxygen is a Group 16 element.

Context: part (b) (introduction)

- (b)** H_2O reacts with both inorganic and organic compounds.

- (ii)** Write an equation for the reaction of CH_3CN with H_2O in acidic conditions.




[1]

10. O/N 24/P23/Q4(b)

Context

- 4** Propanone, CH_3COCH_3 , is an important organic reagent. Fig. 4.1 shows some reactions of propanone and its derivatives.

**Fig. 4.1**


- (b)** Suggest the reagents and conditions for reaction 2.

[1]

11. M/J 24/P23/Q5(d)(i)-(ii)

Context: part (a)

- 5** **(a)** **M** reacts to form **R** by the addition of one reagent, as shown in Fig. 5.1.


**Fig. 5.1**

Identify the reagent and conditions for this reaction.

[1]

Context: part (b) (introduction)

- (b)** **R** is also made from **M** by two steps, as shown in Fig. 5.2.

**Fig. 5.2**

Context: part (d) (introduction)

- (d)** **Y** is made from **Q** in a three-step reaction.




Fig. 5.4

- (i) Draw the structure of **W** in the box in Fig. 5.4. [1]
- (ii) In step 2, **W** is heated with HCl(aq) to produce **X** and an inorganic product. Identify the formula of the inorganic product. [1]

12. MAR 24/P22/Q4(b)(iii)

Context

- 4 Compound **S** is used in food flavourings. A possible synthesis of **S** is shown in Fig. 4.1.





Fig. 4.1

- (iii) Complete the equation for reaction 3. **R** is represented as $\text{C}_4\text{H}_9\text{CN}$.




[1]

13. M/J 23/P22/Q3(a)(iii)

Context

- 3** Fig. 3.1 describes a sequence of reactions that can be used to produce a food additive, compound **Y**, from $\text{CH}_3\text{CH}_2\text{Cl}$.

**Fig. 3.1**

- (iii) Identify the type of reaction that occurs when dilute acid is added to **X** in step 2.

[1]

14. M/J 23/P23/Q6(e)

Context

- 6** Compound **W**, $\text{CH}_2=\text{CHCN}$, is used to make an addition polymer which is present in carbon fibres.

- (e) Propanenitrile is heated with hydrogen gas and a platinum catalyst. The only product is propylamine.

Construct an equation for this reaction.

[1]


15. M/J 23/P23/Q6(f)(ii)

Context

- 6** Compound **W**, $\text{CH}_2=\text{CHCN}$, is used to make an addition polymer which is present in carbon fibres.


Context: part (b) (introduction)

- (b) CH_3CHO is used in a two-step synthetic route to form **W**, as shown in Fig. 6.1. In step 1, CH_3CHO is heated with HCN in the presence of KCN .

**Fig. 6.1**

Context: part (f) (introduction)

- (f) Propylamine can also be formed in a two-step synthesis from propan-1-ol, as shown in Fig. 6.4.

**Fig. 6.4**

- (ii) Identify the reagent and conditions for step 2 in Fig. 6.4.

[2]

16. MAR 23/P22/Q4(c)(ii)

Context

4 Fig. 4.1 shows some reactions of compound **D**, 2-bromobutane.




Fig. 4.1

(ii) Reaction 5 is similar to the reaction of LiAlH_4 with carboxylic acids to form alcohols.


Suggest the role of LiAlH_4 in reaction 5.

[1]

17. M/J 22/P22/Q5(c)(ii)


Context

5 Fig. 5.1 shows three reactions of 2-bromopropane, $\text{CH}_3\text{CH}(\text{Br})\text{CH}_3$.

**Fig. 5.1**

Context: part (c) (introduction)

(c) Fig. 5.2 shows how butan-1-ol can be made from 1-bromopropane in three steps.

**Fig. 5.2**

(ii) In step 2, butanenitrile is heated with $\text{HCl}(\text{aq})$. A hydrolysis reaction occurs.

Construct an equation for the reaction in step 2.

[1]

Answers Section

1. M/J 26/P24/Q4(d)(ii)

4(d)(ii)	$ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & & \text{H} & & \text{O} \\ & & & & & & & & & // \\ \text{H} & - \text{C} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{C} \\ & & & & & & & & & \backslash \\ & \text{H} & & \text{H} & & \text{H} & & \text{H} & & \text{O} - \text{H} \end{array} $	1	R-OH
----------	---	---	------

2. MAR 26/P22/Q4(a)(iv)

4(a)(iv)	CH ₂ (COO) ₂ Na ₂ OR CH ₂ (COONa) ₂	1
----------	---	---

3. O/N 25/P24/Q4(b)(ii)

4(b)(ii)	(C ₂ H ₅) ₂ CHCN + 2H ₂ O + HCl → (C ₂ H ₅) ₂ CHCOOH + NH ₄ Cl	1
----------	--	---

Answer for context part (a)(v)

4(a)(v)	M1 (reaction proceeds via a) more stable intermediate / carbocation M2 because of greater positive inductive effect of more alkyl groups	2
---------	---	---

4. M/J 25/P22/Q4(b)(iii)-(iv)

4(b)(iii)	CH ₃ CH ₂ CH ₂ CN + NaOH + H ₂ O → CH ₃ CH ₂ CH ₂ CO ₂ Na + NH ₃ OR CH ₃ CH ₂ CH ₂ CN + NaOH + 2H ₂ O → CH ₃ CH ₂ CH ₂ CO ₂ Na + NH ₄ OH M1 correct organic product (as molecule or anion) M2 rest of equation is correct	2
4(b)(iv)	(alkaline) hydrolysis	1

5. M/J 25/P23/Q6(c)(ii)-(iii)

6(c)(ii)	(acid) hydrolysis	1
6(c)(iii)	HOCH ₂ CH ₂ CN + 2H ₂ O + HCl → HOCH ₂ CH ₂ COOH + NH ₄ Cl	1

6. M/J 25/P24/Q4(d)(i)

4(d)(i)	CH ₃ CN + HCl + 2H ₂ O → CH ₃ COOH + NH ₄ Cl	1
---------	--	---

7. MAR 25/P22/Q4(b)(ii)

4(b)(ii)	hydroxynitrile	1
----------	----------------	---

8. O/N 24/P21/Q4(b)

4(b)	H ₂ SO ₄ (aq) or HCl(aq)	1
------	--	---

9. O/N 24/P22/Q2(b)(ii)

2(b)(ii)	CH ₃ CN + 2H ₂ O + (1)H ⁺ → CH ₃ COOH + NH ₄ ⁺	1
----------	--	---

10. O/N 24/P23/Q4(b)

4(b)	H ₂ SO ₄ (aq) or HCl(aq)	1
------	--	---

11. M/J 24/P23/Q5(d)(i)-(ii)

5(d)(i)	W =	1
5(d)(ii)	NH_4Cl	1

Answer for context part (a)

5(a)	cold dilute KMnO_4 OR cold dilute potassium manganate(VII)	1
------	--	----------

12. MAR 24/P22/Q4(b)(iii)

4(b)(iii)	$\text{C}_4\text{H}_9\text{CN} + 2\text{H}_2\text{O} + \text{H}^+ \rightarrow \text{C}_4\text{H}_9\text{COOH} + \text{NH}_4^+$	1
-----------	--	----------

13. M/J 23/P22/Q3(a)(iii)

3(a)(iii)	(acid) hydrolysis	1
-----------	-------------------	----------

14. M/J 23/P23/Q6(e)

6(e)	$\text{CH}_3\text{CH}_2\text{CN} + 2\text{H}_2 \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$	1
------	--	----------


15. M/J 23/P23/Q6(f)(ii)

6(f)(ii)	M1 ammonia / NH_3 M2 heat + (M1 in ethanol) under pressure	2
----------	--	----------

16. MAR 23/P22/Q4(c)(ii)

4(c)(ii)	reducing agent	1
----------	----------------	----------

17. M/J 22/P22/Q5(c)(ii)

5(c)(ii)	$\text{CN}(\text{CH}_2)_2\text{CH}_3 + 2\text{H}_2\text{O} + \text{HCl} \rightarrow \text{HOOC}(\text{CH}_2)_2\text{CH}_3 + \text{NH}_4\text{Cl}$ OR 	1
----------	---	----------