

Matrices

4

□ Matrix

- ▣ Array of numerical values, e.g.:

$$\mathbf{A} = \begin{bmatrix} -7 & 0 & 1 & 4 \\ 4 & -2 & 9 & 5 \\ 8 & 3 & 4 & 0 \end{bmatrix}$$

- ▣ The variable, \mathbf{A} , is a ***matrix***
- An $m \times n$ matrix has m ***rows*** and n ***columns***
- These are the ***dimensions*** of the matrix
 - ▣ \mathbf{A} is a 3×4 matrix

Matrix Dimensions and Indexing

5

- An $m \times n$ matrix:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Use indices to refer to individual elements of a matrix
 - ▣ a_{ij} : the element of \mathbf{A} in the i^{th} row and the j^{th} column

Vectors

6

□ Vectors

- A matrix with one dimension equal to one
- A matrix with ***one row*** or ***one column***

□ ***Row vector***

- One row – a $1 \times n$ matrix, e.g.:

$$x = [-9 \quad 1 \quad -4]$$

- A 1×3 row vector

□ ***Column vector***

- One column – an $m \times 1$ matrix, e.g.:

$$x = \begin{bmatrix} 5 \\ 1 \\ 8 \end{bmatrix}$$

- A 3×1 column vector

Scalars

7

□ Scalar

- A 1×1 matrix

- The numbers we are familiar with, e.g.:

$$b = 4, \quad x = -3 + j5.8, \quad y = -1 \times 10^{-9}$$

- We understand simple mathematical operations involving scalars

- Can add, subtract, multiply, or divide any pair of scalars

- Not true for matrices

- Depends on the matrix dimensions

8

Mathematical Matrix Operations

Matrix Addition and Subtraction

9

- As long as matrices have the ***same dimensions***, we can add or subtract them
 - ▣ ***Addition*** and ***subtraction*** are done ***element-by-element***, and the ***resulting matrix is the same size***

$$\begin{bmatrix} 4 & 8 \\ 0 & 3 \end{bmatrix} + \begin{bmatrix} 1 & -4 \\ 6 & -1 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 6 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 8 \\ 0 & 3 \end{bmatrix} - \begin{bmatrix} 1 & -4 \\ 6 & -1 \end{bmatrix} = \begin{bmatrix} 3 & 12 \\ -6 & 4 \end{bmatrix}$$

- We can also add ***scalars*** to (or subtract from) matrices

$$\begin{bmatrix} 1 & -4 \\ 6 & -1 \end{bmatrix} + 5 = \begin{bmatrix} 6 & 1 \\ 11 & 4 \end{bmatrix}$$

Matrix Addition and Subtraction

10

- If matrices are not the same size, and neither is a scalar, addition/subtraction are not defined
 - ▣ The following operations cannot be done

$$\begin{bmatrix} 4 & 8 \\ 0 & 3 \end{bmatrix} + \begin{bmatrix} 1 & -4 & 6 \\ 6 & -1 & 9 \end{bmatrix} = ?$$

$$\begin{bmatrix} 8 \\ 3 \end{bmatrix} - \begin{bmatrix} 1 & -4 \\ 6 & -1 \end{bmatrix} = ?$$

- Addition is commutative (order does not matter):

$$\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A} = \mathbf{C}$$

$$\begin{bmatrix} 4 & 8 \\ 0 & 3 \end{bmatrix} + \begin{bmatrix} 1 & -4 \\ 6 & -1 \end{bmatrix} = \begin{bmatrix} 1 & -4 \\ 6 & -1 \end{bmatrix} + \begin{bmatrix} 4 & 8 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 6 & 2 \end{bmatrix}$$

Matrix Multiplication

11

- In order to multiply matrices, their *inner dimensions* must agree
- We can multiply $\mathbf{A} \cdot \mathbf{B}$ only if the *number of columns* of \mathbf{A} is equal to the *number of rows* of \mathbf{B}
- Resulting Matrix has same number of rows as \mathbf{A} and same number of columns as \mathbf{B}

$$\begin{array}{c} \mathbf{A} \cdot \mathbf{B} = \mathbf{C} \\ \nearrow \quad \nearrow \quad \nearrow \\ (m \times n) \cdot (n \times p) = (m \times p) \end{array}$$

Matrix Multiplication – $\mathbf{A} \cdot \mathbf{B} = \mathbf{C}$

12

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & \cdots & b_{1p} \\ \vdots & \ddots & \vdots \\ b_{n1} & \cdots & b_{np} \end{bmatrix} = \begin{bmatrix} c_{11} & \cdots & c_{1p} \\ \vdots & \ddots & \vdots \\ c_{m1} & \cdots & c_{mp} \end{bmatrix}$$

- The (i, j^{th}) entry of \mathbf{C} is the **dot product** of the i^{th} row of \mathbf{A} with the j^{th} column of \mathbf{B}

$$c_{ij} = \sum_{k=1}^n a_{ik} \cdot b_{kj}$$

- Consider the multiplication of two 2×2 matrices:

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

Matrix Multiplication – Examples

13

- A 2×2 and a 2×3 yield a 2×3

$$\begin{bmatrix} 1 & 4 \\ 2 & 1 \end{bmatrix} \cdot \begin{bmatrix} 3 & -1 & 5 \\ 6 & 2 & 0 \end{bmatrix} = \begin{bmatrix} 27 & 7 & 5 \\ 12 & 0 & 10 \end{bmatrix}$$

- A 3×3 and a 3×1 result in a 3×1

$$\begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \\ 2 & 7 & 3 \end{bmatrix} \cdot \begin{bmatrix} 6 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 11 \\ 20 \\ 25 \end{bmatrix}$$

Matrix Multiplication – Properties

14

- ***Matrix multiplication is not commutative***

- Order matters
- Unlike scalars

- In general,

$$\mathbf{A} \cdot \mathbf{B} \neq \mathbf{B} \cdot \mathbf{A}$$

- If A and/or B is not square then one of the above operations may not be possible anyway
 - Inner dimensions may not agree for both product orders

Matrix Multiplication – Properties

15

□ ***Matrix multiplication is associative***

- ▣ Insertion of parentheses anywhere within a product of multiple terms does not affect the result:

$$(\mathbf{A} \cdot \mathbf{B}) \cdot \mathbf{C} = \mathbf{D}$$

$$\mathbf{A} \cdot (\mathbf{B} \cdot \mathbf{C}) = \mathbf{D}$$

□ ***Matrix multiplication is distributive***

- ▣ Multiplication distributes over addition
- ▣ Must maintain correct order, i.e. left- or right-multiplication

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$$

$$(\mathbf{B} + \mathbf{C})\mathbf{A} = \mathbf{BA} + \mathbf{CA}$$

Identity Matrix

16

- Multiplication of a scalar by 1 results in that scalar

$$a \cdot 1 = 1 \cdot a = a$$

- The matrix version of 1 is the ***identity matrix***
 - ▣ Ones along the diagonal, zeros everywhere else
 - ▣ Square ($n \times n$) matrix
 - ▣ Denoted as **I** or **I_n**, where **n** is the matrix dimension, e.g.

$$\mathbf{I}_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Left- or right-multiplication by an identity matrix results in that matrix, unchanged

$$\mathbf{A} \cdot \mathbf{I} = \mathbf{I} \cdot \mathbf{A} = \mathbf{A}$$

Identity Matrix

17

- Right-multiplication of an $n \times n$ matrix by an $n \times n$ identity matrix, \mathbf{I}_n

$$\begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \\ 2 & 7 & 3 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \\ 2 & 7 & 3 \end{bmatrix}$$

- Same result if we left-multiply by \mathbf{I}_n

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \\ 2 & 7 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \\ 2 & 7 & 3 \end{bmatrix}$$

Identity Matrix

18

- Right-multiplication of an $m \times n$ matrix by an $n \times n$ identity matrix

$$\begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \end{bmatrix}$$

- Same result if we left-multiply the $m \times n$ matrix by an $m \times m$ identity matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \end{bmatrix} = \begin{bmatrix} 1 & 5 & 0 \\ 0 & 4 & 8 \end{bmatrix}$$

Vector Multiplication

19

- Vectors *are* matrices, so inner dimensions must agree
- Two types of vector multiplication:
- ***Inner product (dot product)***
 - ▣ Result is a scalar

$$\begin{bmatrix} a_{11} & a_{12} \end{bmatrix} \cdot \begin{bmatrix} b_{11} \\ b_{21} \end{bmatrix} = a_{11}b_{11} + a_{12}b_{21}$$

- ***Outer product***
 - ▣ Result for n-vectors is an n x n matrix

$$\begin{bmatrix} a_{11} \\ a_{21} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} & a_{11}b_{12} \\ a_{21}b_{11} & a_{21}b_{12} \end{bmatrix}$$

Exponentiation

20

- As with scalars, raising a matrix to the power, n , is the multiplication of that matrix by itself n times

$$\mathbf{A}^3 = \mathbf{A} \cdot \mathbf{A} \cdot \mathbf{A}$$

- What must be true of a matrix for exponentiation to be allowable?
 - ▣ Inner matrix dimensions must agree
 - ▣ Rows of \mathbf{A} must equal columns of \mathbf{A} – $n \times n$
 - ▣ ***Matrix must be square***

Matrix Transpose

25

- The ***transpose*** of a matrix is that matrix with ***rows and columns swapped***
 - ▣ First row becomes the first column, second row becomes the second column, and so on
- For example:

$$\mathbf{A} = \begin{bmatrix} 0 & 9 \\ 2 & 7 \\ 6 & 3 \end{bmatrix} \quad \mathbf{A}^T = \begin{bmatrix} 0 & 2 & 6 \\ 9 & 7 & 3 \end{bmatrix}$$

- Row vectors become column vectors and vice versa

$$\mathbf{x} = \begin{bmatrix} 7 \\ -1 \\ -4 \end{bmatrix} \quad \mathbf{x}^T = [7 \quad -1 \quad -4]$$

Why Do We Use Matrices?

26

- Vectors and matrices are used extensively in many engineering fields, for example:
 - ▣ Modeling, analysis, and design of dynamic systems
 - ▣ Controls engineering
 - ▣ Image processing
 - ▣ Etc. ...
- Very common usage of vectors and matrices is to represent ***systems of equations***
 - ▣ These regularly occur in *all* fields of engineering

Systems of Equations

27

- Consider a system of three equations with three unknowns:

$$\begin{aligned}3x_1 + 5x_2 - 9x_3 &= 6 \\ -3x_1 + 7x_3 &= -2 \\ -x_2 + 4x_3 &= 8\end{aligned}$$

- Can represent this in **matrix form**:

$$\begin{bmatrix} 3 & 5 & -9 \\ -3 & 0 & 7 \\ 0 & -1 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 6 \\ -2 \\ 8 \end{bmatrix}$$

- Or, more compactly as:

$$\mathbf{Ax} = \mathbf{b}$$

- Perform algebra operations as we would if \mathbf{A} , \mathbf{x} , and \mathbf{b} were scalars
 - Observing matrix-specific rules, e.g. multiplication order, etc.