

Introduction to Categorical Data Analysis



DATA SCIENCE BOOTCAMP

Categorical Data Analysis

Methods for response (y) variable having scale that is categorical


Examples:

- Binary (Democrat or Republican)
(Success or Not Success)
- Nominal (Favorite music genre: rock, classical, jazz)
- Ordinal (Rank: 1st, 2nd, 3rd)

For these problems, linear regression is not the correct approach.

Probability Distributions for Categorical Data


In linear regression: response y is continuous, and the (conditional) response $y | x$ is normally distributed.



Probability Distributions for Categorical Data

In linear regression: response y is continuous, and the (conditional) response $y | x$ is normally distributed.

In models where y is categorical, the response $y | x$ can take other distributions.



Let's first start simple.

Binomial Experiments

The experiment of tossing a coin n times is an example of a binomial experiment.

Binomial Experiments

The experiment of tossing a coin n times is an example of a binomial experiment.

Three conditions:

- n independent trials
- The outcome of a single trial is classified as 1 of 2 types:
 - *Success*
 - *Not Success*
- For each trial, the probability of success is the same:
 - $P(\text{Success}) = \pi$
 - Note: π is called the parameter of the experiment.

Binomial Experiments

Y : random variable, represents the # of successes out of n trials

Example:

- 200 voters
- Each person votes:
 - Democrat – with $p = 0.55$
 - Republican – with $p = 0.45$

Binomial Experiments

Y : random variable, represents the # of successes out of n trials

Example:

- 200 voters
- Each person votes:
 - Democrat – with $p = 0.55$
 - Republican – with $p = 0.45$
- Then # Democrat votes $\sim \text{Binomial} (n=200, \pi = 0.55)$

Binomial Experiments

Y : random variable, represents the # of successes out of n trials

Example:


- 200 voters
- Each person votes:
 - Democrat – with $p = 0.55$
 - Republican – with $p = 0.45$
- Then # Democrat votes $\sim \text{Binomial} (n=200, \pi = 0.55)$
- Note: if $n = 1$ (just one trial), then $Y \sim \text{Bernoulli}$ distribution

Binomial Experiments

Y : random variable, represents the # of successes out of n trials

Example:

- 200 voters
- Each person votes:
 - Democrat – with $p = 0.55$
 - Republican – with $p = 0.45$
- Then # Democrat votes $\sim \text{Binomial} (n=200, \pi = 0.55)$
- Note: if $n = 1$ (just one trial), then $Y \sim \text{Bernoulli}$ distribution



The Binomial Distribution

More generally, the random variable

$$Y \sim \textit{Binomial}(n, p)$$

has probability mass function:

$$P(Y = y) = \frac{n!}{y!(n-y)!} \pi^y (1 - \pi)^{n-y}$$

The Binomial Distribution

More generally, the random variable

$$Y \sim \text{Binomial}(n, \pi)$$

has probability mass function:

$$P(Y = y) = \frac{n!}{y!(n-y)!} \pi^y (1 - \pi)^{n-y}$$

In general, for a binomial random variable Y ,

- The Expected Value is:

$$E(Y) = \mu_Y = n\pi$$

- The Variance is:

$$\text{Var}(Y) = \sigma_Y^2 = n\pi(1-\pi)$$

The Binomial Distribution

More generally, the random variable

$$Y \sim \text{Binomial}(n, \pi)$$

has probability mass function:

$$P(Y = y) = \binom{n}{y} \pi^y (1 - \pi)^{n-y}$$

In general, for a binomial random variable Y ,


- The Expected Value is:

$$E(Y) = \mu_Y = n\pi$$

- The Variance is:

$$\text{Var}(Y) = \sigma_Y^2 = n\pi(1-\pi)$$

The Binomial Distribution



Example of a Binomial Distribution

Example of a Binomial Distribution

Let $Y = \#$ of people who voted Democrat.

$n = 3$ voters.

Suppose $P(\text{vote Dem}) = 0.5$

Possible values of Y are:

Example of a Binomial Distribution

Let $Y = \#$ of people who voted Democrat.

$n = 3$ voters.

Suppose $P(\text{vote Dem}) = 0.5$

Possible values of Y are: $0, 1, 2, 3$

Example of a Binomial Distribution

Let $Y = \#$ of people who voted Democrat.
 $n = 3$ voters.

Suppose $P(\text{vote Dem}) = 0.5$

Possible values of Y are: $0, 1, 2, 3$

Then Y follows a binomial distribution:

$$Y \sim \text{Binomial}(n = 3, \pi = 0.5)$$

Example of a Binomial Distribution

Let $Y = \#$ of people who voted Democrat.


$n = 3$ voters.

Suppose $P(\text{vote Dem}) = 0.5$

Possible values of Y are: $0, 1, 2, 3$

Then Y follows a binomial distribution:

$$Y \sim \text{Binomial}(n = 3, \pi = 0.5)$$



Example of a Binomial Distribution

Let $Y = \#$ of people who voted Democrat.

$n = 3$ voters.

Suppose $P(\text{vote Dem}) = 0.5$


Possible values of Y are: 0, 1, 2, 3

Then Y follows a binomial distribution:

$Y \sim \text{Binomial}(n = 3, \pi = 0.5)$

$$P(Y = y) = \frac{3!}{y!(3-y)!} (1/2)^y (1/2)^{3-y}$$

$$= \frac{3!}{y!(3-y)!} (1/8)$$



Thinking with Likelihoods



Thinking with Likelihoods

Before, we were thinking about the probability of various possible outcomes of Y , given a parameter π (and n data points).

Given π , what is the probability of seeing y # *successes*, $P(Y=y)$?

Given $P(\text{vote Dem}) = 0.5$, what is the probability of seeing y # *Democrat votes*, $P(Y=y)$?

Thinking with Likelihoods

Before, we were thinking about the probability of various possible outcomes of Y , given a parameter π .

Given π , what is the probability of seeing y # *successes*, $P(Y=y)$?

In real life, we're often more interested in going the other way. So now let's flip our thinking:

Given $Y = y$ # *successes*, what is the underlying proportion π ?

Thinking with Likelihoods

Before, we were thinking about the probability of various possible outcomes of Y , given a parameter π .

Given π , what is the probability of seeing y # *successes*, $P(Y=y)$?

In real life, we're often more interested in going the other way. So now let's flip our thinking:

Given $Y = y$ # *successes*, what is the underlying proportion π ?

Example: on Wall St, I count 100 people, 15 of whom are women. What is the proportion of women on Wall St?

Thinking with Likelihoods: The Idea

Thinking with Likelihoods: The Idea

- There is an unknown parameter π (like a “state of nature”)
- Have data, i.e. observations Y (or Y_1, Y_2, Y_3, \dots)
- Assume a model, i.e. $Y \sim \text{Binomial}(n, \pi)$

Thinking with Likelihoods: The Idea

- There is an unknown parameter π (like a “state of nature”)
- Have data, i.e. observations Y (or Y_1, Y_2, Y_3, \dots)
- Assume a model, i.e. $Y \sim \text{Binomial}(n, \pi)$

Goal: want $P(\pi \mid Y = y)$

Thinking with Likelihoods: The Idea

- There is an unknown parameter π (like a “state of nature”)
- Have data, i.e. observations Y (or Y_1, Y_2, Y_3, \dots)
- Assume a model, i.e. $Y \sim \text{Binomial}(n, \pi)$

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

Thinking with Likelihoods: The Idea

- There is an unknown parameter π (like a “state of nature”)
- Have data, i.e. observations Y (or Y_1, Y_2, Y_3, \dots)
- Assume a model, i.e. $Y \sim \text{Binomial}(n, \pi)$

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

In particular, we'd like to find the probability that is *maximal*.

Thinking with Likelihoods: The Idea

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

Two ways:

Thinking with Likelihoods: The Idea

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

Two ways:

1. Bayes Theorem.

Thinking with Likelihoods: The Idea

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

Two ways:

1. Bayes Theorem.

$P(\pi \mid Y = y)$ is proportional to $P(\pi) P(Y = y \mid \pi)$

Thinking with Likelihoods: The Idea

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

Two ways:

1. Bayes Theorem.

$P(\pi \mid Y = y)$ is proportional to $P(\pi)$ $P(Y = y \mid \pi)$

prior likelihood



Thinking with Likelihoods: The Idea

Goal: want $P(\pi \mid Y = y)$

After we have data, we want to know the probability of seeing various values of π .

Two ways:

1. Bayes Theorem.
 $P(\pi \mid Y = y)$ is proportional to $\overset{\text{prior}}{\downarrow} P(\pi) \overset{\text{likelihood}}{\downarrow} P(Y = y \mid \pi)$

Some controversy in getting the prior:

- How to get it, what it means, subjective disagreements, subjective biases

Thinking with Likelihoods: The Idea

2. Maximum Likelihood

No priors; more limited, working only with the likelihood

Thinking with Likelihoods: The Idea

2. Maximum Likelihood

No priors; more limited, working only with the likelihood

We can conduct inferences about parameters using maximum likelihood.

Thinking with Likelihoods: The Idea

2. Maximum Likelihood

No priors; more limited, working only with the likelihood

We can conduct inferences about parameters using maximum likelihood.

Definition. The likelihood function is the probability of the observed data occurring, expressed as a function of the parameter value.

Thinking with Likelihoods: The Idea

2. Maximum Likelihood

No priors; more limited, working only with the likelihood

We can conduct inferences about parameters using maximum likelihood.

Definition. The likelihood function is the probability of the observed data occurring, expressed as a function of the parameter value.

Example.

Of the $N = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial}(n = 2, \pi = ?)$

Maximum Likelihood Example, continued.

Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial}(n = 2, \pi = ?)$

Maximum Likelihood Example, continued.

Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

Binomial Distribution:

$$P(Y = y \mid n=2) = \frac{2!}{y!(2-y)!} (\pi)^y (1-\pi)^{2-y}$$

Maximum Likelihood Example, continued.

Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

Binomial Distribution:

$$P(Y = y \mid n=2) = \frac{2!}{y!(2-y)!} (\pi)^y (1-\pi)^{2-y}$$

$$P(Y = 1 \mid n=2) = 2\pi(1-\pi)$$

Maximum Likelihood Example, continued.

Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

Binomial Distribution:

$$P(Y = y \mid n=2) = \frac{2!}{y!(2-y)!} (\pi)^y (1-\pi)^{2-y}$$

$$P(Y = 1 \mid n=2) = 2\pi(1-\pi)$$

The likelihood function, below, is the probability of observing $y=1$, expressed as a function of parameter value π :

$$L(\pi) = 2\pi(1-\pi)$$


Maximum Likelihood Example, continued.

Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial}(n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$




Maximum Likelihood Example, continued.

Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial}(n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$



Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$

If $\pi = 1$, the likelihood function represents the probability of observing $y = 1$.

Of the $n = 2$ people on Wall Street, 1 was a woman.


Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$

If $\pi = 1$, the likelihood function represents the probability of observing $y = 1$.

$$\begin{aligned} L(\pi = 1) &= 2 \times 1 \times (1 - 1) \\ &= 0 \end{aligned}$$



Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$

If $\pi = 0$, *the likelihood function represents the probability of observing $y = 1$.*

Of the $n = 2$ people on Wall Street, 1 was a woman.


Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$

If $\pi = 0$, the likelihood function represents the probability of observing $y = 1$.

$$\begin{aligned} L(\pi = 0) &= 2 \times 0 \times (1 - 0) \\ &= 0 \end{aligned}$$



Of the $n = 2$ people on Wall Street, 1 was a woman.

Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$

$$L(\pi) = 2\pi(1 - \pi)$$

If $\pi = 1/2$, the likelihood function represents the probability of observing $y = 1$.

Of the $n = 2$ people on Wall Street, 1 was a woman.


Then $Y = \# \text{ successes} = 1$.

$Y \sim \text{Binomial} (n = 2, \pi = ?)$


$$L(\pi) = 2\pi(1 - \pi)$$

If $\pi = 1/2$, the likelihood function represents the probability of observing $y = 1$.

$$\begin{aligned} L(\pi = 1/2) &= 2 \times 1/2 \times (1 - 1/2) \\ &= 1/2 \end{aligned}$$



This is the maximum likelihood estimate.




Definition. The Maximum Likelihood (ML) estimate of a parameter π is the parameter value at which the likelihood function takes its maximum.

Intuitively, the Maximum Likelihood estimate $\hat{\pi}$ is your guess for the “state of nature” that best explains the data. It is the estimate for which the data is *most likely* to occur.

Definition. The Maximum Likelihood (ML) estimate of a parameter π is the parameter value at which the likelihood function takes its maximum.

Intuitively, the Maximum Likelihood estimate $\hat{\pi}$ is your guess for the “state of nature” that best explains the data. It is the estimate for which the data is *most likely* to occur.

Example: $L(\pi) = 2\pi(1 - \pi)$ is maximized at $\pi = 1/2$.
Thus the Maximum Likelihood estimate for $L(\pi)$ is $\hat{\pi} = 1/2$.



Notes.

For Binomial random variable Y : $\hat{\pi} = y/n$

Mathematically, in Ordinary Least Squares ($Y|X \sim \text{Normal}$), the least squares estimates for the β parameters are the Maximum Likelihood estimates.