


# GENERALIZED LINEAR MODELS




# Motivation: Normal Linear Models



Let's start with some data...



# Motivation: Normal Linear Models




Let's start with some data...

Here's an attempt to fit the data. How do we judge it?



# Motivation: Normal Linear Models




Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$



# Motivation: Normal Linear Models



Let's start with some data...


Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost



# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?


Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost

We often plot residuals on the y-axis



# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost


We often plot residuals on the y-axis

Some things to notice:

- Equal-magnitude residuals make the same contribution to the cost
- A positive (or negative) residual means the data is greater than (or less than) we predicted



# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost

We often plot residuals on the y-axis

Some things to notice:


- Equal-magnitude residuals make the same contribution to the cost
- A positive (or negative) residual means the data is greater than (or less than) we predicted

Features are usually higher-dimensional, tough to plot





# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost

We often plot residuals on the y-axis

Some things to notice:


- Equal-magnitude residuals make the same contribution to the cost
- A positive (or negative) residual means the data is greater than (or less than) we predicted

Features are usually higher-dimensional, tough to plot

So you'll often see "predicted values" on the x-axis



# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost

We often plot residuals on the y-axis

Some things to notice:

- Equal-magnitude residuals make the same contribution to the cost
- A positive (or negative) residual means the data is greater than (or less than) we predicted


Features are usually higher-dimensional, tough to plot

So you'll often see "predicted values" on the x-axis

We can also check for patterns this way



# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost

We often plot residuals on the y-axis

Some things to notice:

- Equal-magnitude residuals make the same contribution to the cost
- A positive (or negative) residual means the data is greater than (or less than) we predicted

Features are usually higher-dimensional, tough to plot


So you'll often see "predicted values" on the x-axis

We can also check for patterns this way

Would an  $x^2$  feature improve the fit?



# Motivation: Normal Linear Models



Let's start with some data...

Here's an attempt to fit the data. How do we judge it?

Find the residuals,  $r = y - \hat{y}$

Each residual gives a contribution (its square) to the cost

We often plot residuals on the y-axis

Some things to notice:

- Equal-magnitude residuals make the same contribution to the cost
- A positive (or negative) residual means the data is greater than (or less than) we predicted

Features are usually higher-dimensional, tough to plot

So you'll often see "predicted values" on the x-axis


We can also check for patterns this way

Would an  $x^2$  feature improve the fit?

Looks like a yes!




# Can we Generalize?



Say we have some count data like this



# Can we Generalize?




Say we have some count data like this

To this point we've seen conditional expectations that are...

- Normal (linear regression)
- Bernoulli or binomial (logistic regression)



# Can we Generalize?



Say we have some count data like this

To this point we've seen conditional expectations that are...

- Normal (linear regression)
- Bernoulli or binomial (logistic regression)

It would be great to...

- Find something similar for count data
- Find the analog of residuals and squared residuals



# Poisson Regression: Just Another GLM

Here are three generalized linear models:

Linear regression	identity $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Normal}(X\beta, \sigma)$
Logistic regression	logit $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Bernoulli}(\text{logit}^{-1}(X\beta))$
Poisson regression	log $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Poisson}(\exp(X\beta))$





# Poisson Regression: Just Another GLM

Here are three generalized linear models:

Linear regression	identity $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Normal}(X\beta, \sigma)$
Logistic regression	logit $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Bernoulli}(\text{logit}^{-1}(X\beta))$
Poisson regression	log $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Poisson}(\exp(X\beta))$

$X\beta$  is the *systematic component*



# Poisson Regression: Just Another GLM

Here are three generalized linear models:

Linear regression	identity $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Normal}(X\beta, \sigma)$
Logistic regression	logit $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Bernoulli}(\text{logit}^{-1}(X\beta))$
Poisson regression	log $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Poisson}(\exp(X\beta))$

The distribution of  $y$  is the *stochastic component*



# Poisson Regression: Just Another GLM

Here are three generalized linear models:


Linear regression	identity $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Normal}(X\beta, \sigma)$
Logistic regression	logit $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Bernoulli}(\text{logit}^{-1}(X\beta))$
Poisson regression	log $\mathbb{E}(y   X) = X\beta$	$y \sim \text{Poisson}(\exp(X\beta))$



The *link function* connects the systematic and stochastic components




# Like Counting Fish in a Barrel



Back to our count data



# Like Counting Fish in a Barrel




Back to our count data

The Poisson regression model is

$$y \sim \text{Poisson}(\exp(X\beta))$$



# Like Counting Fish in a Barrel



Back to our count data


The Poisson regression model is

$$y \sim \text{Poisson}(\exp(X\beta))$$

Here's the conditional expectation



# Like Counting Fish in a Barrel



Back to our count data

The Poisson regression model is


$$y \sim \text{Poisson}(\exp(X\beta))$$

Here's the conditional expectation

What about residuals?



# These Aren't the Residuals You're Looking For




Why do we usually take “observed minus predicted”?





# These Aren't the Residuals You're Looking For




Why do we usually take “observed minus predicted”?

Likelihood for a normal looks like this:

$$L_{\text{Normal}} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - \mu)^2}{2\sigma^2}\right)$$



# These Aren't the Residuals You're Looking For



Why do we usually take “observed minus predicted”?

Likelihood for a normal looks like this:


$$L_{\text{Normal}} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - \mu)^2}{2\sigma^2}\right)$$

But here's a Poisson likelihood:

$$L_{\text{Poisson}} = \frac{\mu^y e^{-\mu}}{y!}$$



# These Aren't the Residuals You're Looking For



Why do we usually take “observed minus predicted”?

Likelihood for a normal looks like this:

$$L_{\text{Normal}} = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - \mu)^2}{2\sigma^2}\right)$$


But here's a Poisson likelihood:

$$L_{\text{Poisson}} = \frac{\mu^y e^{-\mu}}{y!}$$

“Residuals” will be different, but we can still calculate the contribution of each point to the overall cost



# Unit Deviance



Here are the contributions of each point to the total cost

For a linear model, a perfectly-fit point has zero contribution (squared residual)

Here, we subtract cost for a hypothetical “saturated model” to make this work (more on this soon)

Resulting *unit deviance* values play the role of squared residuals


Remember the linear case:

- Squared residuals give contribution to cost
- Sign indicates data above or below prediction

Can we make that work out here? Spoiler: yes!



# Deviance Residuals



Here's the result!

Use these *deviance residuals* just like you'd use residuals for a linear model

There can be some unavoidable patterns, like the “Hawaii” in the lower-left



# Linear Models: Some more details

- For a normal linear model, the likelihood and log-likelihood for each data point are

$$L = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - \hat{y})^2}{2\sigma^2}\right)$$

$$\ell = -\frac{1}{2} \log(2\pi\sigma^2) - \frac{(y - \hat{y})^2}{2\sigma^2}$$



# Linear Models: Some more details

- For a normal linear model, the likelihood and log-likelihood for each data point are

$$L = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - \hat{y})^2}{2\sigma^2}\right)$$

$$\ell = -\frac{1}{2} \log(2\pi\sigma^2) - \frac{(y - \hat{y})^2}{2\sigma^2}$$

- So the total log-likelihood is

$$\ell = -\frac{N}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{j=1}^N (y_j - \hat{y}_j)^2$$



# Linear Models: Some more details

- For a normal linear model, the likelihood and log-likelihood for each data point are

$$L = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y - \hat{y})^2}{2\sigma^2}\right)$$

$$\ell = -\frac{1}{2} \log(2\pi\sigma^2) - \frac{(y - \hat{y})^2}{2\sigma^2}$$

- So the total log-likelihood is

$$\ell = -\frac{N}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{j=1}^N (y_j - \hat{y}_j)^2$$

- We've seen the **sum of squared residuals**, but what happened to **the first term**?





# The Saturated Model, and Deviance

Imagine we could fit a model perfectly. Our log-likelihood

$$\ell = -\frac{N}{2} \log(2\pi\hat{\sigma}^2) - \frac{1}{2\hat{\sigma}^2} \sum_{j=1}^N (y_j - \hat{y}_j)^2$$

would become

$$\ell_S = -\frac{N}{2} \log(2\pi\hat{\sigma}^2)$$

*Deviance* measures how far we are from a hypothetical perfect fit:

$$D = 2(\ell_S - \ell) = \frac{1}{\hat{\sigma}^2} \sum_{j=1}^N (y_j - \hat{y}_j)^2 = \sum_{j=1}^N \left( \frac{y_j - \hat{y}_j}{\hat{\sigma}} \right)^2$$

For a linear model, deviance is the sum of *Studentized* residuals!



# GLM Hypothesis Testing

- For a linear model, deviance is

$$D = \frac{1}{\sigma^2} \sum_{j=1}^N (y_j - \hat{y}_j)^2$$

- If  $\mathcal{M}_0$  is a “submodel” of  $\mathcal{M}$  (so  $\mathcal{M}_0 \subset \mathcal{M}$ ), which will fit training data better?  $\mathcal{M}$
- What does this mean about the deviance?  $D_0 > D$
- How do we measure “how much better”  $D$  fits? **With a  $\chi^2$  test!**

$$D_0 - D \sim \chi_{\Delta p}^2$$

↑

“Degrees of freedom” is difference in number of parameters

