A

Wt (Sanger)
n = 409

From \ To	A	T	C	G
A	—	46 11.2%	31 7.6%	63 15.4%
T	39 9.5%	—	57 13.9%	23 5.6%
C	10 2.4%	40 9.8%	—	11 2.7%
G	48 11.7%	18 4.4%	23 5.6%	—

Wt (ONT)
n = 548

From \ To	A	T	C	G
A	—	49 8.9%	58 10.6%	91 16.6%
T	53 9.7%	—	93 17.0%	36 6.6%
C	18 3.3%	44 8.0%	—	17 3.1%
G	45 8.2%	17 3.1%	27 4.9%	—

0 2 4 6 8 10 12 14 16
% of unique mutations

B

Platform × type interaction (repeated measures): F(11,22) = 1.53, p = 0.190

Sanger (n = 409) ONT (n = 548)

adj. p >0.99 | adj. p >0.99 | adj. p >0.99 | adj. p >0.99 | adj. p 0.92 | adj. p 0.99 | adj. p 0.76 | adj. p 0.97 | adj. p 0.95 | adj. p 0.44 | adj. p >0.99 | adj. p >0.99

Points: each mouse, area ∝ its mutations.
Sanger Wt2: 6 mutations
(the smallest points)

% of unique mutations

A>T A>C A>G T>A T>C T>G C>A C>T C>G G>A G>T G>C

C

r = 0.90 [0.69–0.97]
slope = 1.00

T>C A>G A>C A>T G>A

ONT, % of unique mutations
Sanger, % of unique mutations

D

Sanger ONT

adj. p = 0.20
adj. p = 1.00

% of unique mutations
G:C transitions
G:C transversions

E

Sanger
n = 93
13% 43% 14% 13% 17%

ONT
n = 1,491
15% 50% 14% 17%

Mutations per sequence
0
1–3
4–10
11–20
20+