

AID-competent library
AID-null control library
0 calls (below axis break)
A
JH4 intron
3 Wt vs 3 Aicda−/− mice
B
N114 VH
3 vs 3 subclones
C
Sμ, LPS blasts
3 vs 3 libraries
D
Sμ, CH12
4 vs 3 libraries
E
Pim1, LPS blasts
3 vs 3 libraries
Unique variants per 10^5 bases
SHMtrace
Geneious
LoFreq
LoFreq -BAQ
LoFreq perm.
F
Case / control separation
JH4 141 44 2.2 40 38
N114 VH 85 62 10 13 34
Sμ LPS 66 16 10 14 8.6
Sμ CH12 94 26 6.5 9.7 7.0
Pim1 1.8 1.6 1.0 1.0 1.3
CH12 OT1 Wt 59 4.5 3.3 8.5 2.5
CH12 OT1 dKO 1.6 1.0 1.0 1.0 0.9
CH12 OT2 Wt 28 8.7 1.6 2.5 1.6
CH12 OT2 dKO 1.0 1.3 1.0 1.0 0.8
1 3 10 30 100
fold (geometric means)
G
Calls in AID-null controls
JH4 2 13 0 5 14
N114 VH 1 5 0 0 1
Sμ LPS 2 20 1 2 17
Sμ CH12 2 11 8 8 21
Pim1 4 11 0 1 40
CH12 OT1 0 44 0 0 41
CH12 OT2 0 11 0 0 50
0 20 40
calls, summed over 3 controls
H
Held-out control calibration
held-out tests
JH4, 529-bp window (724)
JH4, 1,141 bp (1,464)
N114 VH (553)
margin selected
Observed held-out tests with p < 0.01 (%)
1%: if calibrated
Margin on control error rates (×)