

Week 18

Questions

1. Which of the following is not a manifestation of hyperactivity/impulsivity:
 - a. Blurts out answers before question is finished
 - b. Unable to stay seated
 - c. Fidgets
 - d. Careless mistakes
 - e. Difficulty playing quietly
2. T/F: ADHD resolves upon adulthood
3. T/F: The DSM-5 criteria for learning disorders require a test of IQ
4. Which of the following is not a risk factor for learning disabilities:
 - a. Genetics
 - b. Perinatal maternal EtOH consumption
 - c. Presence of lisp
 - d. Low birthweight
5. T/F: Articulation disorders increase with age
6. Which of the following is true about language disorders:
 - a. Language disorders are more common in girls
 - b. Bilingual children with language disorders will preferentially develop one language only
 - c. 90% of late talkers will catch up on their own
 - d. Language disorders are highly heritable
7. Which of the following is true about ASD:
 - a. It is a degenerative disorder
 - b. Anyone with a diagnosis of ASD should have genetic testing with a clinical microarray
 - c. Patients often have a reduced brain size compared to the general population
 - d. 75% of those with ASD have Intellectual Disability
8. Treatment for aggression in ASD include:
 - a. Fluoxetine
 - b. Benzodiazepine
 - c. Risperidone
 - d. Amitriptyline
9. Which of the following is a risk factor for disruptive behaviour disorders:
 - a. Chronic physical illness
 - b. Female sex
 - c. High birth weight
 - d. Overactivity of amygdala
10. Which of the following is not a presentation of conduct disorder?
 - a. Lack of empathy
 - b. Guilt and remorse
 - c. Flat affect

- d. Theft
11. Which of the following is not in the same cluster of personality disorders?
- a. Borderline Personality Disorder
 - b. Obsessive-Compulsive Personality Disorder
 - c. Histrionic Personality Disorder
 - d. Antisocial Personality Disorder
12. What is the most common type of personality disorder?
- a. Antisocial Personality Disorder
 - b. Narcissistic Personality Disorder
 - c. Borderline Personality Disorder
 - d. Schizotypal Personality Disorder
 - e. Paranoid Personality Disorder
13. Which of the following situations meet the criteria A for PTSD?
- a. Failing an Exam
 - b. Almost losing your life from house fire
 - c. Having a big fight with your best friend
 - d. Being suspended from school
14. What are the first line pharmacological therapies used for PTSD?
- a. Fluoxetine, Paroxetine, Sertraline, Venlafaxine
 - b. Amitryptiline, Aripiprazole, Bupropion, Topiramate
 - c. Olanzapine, Fluvoxamine, Mirtazapine, Fluvoxamine
 - d. Fluoxetine, Paroxetine, Sertraline, Citalopram
15. How long do you need to show symptoms of PTSD to be qualified for a diagnosis?
- a. 1 week
 - b. 6 months
 - c. 12 weeks
 - d. 3 months
16. Which of the following disorders are dissociation symptoms concomitantly found in? (Can choose more than 1)
- a. Borderline Personality Disorder
 - b. PTSD
 - c. Panic Disorder
 - d. Narcissistic Personality Disorder
 - e. Autism Spectrum Disorder
17. Which scale is used to evaluate severity of Dissociation Disorder?

- a. PHQ-9
 - b. YBOCS
 - c. MMSE
 - d. Dissociative Experiences Scale II (DES)
18. Average non-clinical population score on DES is...
- a. ≥ 26
 - b. 30
 - c. 11
 - d. 40
19. Which of the following are not one of the risk factors for developing anxiety disorders?
- a. Family history of anxiety
 - b. Personal history of anxiety or mood disorders (i.e., MDD, ADHD)
 - c. Childhood stressful life events/trauma
 - d. Chronic medical illness
 - e. Being male
20. Choose medical conditions that can present as anxiety or panic disorders
- a. Hyper/hypothyroidism
 - b. Caffeine
 - c. Anemia
 - d. Vitamin B12 deficiency
 - e. COPD
21. What is the median age for onset of OCD?
- a. 7
 - b. 15
 - c. 25
 - d. 19
22. T/F: the difference between worry and obsession is that obsession is almost exclusively experienced just as a thought.
23. Choose a combination of SSRI side effects with its appropriate treatments.
- a. Sexual side effects: use benzodiazepine
 - b. Restlessness: use bupropion
 - c. Sleep disturbance: use trazodone
 - d. Indigestion: use pepcid
24. Which one of the following are an appropriate example for each type of exposure therapies for anxiety Disorders?

- a. Situational - hyperventilating to provoke trembling/shaking for panic disorder treatment
- b. Interoceptive - going back to the location where you were mugged, to treat PTSD
- c. Imaginal - running back and forth to simulate symptoms of panic disorder
- d. Virtual - exposure to computer simulation of a war you've fought in, to treat PTSD

Answers

- 1. D
- 2. F. ADHD persists into adulthood in about 60%
- 3. F, but LDAO does require one!
- 4. C
- 5. F
- 6. D
- 7. B
- 8. C
- 9. A
- 10. B
- 11. B
- 12. C
- 13. B
- 14. A
- 15. D
- 16. A,B,C
- 17. D
- 18. C
- 19. E
- 20. A,B,C,D,E
- 21. D (earlier onset in men)
- 22. False; worry is almost exclusively experienced as a thought while obsession is experienced as intrusive thoughts, images, or impulses.
- 23. C
- 24. D