

Chapter 27 : Another appeal to monotonicity.

In which we once again use that nice property.

We are given $f[0..M)$, $g[0..N)$ of int. We are told that f is ascending and g is descending. We are asked to construct a program to compute the number of pairs $f.i$ and $g.j$ whose sum exceeds 37.

$$r = \langle + i,j : 0 \leq i < M \wedge 0 \leq j < N : h.(f.i).(g.j) \rangle$$

where

$$* (0) h.x.y = 1 \iff x + y > 37$$

$$* (1) h.x.y = 0 \iff x + y \leq 37$$

We begin by modelling our domain.

$$* (2) C.m.n = \langle + i,j : m \leq i < M \wedge n \leq j < N : h.(f.i).(g.j) \rangle$$

Emptying the components of the range in turn lead us to the following

$$- (3) C.M.n = 0, 0 \leq n \leq N$$

$$- (4) C.m.N = 0, 0 \leq m \leq M$$

We observe,

$$\begin{aligned} & C.m.n \\ = & \{(2)\} \\ & \langle + i,j : m \leq i < M \wedge n \leq j < N : h.(f.i).(g.j) \rangle \\ = & \{ \text{Split off } i = m \text{ term} \} \\ & \langle + i,j : m+1 \leq i < M \wedge n \leq j < N : h.(f.i).(g.j) \rangle + \langle + j : n \leq j < N : h.(f.m).(g.j) \rangle \\ = & \{(2), (7)\} \\ & C.(m+1).n + D.n \end{aligned}$$

$$- (5) C.m.n = C.(m+1).n + D.n, 0 \leq m < M$$

We observe,

$$\begin{aligned} & C.m.n \\ = & \{(2)\} \\ & \langle + i,j : m \leq i < M \wedge n \leq j < N : h.(f.i).(g.j) \rangle \\ = & \{ \text{Split off } j = n \text{ term} \} \\ & \langle + i,j : m \leq i < M \wedge n+1 \leq j < N : h.(f.i).(g.j) \rangle + \langle + i : m \leq i < M : h.(f.i).(g.n) \rangle \\ = & \{(2), (8)\} \end{aligned}$$

$$C.m.(n+1) + E.m$$

$$- (6) C.m.n = C.m.(n+1) + E.m, 0 < n \leq N$$

$$* (7) D.n = \langle + j : n \leq j < N : h.(f.m).(g.j) \rangle$$

$$-*(8) E.m = \langle + i : m \leq i < M : h.(f.i).(g.n) \rangle$$

We now turn our attention to investigating D and E

$$- (9) D.n = ? \Leftrightarrow f.m + g.n > 37$$

$$- (10) D.n = 0 \Leftrightarrow f.m + g.n \leq 37$$

$$- (11) E.m = M-m \Leftrightarrow f.m + g.n > 37$$

$$- (12) E.m = ? \Leftrightarrow f.m + g.n \leq 37$$

Our postcondition can now be written as

$$\text{Post} : r = C.0.0$$

Invariants.

As invariants we choose

$$P0 : r + C.m.n = C.0.0$$

$$P1 : 0 \leq m \leq M \wedge 0 \leq n \leq N$$

Establish Invariants.

$$R, m, n := 0, 0, 0$$

Upon termination

$$P0 \wedge P1 \wedge (m=M \vee n=N) \Rightarrow \text{Post}$$

Guard.

$$m \neq M \wedge n \neq N$$

Loop body

$$\begin{aligned}
& P0 \\
= & \quad \{\text{definition of } P0\} \\
& r + C.m.n = C.0.0 \\
= & \quad \{(5)\} \\
& r + C.(m+1).n + D.n = C.0.0 \\
= & \quad \{\text{case analysis } f.m + g.n \leq 37 \text{ (10)}\} \\
& r + C.(m+1).n + 0 = C.0.0 \\
= & \quad \{WP\} \\
& (r, m := r+0, m+1).P0
\end{aligned}$$

Giving us

$$\text{if } f.m + g.n \leq 37 \rightarrow r, m := r+0, m+1$$

We also observe

$$\begin{aligned}
& P0 \\
= & \quad \{\text{definition of } P0\} \\
& r + C.m.n = C.0.0 \\
= & \quad \{(11)\} \\
& r + C.m.(n+1) + E.m = C.0.0 \\
= & \quad \{\text{case analysis } f.m + g.n > 37 \text{ (1)}\} \\
& r + C.m.(n+1) + M-m = C.0.0 \\
= & \quad \{WP\} \\
& (r, n := r+(M-m), n+1).P0
\end{aligned}$$

Giving us

$$\text{if } f.m + g.n > 37 \rightarrow r, n := r+(M-m), n+1$$

There is no point in appealing to (12) so we ignore that case

Finished Algorithm.

$$\begin{aligned}
& r, n, m := 0, 0, 0 \{P0 \wedge P1\} \\
& ; \text{ do } m \neq M \wedge n \neq N \rightarrow \{P0 \wedge P1 \wedge m \neq M \wedge n \neq N\} \\
& \quad \text{if } f.m + g.n \leq 37 \rightarrow r, m := r+0, m+1 \\
& \quad [] f.m + g.n > 37 \rightarrow r, n := r+(M-m), n+1 \\
& \quad \text{fi} \\
& \quad \{P0 \wedge P1\} \\
& \text{od} \\
& \{P0 \wedge P1 \wedge (m=M \vee n=N)\}
\end{aligned}$$

