

Lecture 6

(Motion with Constant Acceleration, Free Fall)

Physics 160-01 Fall 2012

Douglas Fields

Derivatives


$$\frac{d}{dt} t^n = n t^{n-1}$$

$$\frac{d}{d\theta} \cos \theta = -\sin \theta$$

$$\frac{d}{d\theta} \sin \theta = \cos \theta$$


CPS Question 6-1

- At which point in the graph below is the velocity negative and acceleration positive?



CPS Question 6-2


- In the position versus time graph shown below, which answer best describes the instantaneous velocity and acceleration at time t_1 ?



- A. velocity is positive, acceleration is zero
- B. velocity is positive, acceleration is positive
- C. velocity is negative, acceleration is negative
- D. velocity is negative, acceleration is zero
- E. velocity is zero, acceleration is zero


Motion With Constant Acceleration

- For many applications, we can consider the acceleration to be constant.
- Acceleration of a falling object near the earth's surface \approx constant (if you ignore air resistance).



Motion With Constant Acceleration

- Now, from our discussions last class, what type of velocity dependence with time gives a constant acceleration?
- Linear behavior: $v_y(t) = a_y t$ $v_y(t) = v_{0y} + a_y t$
- But information is missing, the same slope gives the correct acceleration regardless of where the line starts.



Velocity From Acceleration

- From calculus, we can also get this result by using anti-differentiation (integration):

$$\begin{aligned}a_y(t) &= \frac{dv_y}{dt} \Rightarrow \\a_y(t) dt &= dv_y \Rightarrow \\\int a_y(t) dt &= \int dv_y\end{aligned}$$

- And if a_y is a constant,

$$\begin{aligned}\int a_y(t) dt &= a_y \int dt = \int dv_y \Rightarrow \\a_y t + C &= v_{fy} + C \Rightarrow \\v_{fy} &= v_{oy} + a_y t\end{aligned}$$

Position From Velocity

- We can do this again because we know that the position function is also related to the velocity function:

$$v_y(t) = \frac{dy}{dt} \Rightarrow$$

$$v_y(t) dt = dy \Rightarrow$$

$$\int v_y(t) dt = \int dy \Rightarrow$$

$$\int (a_y t + v_{0y}) dt = \int dy \Rightarrow$$

$$a_y \int t dt + v_{0y} \int dt = \int dy \Rightarrow$$

$$\frac{1}{2} a_y t^2 + v_{0y} t = y(t) + y_0 \Rightarrow$$

$$y_f = y_0 + v_{0y} t + \frac{1}{2} a_y t^2$$

Equations of Motion (constant a)

- We then have the following equations of motion:

$$y_f = y_0 + v_{0y}t + \frac{1}{2}a_y t^2$$

$$v_{fy} = v_{0y} + a_y t$$

- From these, we can solve for t in the second, and substitute into the first:

$$v_y = v_{0y} + a_y t \Rightarrow t = \frac{v_y - v_{0y}}{a_y} \Rightarrow$$

$$y_f = y_0 + v_{0y} \left(\frac{v_y - v_{0y}}{a_y} \right) + \frac{1}{2} a_y \left(\frac{v_y - v_{0y}}{a_y} \right)^2$$

$$2a_y (y_f - y_0) = (v_y - v_{0y})^2 + 2v_{0y} (v_y - v_{0y}) = v_y^2 - 2v_y v_{0y} + v_{0y}^2 + 2v_y v_{0y} - 2v_{0y}^2 \Rightarrow$$

$$v_y^2 - v_{0y}^2 = 2a_y (y_f - y_0)$$

Equations of Motion (constant a)

- We then have the following equations of motion:

$$y_f = y_0 + v_{0y}t + \frac{1}{2}a_y t^2$$

$$v_{fy} = v_{0y} + a_y t$$

$$v_{fy}^2 - v_{0y}^2 = 2a_y (y_f - y_0)$$

$$y_f - y_0 = \left(\frac{v_{fy} - v_{0y}}{2} \right) t$$

- Remember:
 - the “y” is just a name for a direction, I could have used x (like the book) or Nancy...

Example

- A car starts from rest and accelerates with constant acceleration = 2m/s^2 for 5s. What is its velocity after this time?
- What is its position after at $t=4\text{s}$?

CPS 6-3

- A car starts from rest and accelerates with constant acceleration = 4m/s^2 through a distance of 100m. How much time does it take the car to reach this distance?
- A. 25s
- B. 50s
- C. 7s
- D. 10s

Free Fall

- Near the earth's surface, objects fall with nearly constant acceleration of 9.8m/s^2 pointing down (due to the force of gravity).
- Objects that have significant air resistance compared to the force of gravity (feathers, parachutes) fall with lower acceleration.
- If you take away the air...

Free Fall in Vacuum

- Demonstration

Timed Free Fall



- Demonstration

CPS Demonstration Question

Timed Free Fall

- **The timer starts when the ball is dropped, and ends when it hits a plate 2m below. How much time does it take?**
- A) 0.41s
- B) 1.5s
- C) 0.64s
- D) 2.2s
- E) Cannot determine, insufficient information.

Two Objects



$$x_M = 0 + v_{M0x}t + \frac{1}{2}(0)t^2 = v_{M0x}t$$

$$x_P = 0 + (0)t + \frac{1}{2}a_{Px}t^2 = \frac{1}{2}a_{Px}t^2$$

$$v_{M0x}t = \frac{1}{2}a_{Px}t^2$$


$$t = 0 \quad \text{or} \quad t = \frac{2v_{M0x}}{a_{Px}} = \frac{2(15 \text{ m/s})}{3.0 \text{ m/s}^2} = 10 \text{ s}$$

$$v_{Px} = v_{P0x} + a_{Px}t = 0 + (3.0 \text{ m/s}^2)t$$

$$x_M = v_{M0x}t = (15 \text{ m/s})(10 \text{ s}) = 150 \text{ m}$$

Problem 2.66

- The engineer of a passenger train traveling at 25.0 m/s sights a freight train whose caboose is 200 m ahead on the same track. The freight train is traveling at 15.0 m/s in the same direction as the passenger train. The engineer of the passenger train immediately applies the brakes, causing a constant acceleration of -0.100 m/s^2 , while the freight train continues with constant speed. Take $x=0$ at the location of the front of the passenger train when the engineer applies the brakes. Is there a collision? If so, where?



Problem 2.83

- Sam heaves a shot with weight 16-lb straight upward, giving it a constant upward acceleration from rest of 46.0m/s^2 for a height 62.0cm. He releases it at height 2.17m above the ground. You may ignore air resistance.
- What is the speed of the shot when he releases it?
- How high above the ground does it go?
- How much time does he have to get out of its way before it returns to the height of the top of his head, a distance 1.84m above the ground?