



Program: ERRAT2
File: 2ZJQ.pdb
Chain#:A
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:B
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:C

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:D


Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:E
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:F

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:G


Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:H
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:I

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:J


Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:K
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:L


Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:M
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:N

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:O

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:P

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:Q

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:R


Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.


Program: ERRAT2
File: 2ZJQ.pdb
Chain#:S
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.


Program: ERRAT2
File: 2ZJQ.pdb
Chain#:T
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:U
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:V

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:W

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:Z

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:1

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:2

Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.


**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2

File: 2ZJQ.pdb

Chain#:3


Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.


Program: ERRAT2
File: 2ZJQ.pdb
Chain#:4
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.

Program: ERRAT2
File: 2ZJQ.pdb
Chain#:5
Overall quality factor**: 51.408



*On the error axis, two lines are drawn to indicate the confidence with which it is possible to reject regions that exceed that error value.

**Expressed as the percentage of the protein for which the calculated error value falls below the 95% rejection limit. Good high resolution structures generally produce values around 95% or higher. For lower resolutions (2.5 to 3Å) the average overall quality factor is around 91%.